

to the age group 8 to 18 years. One hundred one children underwent surgical fixation of cervical spine under anesthesia. Intravenous anesthetic agents were used for induction, and inhalational anesthetics were used for the maintenance of anesthesia. In majority of the children (62.4%), the airway was secured with fiberoptic intubation. Children in the age group 0 to 8 years (56.3%) had better outcome as compared with those in 8 to 18 years age group (39.6%). Fall from height constituted the major cause of injury ( $n = 68$ , 69.6%), and majority of the children suffered injury at the upper cervical spine (58.9%). Spinal canal was compromised in 77.4% of children and these children had poor outcome. Various complications were observed in 45 (40.2%) children during their hospital stay and 37 (82.2%) of these children had poor outcome. Nine children died owing to different reasons.

**Conclusion:** The overall outcome was good in 42% of children with CSI. The outcome at discharge was poor in children belonging to the age group of 8 to 18 years, who had compromised spinal canal, ASIA scale (A, B, or C) at admission, and those who suffered complications during the hospital stay.

**Keywords:** fiberoptic intubation, CSI, anesthetic

#### References

1. Leonard JC, Jaffe DM, Olsen CS, Kuppermann N. Age-related differences in factors associated with cervical spine injuries in children. *Acad Emerg Med* 2015;22(4):441–446

#### A012 Remote Ischemic Preconditioning in Decreasing Post-Electroconvulsive Therapy-Induced Cognitive Dysfunction

Ravitej Ravitej,<sup>1</sup> V. J. Ramesh,<sup>1</sup> Jagadish Thirthahalli,<sup>1</sup> K. N. Gopalkrishna,<sup>1</sup> Naveen Kumar<sup>1</sup>

<sup>1</sup>Department of Neuroanaesthesia, NIMHANS, Bangalore, Karnataka, India

**Introduction:** Ischemic injury has been implicated in causing post-electroconvulsive therapy (ECT) cognitive dysfunction. Remote ischemic preconditioning (RIPC) causes nonlethal ischemia on a limb, which results in protection of distant organ from ischemic injury. Hence, we studied the role of RIPC in decreasing post-ECT cognitive dysfunction.

**Methodology/Description:** The study was conducted on 80 patients after obtaining their informed consent and ethics committee approval. Patients 18 to 65 years of age, of either gender with a diagnosis of schizophrenia and ASA 1 to 2, were included. The patients were randomized into two groups. The intervention group received RIPC and the control group received sham RIPC before each ECT session for six such ECTs. The RIPC protocol included three 5-minute cycles of upper limb ischemia inflated to a cuff pressure of 30 mm Hg above the systolic BP with 5 minutes intervals of reperfusion in between. The cuff in the control group was inflated to a pressure of 30 mm Hg only. The cognitive assessment was done before the first and after the sixth ECT. Mann-Whitney U test, ANOVA, and paired *t*-test were used for analysis of data.

**Results:** There was no significant statistical difference in the cognitive and memory assessment scores of HMSE test ( $p = 0.908$ ), PGI memory scale test ( $p = 0.123$ ), B4ECT ReCODE test ( $p = 0.308$ ) both within and in between the two groups.

**Conclusion:** RIPC has not been found effective in preventing cognitive and memory impairment post-ECT in schizophrenia patients.

**Keywords:** RIPC, ECT, ASA

#### References

1. Hausenloy DJ, Yellon DM. Remote ischaemic preconditioning: underlying mechanisms and clinical application. *Cardiovasc Res* 2008;79(3):377–386
2. Rao SK, Andrade C, Reddy K, Madappa KN, Thyagarajan S, Chandra S. Memory protective effect of indomethacin against electroconvulsive shock-induced retrograde amnesia in rats. *Biol Psychiatry* 2002;51(9):770–773

#### A013 Clinical Efficacy of Scalp Infiltration with 0.25% Levobupivacaine to Attenuate Hemodynamic Response to Skull Pin Insertion in Supratentorial Craniotomies under General Anesthesia

Paulomi Dey,<sup>1</sup> Veena Ganeriwal<sup>1</sup>

<sup>1</sup>Department of Anesthesiology, Grant Government Medical College and Sir J. J. Hospital, Maharashtra, India

**Introduction:** Sudden hemodynamic changes caused by skull pins in neurosurgery can be prevented by combining scalp block, which blocks both the superficial and deep layers of the scalp. Levobupivacaine has been used in 0.75% and 0.5% concentrations for this purpose. However, no previous study has determined the efficacy of 0.25% levobupivacaine for scalp block.

**Aims:** To determine the effects of scalp block with 0.25% levobupivacaine on the hemodynamic response to head pinning and incision during craniotomy and evaluate the analgesic requirements intraoperatively.

**Methodology/Description:** This prospective observational study included 50 patients, after obtaining approval from ethics committee of Grant Government Medical College and Sir J.J. Hospital and written informed consent from patients (ASA status, I or II; age, 18 to 60 years) who underwent supratentorial craniotomy at this institute. Patients with coagulation disorders, inflammatory skin lesions, pre-existing neuropathies, pregnant and lactating mothers were excluded. Scalp block with 20 mL of 0.25% levobupivacaine was given bilaterally after induction of general anesthesia and prior to application of Mayfield skull pin head holder. The hemodynamic parameters were recorded at baseline, during scalp block and after head pin insertion, incision, and at craniotomy.

**Results:** Values of heart rate and blood pressure before and after anesthesia induction compared with values after painful stimuli were in the 10 to 15% range. Mean intraoperative fentanyl requirement was 150  $\mu$ g (2–3  $\mu$ g/kg).

**Conclusion:** Levobupivacaine (0.25%) when used for scalp block ensures hemodynamic fluctuation of upto