15% during head pinning, incision, and craniotomy without any complications or increased analgesic requirements intraoperatively.

Keywords: craniotomy, levobupivacaine, hemodynamic parameters

References

1. Can BO, Bilgin H. Effects of scalp block with bupivacaine versus levobupivacaine on haemodynamic response to head pinning and comparative efficacies in postoperative analgesia: a randomized controlled trial. *J Int Med Res* 2017;45(2):439–450
2. Bracho Pardey FG, Alne R, Barcelos G, et al. Levobupivacaine scalp nerve block: haemodynamics and anaesthesia requirements in supratentorial craniotomy: P 096. *Eur J Anaesthesiol* 2008;25:33–34

A014 Comparison of 0.375% Levobupivacaine, 0.375% Bupivacaine, and 0.375% Ropivacaine in Terms of Analgesia and Hemodynamic Stability following Scalp Block in Patients Undergoing Awake Craniotomy: A Prospective Randomized Double Blind Study Hitesh Nathani,¹ Hetal Rathod,¹ Bhoomika Thakore,¹ Joseph Monteiro¹

¹Department of Anaesthesiology, P. D. Hinduja Hospital and M. R. C., Mumbai, Maharashtra, India

Introduction: The anesthetic challenges of awake craniotomy are to maintain adequate sedation, analgesia, and hemodynamic stability in an awake patient who should be able to cooperate during intraoperative neurological assessment. Conventionally, scalp block is performed with racemic bupivacaine with epinephrine. Studies on bupivacaine isomers have shown reduced cardiovascular toxicity of its levorotatory form (levobupivacaine). Ropivacaine, the S(–) enantiomer of bupivacaine analogue, is known to have lesser cardiotoxicity with similar pain relief at equivalent analgesic doses in comparison to bupivacaine. This study was undertaken to compare levobupivacaine, bupivacaine, and ropivacaine for analgesia and hemodynamic stability and complications following scalp block in awake craniotomy patients.

Methodology/Description: The study was conducted at the Department of Neuroanesthesiology, P D Hinduja Hospital and M.R.C, Mumbai, after obtaining approval from the Institutional Review Board. Sample size of 42 patients (14 + 14 + 14) was determined using “MedCalc” and patients were randomized into one of the three arms based on computer-generated random numbers. Analgesia and hemodynamic parameters were noted at various time intervals. Results were analyzed by the Kruskal–Wallis test.

Results: All three groups were comparable in terms of demographic characteristics. There was no statistically significant difference in the VAS scores and hemodynamics at Application of Mayfield head pin ($p = 0.54$), skin incision ($p = 0.98$), craniotomy ($p = 0.299$), raising of bone flap ($p = 0.47$), opening of dura ($p = 0.50$), replacing of bone flap ($p = 0.14$), and skin closure ($p = 0.41$) in all three groups.

Conclusion: Levobupivacaine and ropivacaine provided equally good analgesia, hemodynamic parameters and

operating conditions as racemic bupivacaine and can be used routinely in our armamentarium of drugs for scalp block in awake craniotomy with better safety profiles.

Keywords: ropivacaine, levobupivacaine, awake craniotomy

References

1. Sokhal N, Rath GP, Chaturvedi A, Dash HH, Bithal PK, Chandra PS. Anaesthesia for awake craniotomy: a retrospective study of 54 cases. *Indian J Anaesth* 2015;59(5):300–305
2. Sengupta S, Mukherji S, Sheet J, Mandal A, Swaika S. Caudal-epidural bupivacaine versus ropivacaine with fentanyl for paediatric postoperative analgesia. *Anesth Essays Res* 2015; 9(2):208–212

A015 Optic Nerve Sheath Diameter Measured by Ultrasonography: How Well Does It Correlate with Intracranial Pressure According to the New Brain Trauma Foundation Guidelines?

Seelora Sahu,¹ Amlan Swain,¹ Nidhi Panda,² Hemant Bhagat,² Preethy Matthews,² Sunil Kumar Gupta²

¹Department of Anaesthesia and Critical Care, Tata Main Hospital, Jamshedpur, Jharkhand, India

²Department of Anaesthesia and Intensive Care, Post Graduate Institute of Medical Education and Research, Chandigarh, India

Introduction: Bedside ultrasonography (USG), measurement of optic nerve sheath diameter (ONSD) has been proposed as a method to detect raised intracranial pressure (ICP) in various clinical settings. We aimed to evaluate the use of USG in the case of intracranial hypertension and find out the cutoff point that predicts ICP accurately at 22 mm Hg.

Methodology/Description: A retrospective analysis of the data collected from a prospective double-blind study performed by performing ocular ultrasounds in 52 adult patients with features of intracranial hypertension was done. The ONSD was measured by USG under anesthesia and compared with the intraventricular ICP measured simultaneously. The optimum cutoff of ONSD to predict ICP > 22 mm Hg was sought.

Results: There was a significant correlation of sonographic ONSD with ICP ($r = 0.498$, $p = 0.000173$). An ONSD threshold of 6.3 mm predicted ICP > 22 mm Hg with high sensitivity (89%) and specificity (73%, area under ROC curve = 0.814, $p = 0.000424$, CI = 0.672–0.956).

Conclusion: Our study confirms the utility of optic nerve ultrasound in the diagnostic evaluation of patients with known or suspected intracranial hypertension. We recommend an ONSD cutoff of 6.3 mm for predicting ICP > 22 mm Hg.

Keywords: ONSD, ICP, ultrasound

References

1. Dubourg J, Javouhey E, Geeraerts T, Messerer M, Kassai B. Ultrasonography of optic nerve sheath diameter for detection of raised intracranial pressure: a systematic review and meta-analysis. *Intensive Care Med* 2011;37(7):1059–1068