20 patients 11 were ASA I, 3 ASA II, 5 ASA III, and 1 ASA IV grade. 7, 6, 3, 4 of 20 patients had blood loss < 500 ml, 500-1000 ml, 1000-1500 ml, and >1500 ml respectively. All patients received Tranexemic acid as prophylaxis for hemorrhage in a dose of 10 mg/kg bolus followed by 3 mg/kg infusion. Two patients were not given due to cardiac issues and had more blood loss than their counterparts. All patients were shifted to SICU for 24 hours. There were 7 patients with <500 ml blood loss were managed by fluid resuscitation, 6 patients with 500-100ml were managed by fluids alone and if Hb < 10 were given 1PRBC, 3 patients with 1000-1500 ml loss were given 3PRBC if Hb < 10; 2PRBC if Hb > 10 and remaining four patients with >1500 ml loss were managed by three or more PRBC plus three or more FFPs. **Conclusion:** To conclude prophylactic administration of Tranexemic acid is helpful in decreasing the amount of bleed; however, the bleed will depend on the size and duration of surgery. With the above mentioned protocol we can achieve better hemodynamics intra-operatively and better post op Hb.

24. Intraoperative electro-oculographic monitoring to prevent postoperative extra-ocular motor nerve dysfunction during skull base surgeries

Veena Sheshadri, B . Suparna

Department of Anaesthesia, Vikram Hospital, Bangalore, Karnataka, India

Background: To assess the feasibility of Electro-oculographic (EOG) monitoring in preventing extra-ocular motor nerve dysfunction during skull base surgeries. **Materials and Methods:** Intraoperative EOG recordings were obtained using disposable needle electrodes placed on the periorbital skin and the polarity of the waves noted for interpretation. The nerves were stimulated with a monopolar electrode during skull base exposure. Triggered as well as continuous EOG responses were recorded. **Results:** Of the nine cases monitored, oculomotor and abducent nerves were identified in all cases, but the trochlear nerve could not be definitively identified. Five out of nine patients had no pre or postoperative extra-ocular motor nerve dysfunction. The other four patients had pre-existing deficits before surgery, which recovered completely in two, significantly in one, and was unchanged in one patient at 3-6 month follow-up. **Conclusion:** EOG was found to be a simple and non-invasive method of monitoring extra-ocular motor nerves intra-operatively. Further studies in larger patient populations are required to quantitate the predictive value of EOG monitoring in detecting postoperative extra-ocular motor nerve deficits.

25. Pharmacological and non-pharmacological interventions for reducing pain on rocuronium injection - A systematic review

Bhavna Hooda, Hemanshu Prabhakar, Gyaninder Pal Singh, Zulfiqar Ali¹, Mani Kalaivani²

Department of Neuroanaesthesiology, ²Biostatistics, AIIMS, ¹Anaesthesia and Intensive Care, Govt. Medical College, Srinagar, India

Background: Various strategies, both pharmacological and non-pharmacological, have been studied to reduce the discomfort, incidence and severity of pain on rocuronium bromide injection. The objective of this review was to assess the ability of both pharmacological and non-pharmacological interventions designed to reduce or eliminate the pain that accompanies rocuronium bromide administration, both in adults and children, whether in asleep or awake state. **Materials and Methods:** We searched the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE via Ovid SP (1966 to December 2012), and EMBASE via Ovid SP (1980 to December 2012). We included all randomized controlled trials (RCTs) that compared the use of any drug or a non-pharmacological method with control patients, or those receiving no treatment to reduce the severity of pain with rocuronium injection. Our primary outcome was pain on rocuronium bromide injection measured by a pain score assessment. Our secondary outcomes were rise in heart rate and blood pressure following administration of rocuronium and adverse events related to the interventions. We used the standardized methods for conducting a systematic review as described by the *Cochrane handbook for Systematic Reviews of Interventions*. Two authors independently extracted details of trial methodology and outcome data from reports of all trials considered eligible for inclusion. All analyses were made on intention-to-treat basis. We used a fixed-effect model where there was no evidence of significant heterogeneity between studies and a random-effects model if heterogeneity was likely. **Results:** We included 66 studies with 8621 participants in the review, though most analyses were based on data from fewer participants. We noted substantial heterogeneity between trials. Therefore, it was not possible to combine the trials for the outcomes. All studies reported the primary outcome pain; few studies reported adverse events and no study reported heart rate and blood pressure changes after administration of rocuronium. **Conclusion:** Lidocaine was the most popular intervention drug used in 29 trials with 2256 participants. The relative risk (RR) of pain on injection if given lignocaine compared to placebo was 0.21; 95% CI