significant effect on the time to emergence. Of all the factors GCS and temperature at the end of the surgery had the maximum impact on the time to emergence in patients undergoing clipping for ruptured aneurysms.

ISNACC-S-29

Airway management in cervical spine disease - Observations from a tertiary care hospital

A. Karmakar, M. Y. Koura, N. Kumar

Hamad Medical Corporation, Doha, Qatar

Introduction: Airway management in patients with pre-existing cervical spine disease carries potential for further injury and requires careful consideration. Different studies have highlighted the anatomic considerations of different airway techniques. Generally, awake fiberoptic intubation is considered Gold Standard. We sought to study the airway management methods employed in patients presenting for cervical spine surgery at a tertiary care hospital and their impact. **Methods:** We conducted a retrospective observational study at a tertiary care hospital in Qatar as part of an institutional audit. Consecutive cases with cervical spine disease posted for elective spine surgery between July 2012 and December 2014 were observed for type of airway management technique, anesthetic used and signs of immediate post-operative neurological deterioration or any change in status from baseline. **Results:** A total of 98 patients met the inclusion criteria. 3 were excluded due to lack of data. 85% were males and 15% females. 55% had traumatic cervical spine disease and 34% degenerative cervical spine disease. 61 patients had preoperative motor weakness. For airway management, 56/95 patients underwent awake techniques (55 flexible fiberoptic; 1 Glidescope) with majority opting for remifentanyl plus propofol infusion (21/56) or TCI remifentanyl (18/56) for sedation. Remaining 39 patients were anesthetized before intubation. Only 2 patients developed neurologic deterioration after surgery. **Conclusions:** Majority of anesthesiologists had opted for awake fiberoptic technique. Progressive neurologic deterioration after surgery, occurred in only 2.1% of patients. No particular technique has conclusively shown superiority over others. Familiarity with a particular technique and good team communication is recommendable.

ISNACC-S-30

Effect of dexmedetomidine infusion of two different doses on cognitive function after surgery in elderly patients

P. Katariya, M. Kohli

Department of Anaesthesia, KGMU, Lucknow, Uttar Pradesh, India

Introduction: Postoperative cognitive dysfunction (POCD) is a significant social and financial problem occurring in a high percentage of cases in people over 60 years old. POCD is associated with additional costs due to hospital stays and increased need for institutionalization, rehabilitation, and home care. Dexmedetomidine (DEX) has been demonstrated to have a neuroprotective effect. **Methods:** The present study was a randomized, case-controlled double-blind trial of 90 patients who had undergone surgery with general anaesthesia (GA). The cognitive deficit of each patient was assessed using the Mini-Mental State Examination (MMSE). The 90 patients were randomly allocated into three groups: Group I: Patients receiving 0.25 mcg/kg/hr continuous infusion. Group II: Patients receiving 0.50 mcg/kg/hr continuous infusion. Group III: Normal saline group (control group). The SPSS Version 15.0 statistical Analysis Software was used for the statistical analyses. **Results:** Intragroup change in MMSE Score (from Baseline) at different time intervals (Paired 't' test) p values shows that there is no significant changes in MMSE score at 24 hr and at 72 hr in group II patient.

Post-op Group I (<0.001), group II (0.013), group III (<0.001)

At 24 hr op Group I (0.009), group II (0.489), group III (<0.001)

At 72 hr op Group I (0.009), group II (1.000), group III (0.003)

Conclusion: The findings of the present study shows that DEX infusion 0.50 mcg/kg/hr after loading with 1 mcg/kg over 10 min may be an effective method for ameliorating postoperative cognitive impairment in elderly patients who have undergone surgery. Only a small number of patients were recruited. Therefore, larger studies should be performed, expanding the present study to a broader population.

ISNACC-S-31

Anaesthetic management of a patient with permanent pacemaker posted for frontal lobe tumor excision

P. Kate, B. Shah, S. Khatavkar

Dr. D. Y. Patil Medical College, Pimpri, Pune, Maharashtra, India

Introduction: Patients undergoing neuro-anaesthesia with cardiac co-morbidities pose a considerable challenge to anaesthesiologists. Management of patients

with pacemaker in situ and anaesthetic implications are crucial. We report a successful anaesthetic management of a patient with pacemaker for frontal lobe tumor excision under general anaesthesia. **Case Summary:** 64 year old female with complaints of headache and irrelevant speech since 2 weeks, diagnosed to have frontal lobe tumor. Patient had history of syncope 5 years back and was diagnosed to have complete heart block. Thus a permanent pacemaker with VVI synchronised mode with heart rate 70/min was placed. On examination patient was moderately built, drowsy, heart rate was 68/min and rest vitals were within normal limits. Blood investigations were within normal limits. ECG showed pacemaker spikes just before QRS complexes and LVH. 2D Echo showed pacemaker lead in right ventricle and rest findings were normal. Patient accepted for surgery under ASA grade III with appropriate consent. Preoperatively central line was secured. Facility for temporary pacing and pacemaker technician were kept ready. Pacemaker settings were programmed from VVI mode to VVO mode. Patient was premedicated and induced with inj. thiopentone and inj. rocuronium, intubated and ventilated. Intraoperative vitals were maintained within normal limits. At the end of surgery patient was drowsy but maintained SpO₂ 100%. ET tube was kept in situ and patient was put on CPAP mode. VVO mode reprogrammed to VVI mode. Patient shifted to surgical ICU, all investigations were within normal limits. Patient was extubated in SICU 6 hours postoperatively. **Conclusion:** Neurosurgical patient with permanent pacemaker poses a real challenge to an anaesthetist. Thorough pre-op evaluation and team work of anaesthetists, physician and surgeon resulted in successful outcome.

ISNACC-S-32

Evaluation of prognostic factors of outcome in severe traumatic brain injury patients following decompressive craniectomy

A. Kaushal, A. Bindra, N. Kumar, K. Goyal, G. P. Rath

Department of Neuroanaesthesiology and Critical Care, AIIMS, New Delhi, India

Introduction: Severe traumatic brain injury (TBI) is leading cause of death and disability globally. Decompressive craniectomy (DC) is done to treat severe brain edema. We conducted a study to determine predictive factors for prognosis of DC at the time of cranioplasty. We correlated the admission variables, CT scan findings and timing of d/c, hospital stay with functional outcome. **Methods:** The functional outcome was noted prospectively and retrospective admission data was obtained from hospital records. All consenting

patients (15-65 yrs) posted for cranioplasty following DC due to severe TBI were enrolled. The data including Marshall grading, admission Glasgow coma score (GCS), mean arterial pressure (MAP), time of DC, duration of hospital stay, hospital readmission and GCS at discharge. The functional outcome at the time of cranioplasty was measured by Glasgow Outcome Score (GOSE) questionnaire. GOSE 5-8 were classified as good outcome and GOSE 1-4 were classified as poor outcome. **Results:** A total 85 patients (71 male and 14 female) were enrolled. The mean age was 33.42 yrs. Only 36% patients had a good outcome (GOSE 5-8). There was no significant association between age, sex, marshall grading, duration between d/c and cranioplasty, hospital readmission, GCS at admission with outcome. In univariate analysis tracheostomy, duration of hospital stay, MAP, timing of DC and GCS at discharge were predictors of outcome. On multivariate analysis tracheostomy was found to be independent predictor of outcome. **Discussion:** Significant disability is seen among the survivors of DC. Admission variables do not predict outcome. Larger sample size is required.

ISNACC-S-33

Non-neurosurgical complications in traumatic neurosurgical ICU patients: A prospective observational study

A. Khandelwal, K. Goyal, A. Hazarika¹, N. Sokhal, A. Bindra, N. Kumar, G. P. Rath, P. Bithal

Department of Neuroanaesthesiology and Critical Care, JPNATC, AIIMS, New Delhi ¹Department of Anaesthesiology, PGIMER, Chandigarh, India

Introduction: Recognising and treating non-neurological complications occurring in neurotrauma patients during ICU stay is equally challenging. Primary aim: To estimate various non-neurological complications in neurotrauma patients. Secondary aim: To see the effect of these complications on ICU stay, disability and mortality. **Methods:** Prospective observational study at neurotrauma ICU of level 1 trauma center, AIIMS, New Delhi. 200 neurotrauma patients consisting of Traumatic Brain Injury (TBI) and/or Cervical spine injury (CSI) were enrolled. Period of study: From admission to discharge from ICU or demise. Inclusion criteria: Age >16 years, Severe TBI (GCS≤8), CSI requiring mechanical ventilation. **Results:** Non-neurological complications were frequent in neurotrauma ICU patients. We observed respiratory complications to be of highest occurrence (60.5%). Other complications in decreasing order included dyselectrolytemia (40%), cardiovascular (33.5%), coagulopathy (32%), sepsis (24%), abdominal (16.5%) and AKI (3.5%). Presence of systemic