Score Establishment and Brazilian Portuguese version of the Pregnancy Sexual Response Inventory (PSRI)

Definição de escores e versão em português brasileiro do Inventário da Resposta Sexual na Gestação (PSRI)

Cibele Vieira Cunha Rudge¹ Iracema de Mattos Paranhos Calderon¹ Ana Paula Machado de Almeida¹
Fernanda Piculo¹ Marilza Vieira Cunha Rudge¹ Angélica Mércia Pascon Barbosa¹

¹Department of Gynecology and Obstetrics, Universidade Estadual Paulista Júlio de Mesquita Filho, Botucatu, SP, Brazil

Address for correspondence Angélica Mércia Pascon Barbosa, PhD, Av. Higino Muzi Filho, 737, 17525-900, Mirante, Marília, SP, Brazil (e-mail: angelicapascon@gmail.com).

Rev Bras Ginecol Obstet 2018;40:322–331.

Abstract

Objective To establish the Pregnancy Sexual Response Inventory (PSRI) scores for each domain before and during pregnancy, and to publish the Brazilian Portuguese version of the PSRI.

Methods Pregnant women were recruited during antenatal care; the PSRI was administered to 244 women prenatally at Faculdade de Medicina de Botucatu, at Universidade do Estado de São Paulo (UNESP, in the Portuguese acronym). The PSRI scores were estimated based on the Kings Health Questionnaire (KHQ) and the Medical Outcomes Study 36-item short form survey (SF-36). The raw scale type was used to standardize the minimal value and amplitude of each domain. For each domain, the score varied from 0 to 100, and the composite score was obtained as the domain average. The composite score before and during pregnancy was determined by the sum of the scores of all specific domains for each divided by the full domain number. The categorization of the scale into quartiles was established when all PSRI-specific and composite scores were combined.

Results The composite and specific scores for each domain were categorized into quartiles: 0 < 25 as “very bad;” 25 < 50 as “bad;” 50 < 75 as “good” and 75 to 100 as “excellent.” The mean scores were lower during pregnancy than before pregnancy in 8 of the 10 domains. The Brazilian Portuguese PSRI version is presented.

Conclusion This study allowed the establishment of the PSRI composite and specific scores for each domain, and the categorization of scores into quartiles: very bad, bad, good and excellent. In addition, the Brazilian Portuguese version of the PSRI is presented in full for application in the Brazilian population.

Keywords

- pregnancy
- sexuality
- questionnaires

Resumo

Objetivo Estabelecer os escores do Inventário da Resposta Sexual na Gestação (PSRI) para cada domínio antes e durante a gravidez, e publicar a versão do PSRI em português brasileiro.

received
October 24, 2017
accepted
April 19, 2018

DOI <https://doi.org/10.1055/s-0038-1656536>.
ISSN 0100-7203.

Copyright © 2018 by Thieme Revinter
Publicações Ltda, Rio de Janeiro, Brazil

License terms

Métodos Gestantes foram recrutadas durante o cuidado pré-natal; o PSRI foi administrado a 244 mulheres no pré-natal na Faculdade de Medicina de Botucatu da Universidade do Estado de São Paulo (UNESP). Os escores do PSRI foram estimados com base no *Kings Health Questionnaire* (KHQ) e *Medical Outcomes Study 36-item short form survey* (SF-36). O tipo de escala bruta foi utilizado para padronizar o valor mínimo e a amplitude de cada domínio. Para cada domínio, a pontuação variou de 0 a 100, e o escore composto foi obtido pela média do domínio. O escore composto antes e durante a gravidez foi determinado pela somatória dos escores de todos os domínios específicos para cada período dividido pelo número total do domínio. A escala de categorização em quartil foi estabelecida quando todos os escores específicos e compostos do PSRI foram reunidos.

Resultados Os escores compostos e específicos para cada domínio foram categorizados em quartis: $0 < 25$ como “muito ruim;” $25 < 50$ como “ruim;” $50 < 75$ como “bom” e 75 a 100 como “excelente.” As médias dos escores foram menores durante a gravidez do que antes da gravidez em 8 dos 10 domínios. Foi apresentada a versão PSRI em português brasileiro.

Conclusão Este estudo permitiu o estabelecimento dos escores compostos e específicos do PSRI para cada domínio e a categorização dos escores em quartis: muito ruim, ruim, bom e excelente. Além disso, a versão em português do PSRI é apresentada integralmente para aplicação na população brasileira.

Palavras-chave

- ▶ gestação
- ▶ sexualidade
- ▶ questionários

Introduction

There are several maternal adaptations that involve profound anatomical, physiological, and biochemical changes, which may impact the sexual health of partners during pregnancy.¹

A systematic review has found a gradual decrease in vaginal intercourse from prepregnancy to the first and third trimesters,² and many studies have revealed a reduction in sexual function during pregnancy.^{1,3–5}

This topic has attracted researchers' attention due to the increase in the number of epidemiological studies, but data are still limited on the prevalence of sexual dysfunction and concerns about sexual activity in pregnant women, and it remains unclear how to evaluate them. The female sex response cycle proposed by Basson (2000)⁶ starts during a neutral phase, and the rewards of emotional closeness serve as the motivational factors that will activate the cycle the next time. This knowledge needs to be included in the instruments used to evaluate the sexual function during pregnancy. Moreover, there are some attitude changes toward sexual function during pregnancy, such as the different sexual responses proposed by Basson (2000),⁶ but the methodological limitations (sample sizes, unrepresentative samples, and retrospective data) and inconsistent results of published manuscripts may limit their relevance.⁷

Currently, the instrument “Pregnancy and Sexuality Questionnaire (PSQ)” has been developed to evaluate the subjectivity and complexity of sexual function within pregnancy, although the authors did not list the specific items included in their questionnaire within their article.⁸

The “Female Sexual Function Index (FSFI)” was developed to evaluate female sexual response; however, this questionnaire was not developed for pregnant women.⁹ In turn, the Pregnancy and Sexual Function Questionnaire (PSFQ), Portuguese version, was considered adequate for evaluating sexual function during pregnancy.¹⁰

The Pregnancy Sexual Response Inventory (PSRI) was designed based on the PSQ, a validated instrument for studying sexual relations between partners during pregnancy,⁸ and was integrated into the Basson⁶ sexual response. This instrument was developed due to the lack of access to the only instrument validated for studying the sexual relationship of partners within pregnancy.

There were five phases in the development of the PSRI: (1) item selection; (2) item development; (3) determination of internal consistency, reliability and convergence; (4) content validity; and (5) determination of inter-interviewer reliability. Internal consistency and reliability were evaluated using Cronbach's α . Inter-interviewer reliability was assessed by evaluating the responses of 18 academics at various institutions using the Kappa Index and Student *t*-test.¹¹ Furthermore, the PSRI was fully validated in the Brazilian Portuguese language by our current research group and covers different domains of sexual response during pregnancy.¹¹ Although it is a validated questionnaire, the PSRI had not been published in Portuguese, and thus could not be used to support the clinical diagnosis of sexual function during pregnancy in Brazil and other Portuguese-speaking countries.

The aim of this study was to establish the PSRI composite and specific scores for each domain before and during pregnancy, and to publish the Brazilian Portuguese version of the PSRI.

Methods

Study Population

An observational, cross-sectional, single-center study was performed between January and August 2016 at the Department of Gynecology and Obstetrics at the Faculdade de Medicina de Botucatu (FMB-UNESP, in the Portuguese acronym). This hospital is a tertiary center with a perinatal center of the highest level providing health services to medium- and high-risk obstetrical patients from an area with ~ 500,000 inhabitants, and 1,600 deliveries are performed in it per year. Healthy pregnant women seeking antenatal care were recruited to participate in the current study while waiting for their routine medical check-ups. Any patients who presented systemic illnesses, such as diabetes mellitus, hypertension, hyperlipidemia and thyroid dysfunction, and those who conceived by assisted reproduction techniques were excluded from the current study.

The protocol and the objectives of the study were explained to 370 pregnant women; 249 (67.3%) of them provided a signed informed consent just before the administration of the validated instrument of sexual function - the PSRI. The eligibility criteria included healthy pregnant women who were heterosexual, 18 years of age or older, and in the second or third trimester of pregnancy and who had been sexually active in the previous 4 weeks.

Upon signing the informed consent, the eligible women were interviewed by a trained female interviewer using a paper-and-pencil standardized questionnaire. Interviews were conducted at the prenatal clinic in a private room. All women were assessed with a detailed medical history, including partnership status, education level, religion, employment status, parity, smoking habits, drinking, illicit drugs, planned pregnancy, and condom use, and a comprehensive physical examination was also performed for each woman. Our sample was mostly heterosexual, married, and in female-male relationships. The data were cross-sectional, which means we only collected one questionnaire per woman. Approval for the study was given by the local institutional research bureau under protocol number 161/2012.

Questionnaire

Sexual function was assessed using the PSRI. This semi-structured questionnaire contained 38 questions divided into 12 questions about demographic traits and 26 questions about sexual behavior activity before and during pregnancy. The sexual response questions were grouped in 10 domains; eight of them assessed the women's feelings, and two assessed their perception of her partner's sexual interest. All domains included possible distress items, since it is necessary to investigate sexual dysfunction.

► **Table 1** shows the questions grouped by domain for each period.

Outcome Measures

The primary outcomes were to make possible the establishment of scores to adequately evaluate the PSRI responses, and to publish the Portuguese version of the PSRI for application in the Brazilian population.

Table 1 Description of the grouped questions for each domain before and during pregnancy and the sum of all questions per domain

Domains	Questions	Questions	Questions
	Before pregnancy	During Pregnancy	All
PSRI (specific scores)			
Female perception			
Sexual activity frequency	14a	13, 14b, 14c	13, 14a, 14b, 14c
Desire	21a	21b, 22	21a, 21b, 22
Arousal	18a	18b	18a, 18b
Orgasm	23a	23b	23a, 23b
Satisfaction	15a, 17a	15b, 17b	15a, 15b, 17a, 17b
Dyspareunia	24a	24b	24a, 24b
Intercourse start	25a	25b	25a, 25b
Female difficulties	19a	19b	19a, 19b
Female perception of partners			
Male sexual satisfaction	16a	16b	16a, 16b
Male sexual difficulties	26a	26b	26a, 26b

Abbreviation: PSRI, Pregnancy Sexual Response Inventory.

The numbers followed by letters are the number of questions that appear in the PSRI.

PSRI Composite and Specific Score Establishment

The estimated PSRI scores of sexual behavior considered all answers before and during pregnancy, with the answers divided into each domain according to period. Therefore, 11 questions were analyzed before pregnancy, while 14 questions were analyzed during pregnancy. Two composite scores for the PSRI were established according to both analyzed periods. A score was calculated for each domain in both periods. The 20th question was not included in the score calculation because it was only answered if the 19th question was marked "yes." Demographic characteristics were not included in the PSRI score calculation. The PSRI score estimate was based on the Kings Health Questionnaire (KHQ)¹² and the Medical Outcomes Study, a 36-item short-form health survey (SF-36).¹³ The raw scale type was used to standardize the minimal value and amplitude of each domain. For each domain, the score varied from 0 to 100, and the general score was obtained using the domain average. The specific score for each domain was estimated using the SF-36 guidelines.¹³ The composite score comprising the periods before and during pregnancy was determined by adding the score of all specific domains for each period divided by the full domain number. Finally, we established the categorization scale into quartiles, once all the PSRI-specific and composite scores were combined (► **Fig. 1**).

I- Características Demográficas	
1- Idade Materna: Idade do Parceiro:	2- Idade Gestacional:
3- Estado Civil: (1) casada/união estável (2) solteira (3) outro _____	4- Nível Educacional: (1) fundamental (2) ensino médio (3) ensino superior
5- Religião: (1) católica (2) evangélica (3) outras _____	6- Você trabalha? (1) não (2) sim, eu tenho um trabalho (3) sim, mas no momento estou desempregada
7- Você tem filhos? (1) não (2) apenas um (3) dois ou mais	8- Você fuma? (1) sim, com alguma ou muita frequência (2) sim, apenas as vezes (3) não
9- Você bebe? (1) sim, com alguma ou muita frequência (2) sim, apenas às vezes (3) não	10- Você usa drogas ilícitas? (1) sim, com alguma ou muita frequência (2) sim, apenas às vezes (3) não
11- Você planejou sua gravidez? (1) sim (2) não	12- Você usa preservativo? (1) sim (2) não
II- Comportamento/Atividade Sexual – antes e durante a gestação	
13- Na sua opinião, a frequência das suas relações sexuais mudou depois que você engravidou? (1) sim, diminuiu (2) não, é a mesma (3) sim, aumentou	14a- Antes da gestação, quantas vezes por semana você tinha relações sexuais? (1) nenhuma (2) 1-2 vezes (3) 3 ou mais vezes
14b- No primeiro trimestre da gestação, quantas vezes por semana você tinha relações sexuais? (1) nenhuma (2) 1-2 vezes (3) 3 ou mais	14c- No momento, quantas relações sexuais você tem por semana? (1) nenhuma (2) 1-2 vezes (3) 3 ou mais
15a- Como você classificaria sua vida sexual antes de você engravidar? (0 = muito ruim, 10 = muito boa) (1) 0-3 (2) 4-7 (3) 8-10	15b- Como você classificaria sua vida sexual atualmente? (0 = muito ruim, 10 = muito boa) (1) 0-3 (2) 4-7 (3) 8-10

Fig. 1 Full version of the Brazilian Portuguese Pregnancy Sexual Response Inventory.

16a- Como você acha que o seu parceiro classificaria a vida sexual dele antes de você engravidar? (1) 0-3 (2) 4-7 (3) 8-10	16b- Como você acha que o seu parceiro classificaria a vida sexual dele atualmente? (1) 0-3 (2) 4-7 (3) 8-10
17a- Você tinha prazer nas suas relações sexuais antes de engravidar? (1) não (2) eu suponho que estava tudo bem (3) sim	17b- Você tem prazer nas suas relações sexuais durante a gestação? (1) não (2) eu suponho que esteja tudo bem (3) sim
18a- Como você classificaria sua excitação antes da gestação? (1) baixa/muito baixa (2) regular (3) excelente	18b- Como você classificaria sua excitação durante a gestação? (1) baixa/muito baixa (2) regular (3) excelente
19a- Você apresentava alguma dificuldade sexual antes da gestação? (1) sim (2) não	19b- Você tem apresentado alguma dificuldade sexual durante a gestação? (1) sim (2) não
20- Essas dificuldades te deixam angustiada? (1) sim (2) um pouco (3) não	21a- Com que frequência você tinha desejo sexual antes da gravidez? (1) nunca/ raramente (2) algumas vezes por semana (3) uma vez por dia
21b- Com que frequência você tem desejo sexual durante a gravidez? (1) nunca/raramente (2) algumas vezes por semana (3) uma vez por dia	22- O que aconteceu com o seu desejo sexual depois que você engravidou? (1) diminuiu (2) é o mesmo (3) aumentou
23a- Com que frequência você alcançava o orgasmo antes da gestação? (1) nunca/raramente (2) as vezes (3) com frequência/com muita frequência	23b- Com que frequência você alcança o orgasmo durante a gestação? (1) nunca/raramente (2) as vezes (3) com frequência/com muita frequência
24a- Você sentia dor durante a relação sexual antes da gestação? (1) sim (2) não	24b- Você sente dor durante a relação sexual durante a gestação? (1) sim (2) não

Fig. 1 (Continued)

25a- A relação sexual antes da gestação era iniciada: (1) forçada, sem nenhum desejo (2) geralmente iniciada pelo parceiro (3) espontaneamente ou espontaneamente com estímulo	25b- A relação sexual durante a gestação é iniciada: (1) forçada, sem nenhum desejo (2) geralmente iniciada pelo parceiro (3) espontaneamente ou espontaneamente com estímulo
26a- Na sua opinião, você acha que o seu parceiro estava apresentando alguma dificuldade sexual antes da gestação? (1) sim (2) não	26b- Na sua opinião, você acha que o seu parceiro está apresentando alguma dificuldade sexual durante a gestação? (1) sim (2) não

Fig. 1 (Continued)

Portuguese Version

The Brazilian Portuguese version of the PSRI is presented in the same format as the English one.

Statistical Analyses

The sample size was calculated according to the 40% prevalence of sexual dysfunction in pregnant women, with a margin of error of 10% and a reliability of 95%.¹⁴ Thus, the minimum sample size was determined to be 184 participants.

Comparisons between means of the domain values classified by both analyzed periods were assessed by paired *t*-test at a significance level of 5%. All data were analyzed using the software Statistical Analysis System (SAS) for Windows, version 9.2 (SAS Institute Inc., Cary, NC, USA).

Results

► **Fig. 2** provides an overview of the study sample collection.

The Brazilian Portuguese PSRI, a validated questionnaire, is shown in ► **Fig. 1**. Two hundred and forty-nine pregnant women completed the PSRI, with 49 in the second trimester of pregnancy, 200 in the third trimester of pregnancy and 5 excluded from the final sample because their questionnaires were incomplete. ► **Table 2** represents the demographic features of our full sample. The mean maternal age of the 244 participants was 26 years (SD = 5.4, Min = 20.6, Max = 31.4). At study inclusion, the mean gestational age was 34.8 weeks of pregnancy (SD = 3.5, Min = 25.0, Max = 42.0). The majority of our sample (63.1%) was married

Fig. 2 Flow-diagram describing the process for recruitment of the pregnant women.

Table 2 Descriptive demographic characteristics of pregnant women

Variables	f (%)
Partnership status	
Married/Living together	154 (63.1)
Single	71 (2.1)
Other	19 (7.8)
Sociodemographic factors	
Education Level	
Basic Level	99 (40.6)
High School	120 (49.2)
College/University	25 (1.2)
Religion	
Catholic	113 (46.3)
Brazilian Protestants	95 (38.9)
Other/No religion	36 (14.8)
Employment status	
Student	72 (29.5)
Employed	86 (35.2)
Not employed	86 (35.2)
Children	
No	128 (52.5)
Just one	71 (29.1)
Two or more	45 (18.4)
Smoke	
Often/Very often	20 (8.2)
Sometimes	21 (8.6)
No	203 (83.2)
Drink	
Often/Very often	2 (0.8)
Sometimes	11 (4.5)
No	231 (94.7)
Illicit drugs	
Often/Very often	5 (2.0)
Sometimes	2 (0.8)
No	237 (97.1)
Family planning knowledge	
Planned pregnancy	
Yes	109 (44.7)
No	135 (55.3)
Contraceptive methods*	
No	199 (81.6)
Yes, stopped before pregnancy	28 (11.5)
Very often	17 (7.0)

Abbreviation: f, frequency of clinical characteristics of the study population.

or living together, primigravida (52.5%) and had studied until elementary school (59.4%). From our sample, 40.6% were Catholic, 38.9% were Brazilian Protestants, and the rest answered another or no religion. A high proportion of the

respondents were students (29.5%) and employed full- or part-time (35.2%). Only a small percentage (16.8%) reported smoking at least half a pack of cigarettes per day, and 94.7% responded that they did not drink alcohol even socially. A history of illicit drug use was observed in 2.8% of all respondents. A high percentage of our sample (55.3%) declared that pregnancy was unplanned, and 81.6% did not use condoms. Additional assessed demographics can be seen in ►Table 2.

Composite and Specific Scores Measured by Domains for PSRI

►Fig. 3 shows the questions grouped by each domain and by each period, and the composite score for the PSRI specific score measurements before and during pregnancy in the studied population. As the options for the PSRI answers are graduated from minimal to maximal values, “0” is considered the worst and “100” the best. These values are the inverse of the KHQ, for which the answer options are graduated from “best” to “worse” values.

The score was categorized into quartiles by sexual response as follows: $0 < 25$ as “very bad,” $25 < 50$ as “bad,” $50 < 75$ as “good” and 75 to 100 as “excellent.” Using this established quartile-categorized score for PSRI composite scores before and during pregnancy allowed us to accurately identify the quality of the answers of each domain and the sum of the domains of the composite score (►Fig. 3).

Influence of Pregnancy on Sexual Response as Evaluated by the PSRI

►Table 3 shows the results of the specific and composite scores before and during pregnancy. During pregnancy, the specific scores were lower than before pregnancy in almost all of the PSRI domains (sexual activity frequency, arousal, orgasm, satisfaction, dyspareunia, intercourse start, female difficulties and male sexual satisfaction) ($p < 0.05$), thus suggesting a negative impact of pregnancy on sexual function response. A significant increase in the desire score was observed, but no significant difference in male sexual difficulties was shown between the periods. The composite score of sexual activity as evaluated by the PSRI showed a significant decrease from pre-pregnancy (mean score = 83 “excellent”) to during pregnancy (mean score = 66 “good”).

Discussion

Sexual function during pregnancy is an aspect of quality of life. The World Health Organization defined sexual health as “a state of physical, emotional, mental, and social wellbeing related to sexuality.”¹⁵ Sexual dysfunctions are defined as disorders related to both sexual desire and sexual satisfaction for several reasons.¹⁶

Pregnancy is a process of alteration experienced by women, and as a consequence, sexual life also changes during pregnancy,¹⁷ although there is a lack of specific instruments in the literature to confirm the influence of pregnancy on sexual function. Many non-specific questionnaires to characterize this adjustment of sexual function in pregnant women have been published.¹⁸ The FSFI questionnaire has been used to

PSRI Specific Score	
Before pregnancy	During pregnancy
Frequency score $FS = \frac{(Q14a - 1)}{2} \times 100$	Frequency Score $FS = \frac{[(Q13 + Q14b + Q14c) - 3]}{6} \times 100$
Desire Score $DS = \frac{(Q21a - 1)}{2} \times 100$	Desire Score $DS = \frac{[(Q21b + Q22) - 2]}{4} \times 100$
Arousal Score $AS = \frac{(Q18a - 1)}{2} \times 100$	Arousal Score $AS = \frac{(Q18b - 1)}{2} \times 100$
Orgasm Score $OS = \frac{(Q23a - 1)}{2} \times 100$	Orgasm Score $OS = \frac{(Q23b - 1)}{2} \times 100$
Satisfaction Score $SS = \frac{[(Q15a + Q17a) - 2]}{4} \times 100$	Satisfaction Score $SS = \frac{[(Q15b + Q17b) - 2]}{4} \times 100$
Dyspareunia Score $DyS = (Q24a - 1) \times 100$	Dyspareunia Score $DyS = (Q24b - 1) \times 100$
Intercourse Start Score $ISS = \frac{(Q25a - 1)}{2} \times 100$	Intercourse Start Score $ISS = \frac{(Q25b - 1)}{2} \times 100$
Female Difficulties Score $FDS = (Q19a - 1) \times 100$	Female Difficulties Score $FDS = (Q19b - 1) \times 100$
Male sexual satisfaction score $MSSS = \frac{(Q16a - 1)}{2} \times 100$	Male sexual satisfaction score $MSSS = \frac{(Q16b - 1)}{2} \times 100$
Male difficulties score $MDS = (Q26a - 1) \times 100$	Male difficulties score $MDS = (Q26b - 1) \times 100$
Composite Score* measurements $\text{Composite Score} = \frac{FS + DS + AS + OS + SS + DyS + ISS + FDS + MSSS + MDS}{10}$ <p>* The composite score must be applied for each period: before and during pregnancy. FS: Frequency sexual activity score; DS: Desire score; AS: Arousal score; SS: Satisfaction score; DyS: Dyspareunia score; ISS: Intercourse Start score; FDS: Female difficulties score; MSSS: Male sexual satisfaction score; MDS: Male difficulties score.</p>	

Fig. 3 Pregnancy Sexual Response Inventory I composite and specific scores for each domain before and during pregnancy.

assess sexual function, showing low values in the third trimester.^{19,20} However, it is essential to emphasize that the current and most frequent use of the FSFI is for non-pregnant women, for whom it was designed and validated. The PSRI is a specific questionnaire that was designed to consider the influence of pregnancy on sexual behavior using a self-evaluation before

and during pregnancy. This differences in the design and drafting of the questionnaires need to be taken into account when considering the disparities in the results published in various articles, which result in a lack of consensus.

The findings presented here in our study using the PSRI indicate that the composite and specific scores for each

Table 3 Pregnancy Sexual Response Inventory composite and specific scores before and during pregnancy

Domains	Before pregnancy	During pregnancy	p Value
	Mean $\pm$ SD	Mean $\pm$ SD	
Frequency score	72.95 $\pm$ 28.63	43.83 $\pm$ 29.4	0.00 *
Desire score	48.58 $\pm$ 42.23	63.61 $\pm$ 27.7	0.02 *
Arousal score	79.18 $\pm$ 27.1	54.63 $\pm$ 31.56	0.00 *
Orgasm score	95.55 $\pm$ 16.57	72.95 $\pm$ 34.04	0.00 *
Satisfaction score	86.3 $\pm$ 19.68	64.06 $\pm$ 30.58	0.00 *
Dyspareunia score	89.68 $\pm$ 30.48	70.11 $\pm$ 45.86	0.00 *
Intercourse start score	85.23 $\pm$ 23.24	81.67 $\pm$ 24.5	0.01 *
Female difficulties score	92.52 $\pm$ 26.34	67.61 $\pm$ 46.88	0.00 *
Male sexual satisfaction score	82.74 $\pm$ 30.69	49.46 $\pm$ 40.85	0.00 *
Male sexual difficulties score	97.15 $\pm$ 16.66	95.73 $\pm$ 20.25	0.13
Composite score	82.99 $\pm$ 9.76	66.25 $\pm$ 15.14	0.00 *

Abbreviation: SD, standard deviation. * $P < 0.05$.

domain and from prepregnancy to pregnancy were established. The scores were significantly different and categorized into quartiles by sexual response as follows: 0 < 25 as “very bad,” 25 < 50 as “bad,” 50 < 75 as “good” and 75 to 100 as “excellent” for before and during pregnancy. The results indicated that lower composite and specific scores occurred during pregnancy than before pregnancy in almost all PSRI domains (sexual activity frequency, arousal, orgasm, satisfaction, dyspareunia, intercourse start, female difficulties and male sexual satisfaction).

These results may indicate the negative impact of pregnancy on sexual function response. However, some authors demonstrated no difference in general scores between the 1st and 2nd trimesters but a significant association between decreased intercourse frequency and trimesters.⁴ Galazka et al (2015)⁵ found that desire, arousal, lubrication, orgasm, satisfaction, pain and sexual activity frequency decrease as gestation advances. Most of our findings are in line with the recent literature, which characterizes the perinatal period by a low sex drive.^{21,22} Women also seem to report higher levels of FSD female sexual dysfunction and low sexual desire, which is potentially associated with overall physical discomfort.^{23,24}

Our results suggest that it is possible to quantitatively assess the impact of pregnancy on sexual response through score estimations before and during pregnancy, allowing comparisons of women’s real sexual state during different pregnancy periods. As hypothesized, the PSRI scores could

allow us to understand the influence of pregnancy on sexual health not only in qualitative but also in quantitative parameters for each domain. By using scores, clinicians can better plan and implement strategies and health programs targeted at improving sexual health for pregnant partners.

Identifying pregnant women who experience sexual distress and referring them to appropriate resources could help to minimize sexual and relationship problems during pregnancy.²⁵ These strategies are important not only for clinical assistance but also to teach and train undergraduates of medicine because most of them do not feel comfortable or confident, and they lack specific knowledge and skills to address questions related to sexual problems within pregnancy.²⁶

Despite fears and myths about sexual activity during pregnancy, maintaining sexual interactions throughout the pregnancy and postpartum period can promote sexual health, well-being and a greater depth of intimacy. An open discussion about the expected changes in sexual health could provide guidance for couples, as well as promote rigorously designed, evidence-based studies to further elucidate our understanding of sexual function during pregnancy and postpartum.²⁷

Although far from conclusive, these results are consistent with the hypothesis that a clinical diagnostic assessment using PSRI scores enables and facilitates an understanding of the current pregnancy sexual response and changes in sexual response before and during pregnancy. Our results, in particular, can indicate that clinical scores may represent a key strategy for implementing specific health programs to improve sexual health for pregnant partners.

As with many studies, it is important to consider the potential strengths and weaknesses of the clinical PSRI scores, as well as their use in further clinical practice and research implications.

The current study’s strength relies on the use of a validated instrument to assess sexual function during pregnancy.¹¹ We acknowledge that using additional questionnaires to evaluate the sexual symptoms and quality of life of the participants could have enriched our study. Finally, the current study’s limitations involve our sample, which mostly comprised heterosexual married women, which prevents our findings from being extrapolated to a broader population of pregnant women. More studies involving women of other social and cultural contexts are needed to confirm such findings. As the PSRI is a generic questionnaire, its value for pregnancy comorbidities should be investigated.

Despite these limitations, the current study advances the understanding of the inter-relationships between maternal sexual response before and during pregnancy. As such, our findings regarding the clinical scores for the potential classification of pregnant women’s sexual dysfunction may have implications for evidence-based practice in preventative and intervention efforts, as well as in scientific study. The ultimate goal would be to implement early treatment and support (ideally before pregnancy) to improve the couple’s sexual health outcomes. Further studies are needed to establish the cutoff score to be used to indicate normal sexual function during pregnancy and sexual dysfunction during pregnancy.

Nonetheless, there are several important clinical implications of our findings. First, the current study enriches the literature because a validated questionnaire can establish clinically meaningful scores, supporting the efforts of other nations to translate and apply such instruments in specific pregnancy comorbidities. Additionally, we can encourage healthcare providers to use the PSRI scores for composite and specific domains to determine the influence of pregnancy on each one of the sexual response domains. Finally, the PSRI is a unique validated instrument designed specifically to evaluate at the same time the sexual response before and during pregnancy.

The Brazilian Portuguese version of the PSRI is published within the current manuscript, which allows Portuguese speakers to administer the questionnaire during antenatal care. According to the results, pregnant women or couples would be referred to a sexologist.

Conclusion

This study allowed the establishment of PSRI composite and specific scores for each domain, between 0 and 100, and the categorization of scores into quartiles: very bad, bad, good and excellent. In addition, the Portuguese version of the PSRI is presented in full for application in the Brazilian population.

Contributors

Rudge C. V. C., Calderon I. M. P., Almeida A. P. M., Piculo F., Rudge M. V. C. and Barbosa A. M. P. contributed with the project and interpretation of data, writing of the article, critical review of the intellectual content and final approval of the version to be published.

Conflicts of Interest

No conflicts of interest have been declared by the authors.

Acknowledgments

This study was funded by *Fundação de Amparo à Pesquisa do Estado de São Paulo* (FAPESP, in the Portuguese acronym), grants 2012/25207-4 and 2012/15577-9.

References

- Staruch M, Kucharczyk A, Zawadzka K, Wielgos M, Szymusik I. Sexual activity during pregnancy. *Neuroendocrinol Lett* 2016;37(01):53–58
- Jawed-Wessel S, Sevic E. The impact of pregnancy and childbirth on sexual behaviors: a systematic review. *J Sex Res* 2017;54(4-5):411–423. Doi: 10.1080/00224499.2016.1274715
- Aydin M, Cayonu N, Kadihasanoglu M, Irkilata L, Atilla MK, Kendirci M. Comparison of sexual functions in pregnant and non-pregnant women. *Urol J* 2015;12(05):2339–2344. Doi: 10.22037/uj.v12i5.2881
- Corbacioglu Esmer A, Akca A, Akbayir O, Goksedef BP, Bakir VL. Female sexual function and associated factors during pregnancy. *J Obstet Gynaecol Res* 2013;39(06):1165–1172. Doi: 10.1111/jog.12048
- Gałązka I, Drosdzol-Cop A, Naworska B, Czajkowska M, Skrzypulec-Plinta V. Changes in the sexual function during pregnancy. *J Sex Med* 2015;12(02):445–454. Doi: 10.1111/jsm.12747
- Basson R. The female sexual response: a different model. *J Sex Marital Ther* 2000;26(01):51–65. Doi: 10.1080/009262300278641
- Bartellas E, Crane JM, Daley M, Bennett KA, Hutchens D. Sexuality and sexual activity in pregnancy. *BJOG* 2000;107(08):964–968. Doi: 10.1111/j.1471-0528.2000.tb10397.x
- Barclay L, Bond M, Clark M. Development of an instrument to study the sexual relationship of partners during pregnancy. *Aust J Adv Nurs* 1992–1993;10(02):14–21
- Rosen R, Brown C, Heiman J, et al. The Female Sexual Function Index (FSFI): a multidimensional self-report instrument for the assessment of female sexual function. *J Sex Marital Ther* 2000;26(02):191–208. Doi: 10.1080/009262300278597
- Amaral TL, Monteiro GT. [Translation and validation of the Pregnancy and Sexual Function Questionnaire (PSFQ)]. *Rev Bras Ginecol Obstet* 2014;36(03):131–138. Doi: 10.1590/S0100-72032014000300007
- Rudge CV, Calderon IM, Dias A, et al. Design and validity of a questionnaire to assess sexuality in pregnant women. *Reprod Health* 2009;6:12. Doi: 10.1186/1742-4755-6-12
- Tamanini JT, D'Ancona CA, Botega NJ, Rodrigues Netto N Jr. [Validation of the Portuguese version of the King's Health Questionnaire for urinary incontinent women]. *Rev Saude Publica* 2003;37(02):203–211. Doi: 10.1590/S0034-89102003000200007
- Stewart M. The Medical Outcomes Study 36-item short-form health survey (SF-36). *Aust J Physiother* 2007;53(03):208. Doi: 10.1016/S0004-9514(07)70033-8
- Leite AP, Campos AA, Dias AR, Amed AM, De Souza E, Camano L. Prevalence of sexual dysfunction during pregnancy. *Rev Assoc Med Bras* (1992) 2009;55(05):563–568. Doi: 10.1590/S0104-42302009000500020
- World Health Organization. *Defining Sexual Health: Report of a Technical Consultation on Sexual Health 28–31 January 2002*. Geneva: WHO; 2006
- Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States: prevalence and predictors. *JAMA* 1999;281(06):537–544. Doi: 10.1001/jama.281.6.537
- Gökyıldız S, Beji NK. The effects of pregnancy on sexual life. *J Sex Marital Ther* 2005;31(03):201–215. Doi: 10.1080/00926230590513410
- Leite APL, Moura EA, Campos AAS, Mattar R, Souza E, Camano L. [Validation of the Female Sexual Function Index in Brazilian pregnant women] *Rev Bras Ginecol Obstet* 2007;29:396–401. Doi: 10.1590/S0100-72032007000800003
- Ribeiro MC, Nakamura MU, Torloni MR, Scanavino MdeT, Scomparini FB, Mattar R. Female sexual function of overweight women with gestational diabetes mellitus – a cross-sectional study. *PLoS One* 2014;9(04):e95094. Doi: 10.1371/journal.pone.0095094
- Aslan G, Aslan D, Kizilyar A, Ispahi C, Esen A. A prospective analysis of sexual functions during pregnancy. *Int J Impot Res* 2005;17(02):154–157
- Aslan E, Beji NK, Gungor I, Kadioglu A, Dikencik BK. Prevalence and risk factors for low sexual function in women: a study of 1,009 women in an outpatient clinic of a university hospital in Istanbul. *J Sex Med* 2008;5(09):2044–2052. Doi: 10.1111/j.1743-6109.2008.00873.x
- Erol B, Sanli O, Korkmaz D, Seyhan A, Akman T, Kadioglu A. A cross-sectional study of female sexual function and dysfunction during pregnancy. *J Sex Med* 2007;4(05):1381–1387. Doi: 10.1111/j.1743-6109.2007.00559.x
- DeJucibus MA, McCabe MP. Psychological factors and the sexuality of pregnant and postpartum women. *J Sex Res* 2002;39(02):94–103. Doi: 10.1080/00224490209552128
- Byrd JE, Hyde JS, DeLamater JD, Plant EA. Sexuality during pregnancy and the year postpartum. *J Fam Pract* 1998;47(04):305–308. Doi: 10.1080/00224499609551826
- Vannier SA, Rosen NO. Sexual distress and sexual problems during pregnancy: associations with sexual and relationship satisfaction. *J Sex Med* 2017;14(03):387–395
- Vieira TC, de Souza E, Abdo CH, et al. Brazilian residents' attitude and practice toward sexual health issues in pregnant patients. *J Sex Med* 2012;9(10):2516–2524. Doi: 10.1111/j.1743-6109.2012.02809.x
- Johnson CE. Sexual health during pregnancy and the postpartum. *J Sex Med* 2011;8(05):1267–1284, quiz 1285–1286. Doi: 10.1111/j.1743-6109.2011.02223.x