Figure 1: Overall bispectral index trends in pericraniotomy period

Figure 2: Systolic blood pressure trends compared between poor Glasgow outcome scale and good Glasgow outcome scale cases

Introduction: Poor neurological status in traumatic brain injury (TBI) patients can be due to raised intracranial pressure or underlying brain parenchymal injury or a combination. Surgical decompression would improve the neurological condition of the former but not the later. The purpose of this study is to observe trends of intraoperative haemodynamics and bispectral index (BIS) in head injury cases. **Methods:** In this prospective study, 30 TBI (subdural haematoma or contusion) patients undergoing emergency cranial decompression were to be recruited. BIS electrode was placed in an alternate fashion on contralateral fronto-temporal region. Baseline haemodynamics and BIS were recorded. Further recordings were taken after anaesthesia induction, pre-craniotomy, post-craniotomy (5, 10, 15 min), post-durotomy (5, 10, 15 min) and 10 min at the end of surgery after discontinuation of anaesthetics. Glasgow coma scale before and after surgery, and Glasgow outcome scale (GOS) at hospital discharge were noted. **Results:** Currently, interim analysis of 20 patients showed significant changes between systolic blood

pressure (SBP) ($P = 0.002$) and BIS ($P = 0.007$) during pre-craniotomy, 15 min post-craniotomy and 15 min post-durotomy [Figure 1]. SBP decreased marginally after craniotomy ($P = 0.005$) followed by a decrease after durotomy ($P = 0.017$). BIS showed significant increase post-craniotomy ($P = 0.008$) followed by a significant reduction after durotomy ($P = 0.05$). Mean changes in BIS and SBP after durotomy were 5.08 ± 1.8 and 18.92 ± 5.51 mmHg respectively. Heart rate (HR) and diastolic pressures showed no difference. On sub-classification of cases into good GOS (3, 4, 5) and poor GOS (1, 2), we noted no significant difference in trends of change of BIS ($P = 0.104$) and HR ($P = 0.287$) but a significant difference in SBP trends ($P = 0.008$) [Figure 2].

Discussion: The association between trends of SBP and BIS values indicates impaired cerebral autoregulation. BIS increases immediately after craniotomy and may or may not be related to blood pressures. The outcome of patients seems to be correlated with trends of change of SBP intraoperatively in the pericraniotomy period. Further updated results will be discussed after achieving higher sample size.

ISNACC-S-18

Comparison of midazolam and target control based propofol infusion in refractory status epilepticus

Dheeraj Masapu, G. S. Umamaheswara Rao, Sanjib Sinha¹, K. N. Gopala Krishna

Departments of Neuroanaesthesia and ¹Neurology, National Institute of Mental Health and Neurosciences, Bengaluru, Karnataka, India

Background: Refractory status epilepticus (RSE) is defined as SE that fails to respond to first- and any two drugs in the second-line therapy. Propofol and midazolam are commonly used drugs for RSE; however, there is no consensus regarding superiority of either drug. This study was conceived to compare these two drugs for efficacy for RSE control and clinical outcome. **Methodology:** Twenty-three patients of diagnosed RSE were recruited prospectively and randomised to receive either propofol or midazolam infusion for seizure control, under electroencephalography monitoring according to a standardised protocol. Baseline demographics, time to seizure control, drug utilisation trends, adverse events, haemodynamic, acid-base data and outcome data were collected. Trend based parameters were analysed using mixed models ANOVA, between-group comparison was done using Mann-Whitney U-test or Chi-square test as applicable. Drug trends were analysed using Kaplan-Meier survival curves. **Results:** There was significant difference between the groups for duration of mechanical ventilation (propofol vs. midazolam) (median [interquartile range]) 9 [4.5–12.5] vs. 15 [14–23]

Figure 1: Kaplan–Meier curve shows that the early stoppage of midazolam secondary to the failure of control of refractory status epilepticus or infusion termination due to severe hypotension ($P = 0.005$)

days, $P = 0.006$), duration of Intensive Care Unit stay (11 [6–13.5] vs. 15 [14.25–23] days, $P = 0.006$) and duration of hospital stay (11 [7–13.5] vs. 22 [15.75–43.25] days, $P = 0.006$). The difference of time taken for seizure control between the groups was trending toward significance (4 [3–4.5] vs. 5 [4–12] h, $P = 0.091$). Survival curve analysis showed a significant difference between the groups for stoppage of infusions due to either hypotension or inadequate seizure control, with midazolam recording earlier infusion discontinuation (Breslow's test, $P = 0.005$) [Figure 1]. Termination of infusions due to successful seizure control did not show any significant difference ($P = 0.609$). **Conclusion:** Despite the limitation of small sample size, outcomes seem to be better with propofol infusion rather than midazolam. Future studies with larger sample sizes are likely to validate this finding.

ISNACC-S-19

Bispectral index artefact compendium: Mechanisms explored

Dhritiman Chakrabarti,
Venkatapura J. Ramesh, Deepti B. S.

National Institute of Mental Health and Neurosciences,
Bengaluru, Karnataka, India

Introduction: Bispectral index (BIS), the widely accepted monitoring modality for depth of hypnosis under anaesthesia, has had multiple versions improving upon the original algorithm. Artefact recognition and removal are important aspects of any neuromonitoring modality. This article aims to review the various artefacts recognised during BIS monitoring along with additional unpublished observations and mechanisms for the same. **Methodology:** PubMed was searched using

keywords 'Bispectral index AND (Artifact OR Artefact)' and relevant articles were compiled. Cross referencing from these articles, a list of artefacts concerning BIS were prepared. Wherever mechanisms of artefact processing were not elucidated satisfactorily, a patent review under assignee 'Aspect Medical Systems' was done to obtain additional information. Additional unpublished artefacts were also listed and researched similarly. **Results:** The list of artefacts recognised was either physiological (electrocardiographic, frontalis electromyography [EMG], seizure) or external (electrical - facial nerve stimulator; cautery - bipolar/monopolar, electrical line interference, CUSA, coil detacher artefact, electromagnetic shaver; mechanical - forced air warmer). **Discussion:** Almost all artefacts presented as increase in BIS EMG with or without signal quality index deterioration with or without BIS fluctuation. High-amplitude signal contaminations were usually recognised as artefacts by the BIS monitor while low-amplitude ones were not. Signal contamination occurred even when frequency range of artefact signal was out of bandwidth allowed by BIS filters (may be attributed to Gibbs phenomenon). The exact mechanism behind each artefact is explained and when possible, a method to ameliorate the undesirable signal is suggested.

ISNACC-S-20

Magnetic resonance imaging predictors of intraoperative bleeding in intracranial neurosurgical tumour surgery

Deepti B.S., Dhritiman Chakrabarti,
Radhakrishnan M.

National Institute of Mental Health and Neurosciences,
Bengaluru, Karnataka, India

Introduction: Intraoperative blood loss during intracranial tumour surgery is a common and serious complication and predicting the same is challenging task for the anaesthesiologist. Although radiological scans can be used to qualitatively predict intraoperative blood loss, no current literature explores it systematically. This study is designed with the purpose of elucidating the magnetic resonance imaging (MRI) characteristics predictive of intraoperative bleeding. **Methodology:** Retrospective Observational Single centre study. E-hospital for haemoglobin (Hb) (pre-operative and post-operative) values and InstaRIS PACS MRIs of intracranial tumour surgery patients was collected. Putative predictors of blood loss from MRIs were: (1) Size of tumour, (2) proximity to venous sinuses, (3) proximity to major intracranial arteries and first-degree branches, (4) pattern of contrast enhancement, (5) contrast enhancement ratio, (6) suspected histopathology of tumour, (7) percentage of areas of hypointensity (necrosis) within tumour, (8) average pixel intensity of susceptibility weighted