images, (9) number of flow voids within tumour and (10) peritumoural oedema. The method utilised involved calculation of estimated blood loss from pre-operative and post-operative Hb values and intraoperative blood transfusion weight of patients. Formula used was $(\text{Hb}_{\text{pre}} - \text{Hb}_{\text{post}}) / \text{Hb}_{\text{pre}} \times \text{estimated blood volume (EBV)} + \text{intraoperative BT volume administered}$. $\text{EBV} = 70 \times \text{weight (females)}; 65 \times \text{weight (males)}$. $\text{BT volume} = \text{packed cell volume (PCV)} (0.6/0.35) = \text{PCV} \times 1.714$ or whole blood volume. A sample size of 100 patients was planned for producing a predictive model. Spearman's correlation analysis was used for association between blood loss and MRI characteristics. Linear and logistic regression were used for identifying independent predictors of volume of intraoperative blood loss and predicting high blood loss (>25% of EBV). **Results:** Currently, 20 patients have been recruited. None of the above parameters appear to be significantly correlated with the calculated blood loss. Thus, none of the parameters were used for predictive modelling. Updated results will be discussed after complete data collection. **Conclusion:** The importance of quantitative prediction of expected blood loss in anaesthetic practice cannot be undermined in neurosurgeries. With no precedence of such study in known literature, we expect this study to be useful and hopefully lay background for future detailed research in this niche.

ISNACC-S-21

Hypothermia in traumatic brain injury for control of intracranial hypertension: Standalone therapeutic option or adjunct?

Ashish Bindra, Deepak Gupta¹, Peter Andrews²

Departments of Neuroanaesthesiology and Critical Care and ¹Neurosurgery, JPN Apex Trauma Centre, All India Institute of Medical Sciences, New Delhi, India, ²Department of Anesthesia, University of Edinburgh, UK

Background: Traumatic brain injury (TBI) is a major cause of death and disability across the globe. Raised intracranial pressure (ICP) has been the major detrimental factor while dealing with management of head trauma. Hypothermia has been shown to reduce ICP. Hence, our study planned to see the effect of hypothermia (32–35°C) for ICP reduction after TBI. **Methodology:** In this prospective randomised controlled trial, adult patients with primary closed TBI with raised ICP >20 mmHg for ≥5 min after first line treatments and with no obvious reversible cause, ≤10 days from the initial head injury, with core temperature ≥36°C (at the time of randomisation) and with an abnormal computed tomography scan were randomised to either hypothermia (32–35°C) or normothermia group. **Results:** A total of 27 patients were randomised, 14 in

hypothermia group and 13 in normothermia group. The mean age of the patients was 35.29 and 26.85 years, the mean ICP at the time of randomisation was 22.65 and 24.05 mmHg in hypothermia and control group, respectively. The Glasgow coma score at admission was 6.9 in the two groups. Of 13 patients in the hypothermia group, 4 patients were enrolled following decompressive craniectomy and raised ICP responded well to induction of hypothermia. In rest of the patients in hypothermia group, 5 were managed with hypothermia alone whereas 4 required decompressive surgery. Two patients in each group developed pneumonia. There were no coagulation abnormalities in either group. Mean duration of Intensive Care Unit and hospital stay was 10.15 days, 9.5 days and 19.5, 18.57 days in normothermia and hypothermia group, respectively. **Conclusions:** Hypothermia can be used safely as adjunct to other modalities for controlling ICP in severe head injury patients.

ISNACC-S-22

Craniopharyngioma surgery and various perioperative factors influencing its outcome: A prospective observational study

Ranadhirmitra, Hemanshu Prabhakar, Parmod Bithal, Girija Rath

Department of Neuroanesthesiology, All India Institute of Medical Sciences, New Delhi, India

Introduction/Purpose: The incidence of craniopharyngioma is stated around 1.3/million person-years whereas the incidence is higher in India with 10.2%. This current study is a prospective observational attempt to determine perioperative factors that affect the outcome of patients undergoing surgery for craniopharyngioma. The primary objective being duration of hospital and Intensive Care Unit (ICU) stay and Glasgow outcome scale (GOS) at discharge while the secondary objective being quality of life at 3-month and 6-month post-surgery. **Methods:** All patients aged 5 years and above belonging to either sex scheduled for elective craniopharyngioma surgery from 1st April 2014 to 31st March 2015 in Cardio-Neuro Centre of All India Institute of Medical Sciences, were included in the study while patient's or guardian's refusal or redo-surgery were excluded. The demographics, baseline characteristics (admission Glasgow coma scale, tumour size, hormonal status, location, hydrocephalus, hypothalamic involvement), intraoperative data (anaesthesia and surgery related), GOS at discharge and post-operatively quality of life assessed using the health utility indices (2/3) for a period up to 6 months after surgery were collected. **Results:** Twenty-two patients were included in the study. The median duration of hospital and ICU stay were 17 days (6–64) and 3.5 days (1–25), respectively. The median