Table 2: Intubation score

Criteria	Evaluation	Score	Group D n (%)	Group PF n (%)	P	Mann-Whitney U-test/P
Anesthesia quality	Asleep, deep sedation	2	19 (100)	20 (100)	1.000	136.000/0.039*
	Slight, resistance	0	0	0		
Vocal cords relaxation	No	0	0	0	0.064	
	In part, middle	1	3 (15.8)	0 (0)		
	Relaxed, open	2	16 (84.2)	20 (100)		
ETT tolerance	Bad, disturbing: Cough, swallowing	0	0	0	0.047*	
	Middle; coughing or swallowing not disturbing the procedure	2	7 (36.8)	2 (10)		
	Good, no coughing or swallowing	4	12 (63.2)	18 (90)		

ETT: Endotracheal tube

Fentanyl combination provided better AFOI condition than dexmedetomidine with shorter intubating time, better intubation score and post-operative profile with SAYGO technique. Both BIS and observer's assessment of awareness/sedation OAA/S are reliable indicators of sedation. Blinded could not be possible and small sample size were the limiting factors.

ISNACC-S-25

A retrospective analysis of pre-operative management of patients presenting for awake craniotomy: Sree Chitra Tirunal Institute for Medical Sciences and Technology experience

Nilima R. M., S. Manikandan, Smita V., Mathew Abraham¹

Division of Neuroanesthesiology, Sree Chitra Tirunal Institute for Medical Sciences and Technology, ¹Division of Neurosurgery, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, Kerala, India

Introduction: Awake craniotomy is performed when the lesions (usually tumour/seizure) are located close to the functional areas of the brain. However, awake craniotomy can pose challenges to the entire team of anaesthesiologist, surgeons and patients as well. We have retrospectively evaluated our experience in the management of these patients. **Materials and Methods:** Institutional Ethics Committee approval and patient consent waiver were obtained. Anaesthetic charts of all the patients who underwent the awake craniotomy in our institute from 2000 to August 2015 were analysed. **Results:** A total of 45 patients underwent awake craniotomy in the above period. The mean age of the patients was 33.4 years, male:female was 29:16. Initial presentation was seizure in 22 patients, speech disturbance in 11, sensory

in 4 and motor symptoms in 14. Premedication consisted of T diazepam (17), clonidine (16), pethidine (3) and haloperidol (9). One patient had airway obstruction and as converted to general anaesthesia. Rest of the patients underwent successful awake craniotomy. Scalp block was administered in 35, laryngeal mask airway was used in 3 (asleep awake asleep) technique. For maintenance anaesthesia, propofol with fentanyl infusion was given in 34 patients, dexmedetomidine in 3, midazolam in 2 and each pethidine and haloperidol in 2 patients. Intro complications were high end-tidal carbon dioxide (EtCO₂) in 6, tachycardia in 10 patients, hypertension in 25 patients, bradycardia in 5, hypotension in 6 patients. Pre-operative seizures occurred in 10 patients, brain bulge in 3 patients and intraoperative pain in 5 patients. **Conclusion:** Awake craniotomy is a challenging situation. Our experience showed that propofol with fentanyl infusion combined with scalp block was popular. However, in patients with dexmedetomidine, haemodynamic problems were less compared to propofol fentanyl group but the numbers were small to make a useful comparison.

ISNACC-S-26

Cerebral vascular effects of loading dose of dexmedetomidine: Transcranial colour Doppler study

Soumya Madhusudhan, Arulvelan A., Manikandan S.

Department of Anaesthesia, Division of Neuroanesthesia, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, Kerala, India

Background: Dexmedetomidine has been widely used in critical care settings because of its property of maintaining stable haemodynamics and inducing conscious sedation.

The use of dexmedetomidine is in increasing trend particularly in patients with neurological disorders. A very few studies have focused on the cerebral haemodynamic effects of dexmedetomidine. This study is aimed to address this issue. **Methods:** Thirty patients without any intracranial pathology were included in this study. Middle cerebral artery flow velocity (FV) obtained from transcranial colour Doppler was used to assess the cerebral haemodynamic indices. Mean FV (mFV), pulsatility index (PI), cerebral vascular resistant index (CVRi), estimated cerebral perfusion pressure (eCPP) and zero flow pressure (ZFP) were calculated bilaterally at baseline and after infusion of injection dexmedetomidine 1 mcg/kg over 10 min. **Results:** Twenty-six patients completed the study protocol. After administration of loading dose (LD) of dexmedetomidine, mFV and eCPP values were significantly decreased in both hemisphere ($P < 0.05$); PI, CVRi and ZFP values showed significant increase ($P < 0.05$) after dexmedetomidine infusion. **Conclusion:** Increase in PI, CVRi and ZFP suggests that there is possibility of increase in distal CVR with LD of dexmedetomidine. Decrease in mFV and eCPP along with increase in CVR may lead to decrease in cerebral perfusion. This effect can be exaggerated in patients with pre-existing neurological illness. Further studies are needed to evaluate the effect of dexmedetomidine on various other pathological conditions involving brain such as traumatic brain injury and vascular malformations.

ISNACC-S-27

Cerebral oxygenation during electroconvulsive therapy

Kaushik Theerth, Tanmay Jadhav, Sriganesh Kamath, Madhusudan Reddy

Department of Neuroanaesthesia, National Institute of Mental Health and Neurosciences, Bengaluru, Karnataka, India

Background: Near-infrared spectroscopy is a novel monitor to detect cerebral oxygenation (rSO_2). A very few studies have analysed the effect of electroconvulsive therapy (ECT) on rSO_2 . We hypothesised that atropine pre-medication related activation of systemic and cerebral haemodynamics will exacerbate ECT-induced increase in rSO_2 . This study aimed to assess the effect of atropine and no atropine pre-medication during ECT on rSO_2 and correlate this with haemodynamics and peripheral oxygen saturation. **Methodology:** Psychiatric patients aged between 18 and 60 years requiring modified ECT for their illness were included in this 6-month study. This was a randomised crossover trial, in which patients served as their own controls. For the second ECT session, patients were randomised to receive either 0.01 mg/kg atropine or no atropine preceding anaesthesia.

Anaesthesia was induced by thiopentone 3 mg/kg and muscle relaxation was achieved with suxamethonium 0.5 mg/kg. For the third ECT, the patients were crossed over. Heart rate, blood pressure, SpO_2 and rSO_2 values were recorded at baseline and every 15 s from ECT stimulus till 5 min. **Results:** Data from 41 patients were analysed. There was no difference in baseline rSO_2 in patients with different psychiatric diagnoses. rSO_2 increased consistently in both groups after ECT which did not return to baseline at 5 min. Increase in rSO_2 was independent of increase in blood pressure. rSO_2 was lower in patients who developed desaturation during ECT. **Discussion/Conclusion:** On the contrary, no immediate desaturation after ECT stimulus was seen. rSO_2 increased consistently following ECT in both the groups. Systemic desaturation resulted in lower rSO_2 . Future studies should investigate whether rSO_2 changes associated with therapeutic seizures contributes to post-ECT cognitive changes.

ISNACC-S-28

Evaluation of usefulness of a point of care haemoglobin device in emergency neurosurgery

Pavithra V., Sudhir V.

Department of Neuroanesthesia, National Institute of Mental Health and Neurosciences, Bengaluru, Karnataka, India

Introduction: Blood transfusion and anaemia in traumatic brain injury adversely affect the outcome and also increase length of stay and mortality. Laboratory haemoglobin (Hb) values are considered primary indicators and gold standard to guide blood transfusion, but lab results take time and repetitive measurements may not be feasible. Rapid and accurate measurements of Hb can guide warranted blood transfusions and avoid unnecessary transfusions in emergency neurosurgery. In this regard, point of care testing devices for Hb can be of great use. **Design:** Prospective study to assess the accuracy of Hb values obtained through HemoCue Hb, Sweden analyser compared to laboratory Hb reports in patients undergoing emergency neurosurgery and Intensive Care Unit (ICU). **Conduct:** The study was conducted in 50 patients undergoing emergency neurosurgery at National Institute of Mental Health

Table 1: The differences in haemoglobin values obtained through the device and lab analysis are not significant

Pair 1 haemoglobin device values-haemoglobin lab values			
Difference mean	t	df	P
-0.29	-0.266	48	0.79