Epidemiological and Clinical Features of Diaper Dermatitis in Infants at the Yaounde Gynaeco-Obstetric and Pediatric Hospital, Cameroon

Andreas Chiabi^{1,2} Jenny Christelle Ngamgo Kamdem³ Anita Grâce Nkoro¹ Hippolyte Siyou¹
Evelyn Mah^{1,2} Félicitée Dongmo Nguéack^{1,2} Seraphin Nguéack^{1,2} Fru Angwafo III^{1,2}

¹Yaounde Gynaeco-Obstetric and Pediatric Hospital, Yaounde, Cameroon

²Department of Pediatrics, Faculty of Medicine and Biomedical Sciences, University of Yaounde I, Yaounde, Cameroon

³Université de Montagnes, Bangangté, Cameroun

Address for correspondence: Andreas Chiabi, MD, Yaounde Gynaeco-Obstetric and Pediatric Hospital, Cameroon, Faculty of Medicine and Biomedical Sciences, University of Yaounde I, P.O. Box 4362, Yaounde, Cameroon (e-mail: andy_chiabi@yahoo.co.uk).

J Child Sci 2018;8:e46–e49.

Abstract

Diaper dermatitis is defined as an inflammatory reaction localized on the skin that is usually covered by the diaper (genitals, buttocks, and anus). The aim of this study was to determine the prevalence of diaper dermatitis and to describe the clinical presentation and practices related to hygiene in infants with diaper rash at the Yaounde Gynaeco-Obstetric and Pediatric hospital. It was a descriptive cross-sectional study undertaken in 1,004 mother–children pairs. Data were collected using structured questionnaires during face-to-face interviews with the parents. The prevalence of diaper dermatitis was 18.4%, and infants were aged 7 days to 24 months with a mean age of 5.91 months. The most represented age group was 0 to 6 months (67.6%). Infants living in urban areas (59.5%), of unemployed mothers or guardians (30.3%), and those of mothers living as a couple (75.1%) were most represented in our study sample. The majority of infants used both disposable and reusable diapers (48.6%). In addition, 72.7% mothers used palm kernel oil, and 27.3% rubbed other ointments before changing the baby's diapers. The most frequent clinical form was the chaffing form (48.1%). Healthcare personnel should inform, educate, and communicate with mothers on body hygiene of their babies to avoid this disorder, which can cause severe discomfort and sometimes lead to complications.

Keywords

- diaper rash
- infant
- epidemiology
- clinical features

Introduction

The terms “diaper rash” or “nappy rash,” or “diaper dermatitis” are used to designate any skin irritation in infants appearing on the area covered by the diaper.^{1,2} It is one of the most common disorders in infants and newborns, accounting for 10 to 20% of all skin diseases.³ The precise and specific etiology of diaper dermatitis has not yet been determined.⁴ However, irritation

with skin maceration due to prolonged or frequent contact with soiled diapers is often indicated.⁵

Knowing the epidemiological and clinical features of this disorder is important for effective prevention. This study was thus undertaken at the Yaounde Gynaeco-Obstetric and Pediatric hospital, which is a mother–child hospital, to assess the epidemiological and clinical aspects of this disorder in an urban Cameroonian context.

received
April 15, 2018
accepted after revision
July 25, 2018

DOI <https://doi.org/10.1055/s-0038-1669424>.
ISSN 2474-5871.

Copyright © 2018 Georg Thieme Verlag
KG Stuttgart · New York

License terms

Materials and Methods

This is a descriptive cross-sectional study, at the vaccination and outpatient consultation units of the Yaounde Gynaeco-Obstetrics and Pediatric hospital. The study was conducted from February 1, 2017, to June 30, 2017 (5 months).

We consecutively enrolled all infants aged less than 24 months, brought for vaccination or pediatric consultation, over the study period. The exclusion criteria were: children with a life-threatening emergency requiring immediate hospitalization and children whose parents refused to participate in the study.

The variables noted were: sex and age of the infant, age of the mother, frequency of change of the diapers, frequency of baths, sociodemographic data of the mothers and infants, hygiene practices performed for the infant, and clinical features.

All infants enrolled at the vaccination and/or outpatient consultations were undressed and examined. At the end of the physical examination, the elementary lesions observed, as well as their location, permitted us to describe the different types of diaper dermatitis if found. The dermatitis was noted as “W” type diaper dermatitis or chaffing dermatitis, or frictional dermatitis, if erythema was found on the inner part of the thighs and extends in girls to the labia majora and in boys to the inguinoscrotal region, not affecting the inguinal folds. It was noted as “Y” type dermatitis or *Candida albicans* dermatitis when the erythema was present on the inguinal and intergluteal folds, and irritant contact dermatitis when the erythema affected the intergluteal folds, the gluteal, perianal, and the pubic regions.

Data were entered and analyzed with the software Epi info version 3.5.4.

Authorization and ethical clearance to perform the study were obtained from the administration of the hospital. Informed consent to participate in the study was obtained from the parents before enrollment.

Results

During the study period, we received 1,155 infants aged from 7 weeks to 24 months. We examined 1,004 infants, of which 185 had a diaper rash, giving a prevalence of 18.4%.

The sociodemographic profile of infants is summarized in ►Table 1.

Table 1 Sociodemographic profile of the infants (total number of patients: 185)

	Number	Percentages (%)
Age group		
0–6 months	125	67.6
6–12 months	48	25.9
12–18 months	12	6.5
Sex		
Male	91	49.2
Female	94	50.8

The most represented age group was 0 to 6 months. The mean age was 5.91 months with extremes from 7 weeks to 24 months. Out of 185 infants with diaper rash, 50.8% were female, and 49.2% were male giving a sex ratio of 0.97. The sociodemographic profile of parents is summarized in ►Table 2.

In our study population, 59.5% resided in urban areas (►Table 3). The most represented maternal age group was between 20 and 30 years, i.e., 51.9%. Infants whose parents lived together were represented in 75.1% of the cohort. Regarding the practices of hygiene care, 48.6% used both types of diaper (reusable and disposable). Infants having a frequency of diaper change less than 6 times a day were the most affected by diaper rash (76.8%), and 90.3% were not cleaned after passing urine. The most frequently used topical treatment was palm kernel oil (72.7%). Wipes were the most used cleaning agents for the skin at 75.7%. The most frequent clinical form of diaper dermatitis was the “W” or chaffing or frictional form in 89 (48.10%) cases. The irritant contact dermatitis and the “Y” or *Candida* dermatitis occurred in 52 and 44 cases, respectively, i.e., 28.1% and 23.8%, respectively.

Discussion

In our study, 18.4% of infants had diaper rash, which is lower than that found by Li et al in 2012 in China, which was 43.8%.⁶ Indeed, the study by Li et al focused on the study of risk factors in infants from 1 to 24 months in China and was conducted in five different hospitals, while our study was conducted in a single hospital.

The age of the infants ranged from 1 week to 24 months with a mean age of 5.91 months, and the most represented age group was 0 to 6 months. Our results are similar to those of Wanjiku et al in 2016 in Kenya.⁷ Infants in the 6 to 12 months age group were predominant in the study of Biranjia and Pandamikum in 2015 in the Mauritius island,⁸ whose study focused on diaper rash in babies less than 36 months of age. In the study in Mauritius, infants who

Table 2 Sociodemographic profile of the mothers or guardians (total number of patients: 185)

	Number	Percentages (%)
Age group of the mother or guardian		
<20 years	8	4.3
20–30 years	96	51.9
30–40 years	77	41.6
40–50 years	4	2.2
Area of residence		
Urban	110	59.5
Rural	75	40.5
Marital status		
Living as a couple	139	75.1
Living alone	46	24.9

Table 3 Hygiene practices (total number of patients: 185)

	Number	Percentages (%)
Diaper type		
Disposable	72	38.9
Reusable	23	12.4
Both	90	48.6
Frequency of change		
<6 times a day	142	76.8
≥6 times a day	43	23.2
Do you clean the child when he urinates?		
Yes	18	9.7
No	167	90.3
Cleaning agents for the skin		
Wipes	140	75.7
Wet diapers	15	8.1
Toilet paper	2	1.1
Topical treatments		
Palm kernel oil	133	72.7
Ointment ^a	50	27.3
Cream ^b	13	7.1
Powder ^c	13	7.1
Olive oil	7	3.8
Milk	3	1.6
Perfume	3	1.6

^aAntibiotics, anti-fungal agents, and anti-inflammatory.^bTraditional and pharmaceutical.^cPharmaceutical and baby powders.

had had at least one previous episode of diaper rash were enrolled, unlike in ours.

Infants residing in urban areas were the most represented. Our results are contrary to those of Li et al in 2012 in China⁶ who noted that most of their patients were from the rural areas. This difference could be due to the study setting with respect to the residence of the mother. Patients will tend to go to the nearest hospital whatever the setting.

In our study, the most represented age group of the mothers was 20 to 29 years of age. This could be explained by the fact that it is within this age range that most women deliver, and most do not have experience in baby hygiene.

The infants of mothers living as couples were the most represented in our study population. These results are similar to those of Wanjiku et al in 2016 in Kenya.⁷ A possible explanation is that women living as part of a couple are more likely to have less time for the hygiene of the infants since they must also cater for the rest of the family.

Regular wearing of diapers was noted in all infants. For hygiene and body care of the infants, most of them used both types of diapers (disposable and reusable), followed by those using the disposables. Several mechanisms could explain the imputability of the nappies in the occurrence of diaper rash—

if the diapers are not properly used, an occlusive effect may occur leading to an increase in skin hydration by inhibiting skin evaporation and transdermal water losses and skin pH.^{9,10}

Diapers, especially the disposable ones, can be irritative due to the poor fabrication quality of the napkins. Reusable diapers could also be irritative if they are of a plastic fabric. Studies conducted by Campbell et al and cited by Moulounguet-Michau,⁹ noted that disposable diapers were less irritating than reusable diapers.

Average frequency of diaper change was 5.76 times per day. In our study, infants whose frequency of diaper change was less than 6 times per day were the most affected. Our results are similar to those noted by Li et al in 2012 in China.⁶ Some authors have noted that an infant urinates 10 to 20 times a day and passes stool 3 to 4 times per day,¹¹ suggesting that on several occasions, the infant had to urinate and stay with its diaper wet. Moulounguet-Michau et al noted that one of the factors which determines the appearance of diaper dermatitis is the time of contact of the skin with the irritating factor.⁹ Diaper change must therefore be done immediately after each episode of passing urine.

Few mothers clean the buttocks of their infants after they had passed urine, not knowing that urine and stools play a very important role in the occurrence of diaper rash. Indeed, the mixture of stools and urine in the diaper will cause a transformation (induced by the bacteria present in the stool) of urea into ammonia, which will increase the pH of the skin in this area. This rise in pH causes a reactivation of the digestive enzymes, which irritate the baby's skin and increases permeability of the skin.^{4,12}

The most frequently used body oil was palm kernel oil (72.7% of cases). This could be due to the fact that this product is locally available and is most often recommended by health personnel in our health facilities for its hypoallergenic properties; nevertheless, the frequency of diaper rash was high in infants using this oil, although the reason for this finding still remains unclear.

The "W" type dermatitis or chaffing dermatitis was the most frequently encountered dermatitis in 48.1% of our patients. According to Moraillon¹¹ and Piérard-Franchimont,¹³ the most frequent diaper rash lesions have the "W" topography or frictional dermatitis, due to mechanical reaction in which occlusion and maceration play a key role. Diaper dermatitis complicated with the presence of *C. albicans* presents a beefy, red, raised-edge skin rash with pinpoint satellite lesions, which often extend into the folds of the skin in the diaper area. Whenever diaper dermatitis fails to respond to treatment, *Candida* diaper dermatitis should be considered as an alternate cause of the presenting rash. Because *C. albicans* is contained in the lower intestine of infants, fecal matter presents the primary source for *Candida* diaper dermatitis. In some instances, in addition to *Candida* diaper dermatitis, oral dermatitis (thrush) may be present, and the infant may require treatment with an oral antifungal agent.¹⁴

Therefore, healthcare personnel should inform, educate, and communicate with mothers on hygiene and body care of

their babies to avoid this disorder that can lead to discomfort and reduced well-being of the infant.

Conflict of Interest

None.

References

- 1 Stevenson J. Getting to the bottom of nappy rash. *Nurse Prescribing*. 2011;9(01):25–27
- 2 Philipp R, Hughes A, Golding J; ALSPAC Survey Team. Avon Longitudinal Study of Pregnancy and Childhood. Getting to the bottom of nappy rash. *Br J Gen Pract* 1997;47(421):493–497
- 3 Ravanfar P, Wallace JS, Pace NC. Diaper dermatitis: a review and update. *Curr Opin Pediatr* 2012;24(04):472–479
- 4 Stamatias GN, Tierney NK. Diaper dermatitis: etiology, manifestations, prevention, and management. *Pediatr Dermatol* 2014;31(01):1–7
- 5 Lagier L, Mazereeuw-Hautier J, Raffin D, Beneton N, Lorette G, Maruani A; Société française de dermatologie pédiatrique. Diaper dermatitis [article in French]. *Ann Dermatol Venerol* 2015;142(01):54–61; quiz 53, 62
- 6 Li CH, Zhu ZH, Dai YH. Diaper dermatitis: a survey of risk factors for children aged 1–24 months in China. *J Int Med Res* 2012;40(05):1752–1760
- 7 Wanjiku A, Ng'ang'a Z, Mbakaya C, Habtu M. Socio-demographic and economic determinants of diaper dermatitis among children aged 0–24 months at Mbaghati District Hospital, Kenya: a cross sectional study. *Int J Health Sci Res*. 2016;6:239–247
- 8 Biranjia-Hurdoyal S, Pandamikum L. A study to investigate the prevalence of nappy rash among babies aged 0 to 36 months old in a tropical country. *Aust J Dermatol* 2015;2(02):1040
- 9 Moulouguet-Michau. Dubertret. *Le Manuel du Resident - Pédiatrie II. Dermatite de siège*. Édition Tsunami. 2009:3455–3465
- 10 Longhi F, Carlucci G, Bellucci R, di Girolamo R, Palumbo G, Amerio P. Diaper dermatitis: a study of contributing factors. *Contact Dermat* 1992;26(04):248–252
- 11 Moraillon I. L'érythème fessier du nourrisson en pratique quotidienne. *Dermatologie Pratique*. 1994;138:1–2
- 12 Beguin A. L'érythème fessier: toujours d'actualité? *Arch Pediatr* 2006;13:6–9
- 13 Piérard-Franchimont GE. La dermatite des langes. *Revue Dermatologie Tropicale*; 1993:605
- 14 Merrill L. Prevention, treatment and parent education for diaper dermatitis. *Nurs Womens Health* 2015;19(04):324–336, quiz 337