Importance of Individual Elements for Perineal Protection in Childbirth: An Interventional, Prospective Trial

Ole Bredahl Rasmussen, MD¹ Annika Yding, CM, MLP¹ Finn Lauszus, MD, PhD¹
Charlotte Sander Andersen, CM¹ Jacob Anhøj, MD, DIT² Jane Boris, MD¹

¹ Department of Obstetrics and Gynaecology, Herning Regional Hospital, Herning, Denmark

² Centre of Diagnostic Investigation, University of Copenhagen, Copenhagen, Denmark

Address for correspondence Ole Bredahl Rasmussen, MD, Department of Obstetrics and Gynaecology, Herning Regional Hospital, Gl. Landevej 61, 7400 Herning, Denmark (e-mail: olebr@dadlnet.dk).

Am J Perinatol Rep 2018;8:e289–e294.

Abstract

Objective To analyze the association between each element of a hands-on intervention in childbirth and the incidence of obstetric anal sphincter injuries (OASIS).

Study Design We conducted a prospective, interventional quality improvement project and implemented a care bundle with five elements at an obstetric department in Denmark with 3,000 deliveries annually. We aimed at reducing the incidence of OASIS. In the preintervention period, 355 vaginally delivering nulliparous women were included. Similarly, 1,622 nulliparous women were included in the intervention period. The association of each element with the outcome was estimated using a regression analysis.

Results The incidence of OASIS went down from 7.0 to 3.4% among nulliparous women delivering vaginally ($p = 0.003$; relative risk = 0.48; 95% confidence interval [CI]: 0.30–0.76). Number needed to treat was 28. Logistic regression analysis showed that using hand on the head of the child significantly reduced the risk of OASIS (odds ratio = 0.28; 95% CI: 0.14–0.58).

Conclusion Using a quality improvement framework, we documented the individual elements of the intervention. Hand on the infant's head reduced the risk of OASIS.

Keywords

- vaginal delivery
- obstetric anal sphincter injury
- hands-on
- quality improvement

An obstetric anal sphincter injury (OASIS) is a serious adverse outcome of vaginal delivery.¹ The consequences may be lifelong and severe.^{2,3} Any measure to reduce the risk during vaginal delivery should be taken into consideration. Nulliparous women have a relatively high risk of OASIS and, therefore, the potential gain of lowering the incidence in this group is high. Two recent studies from the United States found an incidence in nulliparous women delivering vaginally was 9.7 and 8.3%.^{4,5} The odds ratios (OR) for nulliparity were 4.8 and 2.3, respectively. In a national survey from the United Kingdom, the median incidence of OASIS in a similar cohort was 6.1% in 2009.⁶

Several interventional before and after trials have shown a reduction in the rate of OASIS after implementing a hands-on technique in childbirth.^{7–11} In contrast to this, two recent metaanalyses^{12,13} including four randomized controlled trials

(RCTs) did not show a benefit of the hands-on technique.^{14–17} Neither the observational studies nor the RCTs document actual adherence to the specific elements of the intervention.

Using a quality improvement framework with documentation of the use of the elements of the care bundle, we showed that it was feasible to implement a standardized hands-on technique in vaginal delivery, thereby significantly reducing the incidence of OASIS from 7 to 3.4%.¹⁸ In the present study, we aimed at analyzing the association between each element of the intervention and the outcome.

Materials and Methods

In the previously published article covering the quality improvement methodology of the trial, we included

received
June 24, 2018
accepted after revision
September 14, 2018

DOI <https://doi.org/10.1055/s-0038-1675352>.
ISSN 2157-6998.

Copyright © 2018 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 584-4662.

License terms

descriptive statistics on the adherence to process indicators and the corresponding chronology of the outcome measure.¹⁸ The present study includes original descriptive data on the cohorts before and after initiation of the intervention, and analysis of the correlation among the five different elements of the care bundle and the outcome, including a regression analysis.

Around 2,800 to 3,000 women are delivering in our department yearly. We refer very preterm deliveries (before gestational week 28 + 0) and pregnant women with complicated medical illnesses to a tertiary hospital. Our study cohort included all nulliparous women delivering vaginally (including home births with an attending midwife from our department). The baseline/preintervention period was from January through May 2013, while the intervention period was from June 2013 through March 2015.

The main outcome measure was the incidence of OASIS, that is, grade 3 or 4 sphincter ruptures, as defined in the international classification¹⁹ and in correspondence with the reVITALize definition (<https://www.acog.org/About-ACOG/ACOG-Departments/Patient-Safety-and-Quality-Improvement/reVITALize>. Retrieved on April 25, 2018). The definition and the ascertainment of a case with sphincter rupture were the same before and during the intervention period. All cases were ascertained and sutured by a doctor in our department.

The intervention consisted of implementation of a bundle of care and a certification process. The bundle of care comprised four elements: (1) communication settled, that is, a shared decision with the woman about not to push when delivering the head of the child to slow down the speed of delivery, (2) visible brim of the perineum during delivery of the head, (3) hand on the infant's head to slow down the speed of delivery, and (4) manual protection of the perineum during delivery of the head, facilitating the naturally occurring extension of the head. Thus, we used five process indicators in total, including the certification process in which all midwives and doctors were informed about the project and the intervention and were supervised in the use of it. Process and outcome data were collected prospectively after each delivery. Information on clinical characteristics and outcome data on OASIS from the preintervention period as well as data from women with missing data in the intervention period were retrieved from existing electronic records.

The preintervention and intervention periods were compared with respect to the distribution of cohort characteristics, interventions, and the OASIS incidence using a statistical proportions test. The null hypothesis was that the proportions in the two periods were the same. Finally, we performed a logistic regression analysis using stepwise backward elimination of the least significant elements to identify which, if any, of the five elements of the intervention affected the risk of OASIS. A two-sided p -value < 0.05 was considered the level of significance.

Data were analyzed with Open Epi, version 2.3.1 and R Statistical Software v. 3.2.1 (R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, 2015. URL: <http://www.R-project.org>) and graphs were created using the R package qicharts v. 0.2.0 (Anhoej J. QI charts: Quality Improvement Charts. R package version 0.2.0, 2015. URL: <http://CRAN.R-project.org/package=qicharts>).

The study was approved by the Danish Data Protection Agency (2007-58-0010) while approval from the local ethics committee was not found relevant by the committee (1-10-72-127-13). Patient information was anonymized before analysis. No funding was received.

Results

In the preintervention period, 460 nulliparous women gave birth, and in the intervention period, 2,019. The rates of cesarean birth were 22.8 and 19.7%, respectively ($p = 0.13$). Thus, 355 nulliparous women gave birth vaginally in the preintervention period and 1,622 in the intervention period (► **Table 1**).

Registration charts with information on adherence to the bundle of care were collected from 1,594 deliveries (98.3%) and were missing in 28 deliveries (1.7%). None of the missing cases had OASIS. Therefore, they were excluded from the analysis of process indicators but included in the OASIS rate analysis.

None of the characteristics of the cohorts from the preintervention and intervention periods differed, including age at birth, gestational age, birth weight, the incidence of vacuum extraction, and episiotomy. The rate of the different combinations of vacuum extraction and episiotomy did not differ either (► **Table 2**).

Table 1 Mode of delivery in nulliparous women in preintervention and intervention periods, Herning Regional Hospital, Denmark

Mode of delivery	Preintervention N = 460		Intervention N = 2,019		p-Value
	Not in labor (n, %)	In labor (n, %)	Not in labor (n, %)	In labor (n, %)	
SVD		296 (68.8)		1,362 (71.3)	0.32
OVD		59 (13.7)		260 (13.6)	0.94
CB	30	75 (17.4)	108	289 (15.1)	0.23
Total	30 (6.5)	430 (100)	108 (5.3)	1,911 (100)	0.32

Abbreviations: CB, cesarean birth; OVD, operative vaginal delivery; SVD, spontaneous vaginal delivery.

Table 2 Cohort characteristics in nulliparous women delivering vaginally in preintervention and intervention periods, Herning Regional Hospital, Denmark

Characteristic	Preintervention N = 355		Intervention N = 1,622		p-Value
	N	% of column	n	% of column	
Age at birth (y)					
17–24	92	25.9	376	23.2	0.47
25–29	166	46.8	763	47.0	
30–42	97	27.3	483	29.8	
Gestational age (wk + d)					
26 + 1–37 + 6	25	7.1	173	10.7	0.21
38 + 0–39 + 6	113	31.9	520	32.1	
40 + 0–40 + 6	119	33.6	505	31.2	
41 + 0–42 + 4	97	27.4	423	26.1	
Unknown	1		1		–
Birth weight (g)					
860–2,999	53	14.9	297	18.3	0.48
3,000–3,499	138	38.9	590	36.4	
3,500–3,999	121	34.1	550	33.9	
4,000–5,040	43	12.1	185	11.4	
Episiotomy	34	9.6	136	8.4	0.46
OVD	67	18.9	260	16.0	0.18
OVD with episiotomy	13	3.7	44	2.7	0.34
OVD without episiotomy	54	15.2	216	13.3	0.35
SVD with episiotomy	21	5.9	92	5.7	0.84
SVD without episiotomy	267	75.2	1,270	78.3	0.21

Abbreviations: OVD, operative vaginal delivery; SVD, spontaneous vaginal delivery.

In the birth population with vaginal delivery and mixed parity, the rate of OASIS changed significantly from 3.5 to 1.9% ($p = 0.004$). In the study group of nulliparous women with vaginal delivery, the rate of OASIS decreased significantly from 7.0% in the preintervention period to 3.4% in the intervention period ($p = 0.003$, relative risk = 0.48, 95% confidence interval

[CI]: 0.3–0.76). Number needed to treat was 27.8 or 28 nulliparous women delivering vaginally. The rate of OASIS over time in the overall study group is shown in ►Fig. 1. Rates of OASIS in subgroups of the study population are shown in ►Table 3.

The OASIS incidence did not increase in any of the subgroups. In the following subgroups, the incidence decreased

Fig. 1 Rate of OASIS per month in nulliparous women with a vaginal delivery before and after the start of the intervention. P-chart. Horizontal line showing the extended baseline from the preintervention period. Vertical line showing the start of the intervention period, Herning Regional Hospital, Denmark. OASIS, obstetric anal sphincter injury.

Table 3 Incidence of OASIS in subgroups of nulliparous women delivering vaginally in preintervention and intervention periods, Herning Regional Hospital, Denmark

Subgroups	Preintervention			Intervention			p-Value ^a	RR, 95% CI
	n	OASIS (n)	Incidence %	N	OASIS (n)	Incidence %		
Incidence of OASIS	355	25	7.0	1,622	55	3.4	0.003 ^a	0.48, 0.30–0.76
Age at birth (y)								
17–24	92	8	8.7	376	5	1.3	0.001 ^a	0.15, 0.05–0.46
25–29	166	9	5.4	763	32	4.2	0.48	0.77, 0.38–1.59
30–42	97	8	8.2	483	18	3.7	0.07	0.45, 0.20–1.01
Gestational age (wk + d)								
26 + 1–37 + 6	25	2	8.0	173	2	1.2	0.09	0.14, 0.02–0.98
38 + 0–39 + 6	113	3	2.7	520	14	2.7	0.97	1.01, 0.30–3.47
40 + 0–40 + 6	119	13	10.9	505	19	3.8	0.004 ^a	0.34, 0.18–0.68
41 + 0–42 + 4	97	6	6.2	423	20	4.7	0.55	0.76, 0.32–1.85
Unknown	1	1	100	1	0	0	–	
Birth weight (g)								
860–2,999	53	2	3.8	297	7	2.4	0.55	0.62, 0.13–2.93
3,000–3,499	138	2	1.4	590	10	1.7	0.90	1.17, 0.26–5.28
3,500–3,999	121	12	9.9	550	19	3.5	0.006 ^a	0.35, 0.17–0.70
4,000–5,040	43	9	20.9	185	12	6.5	0.008 ^a	0.31, 0.14–0.69
Episiotomy								
– Episiotomy	321	20	6.2	1,486	46	3.1	0.01 ^a	0.50, 0.30–0.83
+ Episiotomy	34	5	14.7	136	9	6.6	0.16	0.45, 0.16–1.26
Mode of delivery								
SVD	288	17	5.9	1,362	33	2.4	0.004 ^a	0.41, 0.23–0.73
OVD	67	8	11.9	260	22	8.5	0.39	0.71, 0.33–1.52
SVD without episiotomy	267	15	5.6	1,270	29	2.3	0.007 ^a	0.41, 0.22–0.75
SVD with episiotomy	21	2	9.5	92	4	4.3	0.39	0.46, 0.09–2.33
OVD without episiotomy	54	5	9.3	216	17	7.9	0.72	0.85, 0.33–2.20
OVD with episiotomy	13	3	23.1	44	5	11.4	0.33	0.49, 0.14–1.79

Abbreviations: CI, confidence interval; OVD, operative vaginal delivery; RR, relative risk; SVD, spontaneous vaginal delivery.

Note: Age at birth: range 17 to 42 years; gestational age: range 26 + 1 to 42 + 4 weeks and days; birth weight: range 860 to 5,040 g.

^ap-Value < 0.05 is considered statistically significant.

significantly: women younger than 25 years, women delivering during the first week after term, women delivering infants of 3,500 g or more, women not having an episiotomy, women with spontaneous vaginal delivery, and the group with spontaneous vaginal delivery without an episiotomy. The incidence of OASIS in operative vaginal deliveries changed from 11.9 to 8.5% ($p = 0.39$). All operative vaginal deliveries were by vacuum extraction. We found one grade 4 sphincter rupture in the preintervention period and three in the intervention period. The incidences were 0.28 and 0.18%, respectively, which was not statistically different.

When analyzing the individual elements of the bundle of care, the lowest incidence of OASIS was 2.3% and was found in the group of women where a certified midwife used hand on the infant's head. The association of the single elements of the process indicators was analyzed further with a logistic regres-

sion analysis. After stepwise removal of nonsignificant elements from the regression analysis, only hand on head remained significant. The OR of OASIS with hand on head was 0.28 with 95% CI of 0.14 to 0.58 corresponding with a risk reduction of 72%. OR and 95% CI for each element: hand on the head 0.28; 0.14 to 0.58, certification 0.71; 0.41 to 1.25, visible perineum 0.85; 0.31 to 2.76, perineal support 1.1; 0.51 to 2.52, and settled communication 1.4; 0.51 to 4.84. The OR and CI for each of the individual elements for perineal protection are shown in **Fig. 2**.

Discussion

We successfully halved the incidence of OASIS in nulliparous women with vaginal delivery after implementing four elements of a hands-on technique and a certification process for

Fig. 2 Association of individual elements for perineal protection on the risk of obstetric anal sphincter injuries (odds ratios and 95% CIs). Regression analysis with stepwise removal of nonsignificant elements. Herning Regional Hospital, Denmark. CIs, confidence intervals; OASIS, obstetric anal sphincter injury.

all midwives and doctors. This reduction is in concord with other results after implementation of a perineal protection training program.^{7–11} Analysis of the elements in different combinations showed that hand on the head by a certified midwife was associated with the lowest incidence of OASIS. Hand on the head of the child resulted in a significant reduction in the risk of OASIS. The rates of cesarean birth and episiotomy were unchanged.

The main strength of this study is that it consisted of a prospective implementation of a standardized intervention in a quality improvement framework. Standardizing the intervention through the certification process may in itself contribute to the benefit of that intervention.²⁰ Using a quality improvement methodology provides control of the process since the use of each element is documented.

The strength of statistical association in the present study is of a clinically important size. The association was seen in different subgroups going in the same direction. We found a relevant temporality of the intervention and outcome relationship. A significant change in the outcome appeared after 3 months and has persisted since then. Presumably, the speed of delivery of the child and a calm approach to vaginal delivery is important for the risk of OASIS.²¹ This factor is addressed by the intervention. Prior to the present project, several initiatives were performed to address the relatively high incidence of OASIS in our department. A possible Hawthorne effect would likely have shown during that period. Thus, the Bradford Hill criteria support an assumption about a causal relationship between the intervention and the outcome.²²

One limitation to the study is that we used a historical cohort for comparison. The only deliberate change that took place during the study came from the intervention, and the cohort characteristics did not change between preintervention and intervention periods (–Table 2). Still, this represents a potential bias, since other differences in the study group may have evolved during the study time, and residual confounding cannot be excluded.

The diagnosis of OASIS is a well-known clinical challenge.²³ During the entire project, we had a focus on how

to diagnose a sphincter rupture for both midwives and doctors. We find it unlikely that the rate of missed diagnoses went up during the intervention period.

The number of women in the study is limited due to the prospective, interventional design. Thus, a type-2 error may be present and may explain the lack of other statistically significant associations between different elements of the intervention and the outcome.

Overreporting of the use of the different elements of the bundle of care may have biased the results. Especially in cases with OASIS, the care provider may feel a pressure to report adherence to all the elements. This recall bias will tend to reduce the association between the different elements and the outcome.

We noticed a high adherence to several of the process indicators from the start of the intervention period. In a before and after study, premature implementation of an intervention will tend to reduce the power.

An apparent limitation to the study is that the all-or-none adherence to the five elements of the intervention leveled out around 80%. Feedback from the midwives and doctors revealed that there were several reasons for this: fast deliveries with not enough time for communication about the proposed intervention, and deliveries where the woman opted for a water birth or other “alternative” birth positions. Also, midwife students and newly employed had to be certified during actual deliveries and therefore were not certified in their first deliveries. Thus, while reducing the association between the intervention and the outcome, at the same time, it strengthens the external validity of our findings, since these situations are part of daily life at a labor ward.

An RCT may also suffer from limitations to providing useful data for recommendation of an intervention.²⁴ The difficulty with blinding the intervention poses a possible bias in non-pharmacologic RCTs.²⁵ Furthermore, personal preferences by the birth attendant or the woman and clinical circumstances may present obstacles to implementing a specific intervention. A lack of difference in the interventions between the two arms in an RCT will reduce or eliminate the power to prove a difference. Irrespective of the design and the specific

intervention, documenting the use of the elements of the intervention will strengthen the evidence.

Applying a quality improvement framework using a specified and standardized care bundle provided a generalizable intervention. The risk of OASIS proved to be modifiable without increasing the rate of cesarean birth or episiotomy. Hand on the head of the child resulted in a significant reduction in the risk of OASIS. This may be applicable also in alternative birth positions and in fast deliveries.

Funding

No funding received.

Conflict of Interest

There is no conflict of interest.

References

- Turner CE, Young JM, Solomon MJ, Ludlow J, Benness C, Phipps H. Vaginal delivery compared with elective caesarean section: the views of pregnant women and clinicians. *BJOG* 2008;115(12):1494–1502
- Desseauve D, Proust S, Carlier-Guerin C, Rutten C, Pierre F, Fritel X. Evaluation of long-term pelvic floor symptoms after an obstetric anal sphincter injury (OASI) at least one year after delivery: a retrospective cohort study of 159 cases. *Gynecol Obstet Fertil* 2016;44(7-8):385–390
- Halle TK, Salvesen KÅ, Volløysaug I. Obstetric anal sphincter injury and incontinence 15-23 years after vaginal delivery. *Acta Obstet Gynecol Scand* 2016;95(08):941–947
- Meister MRL, Cahill AG, Conner SN, Woolfolk CL, Lowder JL. Predicting obstetric anal sphincter injuries in a modern obstetric population. *Am J Obstet Gynecol* 2016;215(03):310.e1–310.e7
- Ramm O, Woo VG, Hung Y-Y, Chen H-C, Ritterman Weintraub ML. Risk factors for the development of obstetric anal sphincter injuries in modern obstetric practice. *Obstet Gynecol* 2018;131(02):290–296
- Thiagamoorthy G, Johnson A, Thakar R, Sultan AH. National survey of perineal trauma and its subsequent management in the United Kingdom. *Int Urogynecol J Pelvic Floor Dysfunct* 2014;25(12):1621–1627
- Laine K, Skjeldestad FE, Sandvik L, Staff AC. Incidence of obstetric anal sphincter injuries after training to protect the perineum: cohort study. *BMJ Open* 2012;2(05):e001649
- Laine K, Pirhonen T, Rolland R, Pirhonen J. Decreasing the incidence of anal sphincter tears during delivery. *Obstet Gynecol* 2008;111(05):1053–1057
- Hals E, Øian P, Pirhonen T, et al. A multicenter interventional program to reduce the incidence of anal sphincter tears. *Obstet Gynecol* 2010;116(04):901–908
- Leenskjoeld S, Høj L, Pirhonen J. Manual protection of the perineum reduces the risk of obstetric anal sphincter ruptures. *Dan Med J* 2015;62(05):A5075
- Basu M, Smith D, Edwards R; STOMP project team. Can the incidence of obstetric anal sphincter injury be reduced? The STOMP experience. *Eur J Obstet Gynecol Reprod Biol* 2016;202:55–59
- Bulchandani S, Watts E, Sucharitha A, Yates D, Ismail KM. Manual perineal support at the time of childbirth: a systematic review and meta-analysis. *BJOG* 2015;122(09):1157–1165
- Aasheim V, Nilsen ABV, Reinart LM, Lukasse M. Perineal techniques during the second stage of labour for reducing perineal trauma. *Cochrane Database Syst Rev* 2017;6:CD006672
- McCandlish R, Bowler U, van Asten H, et al. A randomised controlled trial of care of the perineum during second stage of normal labour. *Br J Obstet Gynaecol* 1998;105(12):1262–1272
- Mayerhofer K, Bodner-Adler B, Bodner K, et al. Traditional care of the perineum during birth. A prospective, randomized, multicenter study of 1,076 women. *J Reprod Med* 2002;47(06):477–482
- Foroughipour A, Firuzeh F, Ghahiri A, Norbakhsh V, Heidari T. The effect of perineal control with hands-on and hand-poised methods on perineal trauma and delivery outcome. *J Res Med Sci* 2011;16(08):1040–1046
- Rezaei R, Saatsaz S, Chan YH, Nia HS. A comparison of the “hands-off” and “hands-on” methods to reduce perineal lacerations: a randomised clinical trial. *J Obstet Gynaecol India* 2014;64(06):425–429
- Rasmussen OB, Yding A, Anhøj J, Sander Andersen C, Boris J. Reducing the incidence of Obstetric Sphincter Injuries using a hands-on technique: an interventional quality improvement project. *BMJ Qual Improv Rep* 2016;5(01):217936
- Sultan AH. Obstetric perineal injury and anal incontinence. *Clin Risk* 1999;5:193–196
- Murray D. What should we do when traditional research fails? *Anaesthesia* 2017;72(09):1059–1063
- Albers LL, Sedler KD, Bedrick EJ, Teaf D, Peralta P. Factors related to genital tract trauma in normal spontaneous vaginal births. *Birth* 2006;33(02):94–100
- Poots AJ, Reed JE, Woodcock T, Bell D, Goldmann D. How to attribute causality in quality improvement: lessons from epidemiology. *BMJ Qual Saf* 2017;26(11):933–937
- Andrews V, Sultan AH, Thakar R, Jones PW. Risk factors for obstetric anal sphincter injury: a prospective study. *Birth* 2006;33(02):117–122
- Frieden TR. Evidence for health decision making - beyond randomized, controlled trials. *N Engl J Med* 2017;377(05):465–475
- Boutron I, Altman DG, Moher D, Schulz KF, Ravaud P; CONSORT NPT Group. CONSORT statement for randomized trials of nonpharmacologic treatments: a 2017 update and a CONSORT extension for nonpharmacologic trial abstracts. *Ann Intern Med* 2017;167(01):40–47