Biomechanics Potential of the Masseter and Temporal Muscles in the Mandibles of Mesofacial Subjects

Ana Paula Chitolina Pradebon¹ Marieli Chitolina Pradebon¹ Guilherme Reghelin Goulart²
Geraldo Pereira Jotz³ Tais Malysz^{3,4} Júlia Niero Páfaró⁵ Magda de Souza Reis¹ Deivis de Campos^{2,5,6}

¹ Department of Nursing and Odontology, Universidade de Santa Cruz do Sul, Santa Cruz do Sul, RS, Brazil

² Department of Biology and Pharmacy, Universidade de Santa Cruz do Sul, Santa Cruz do Sul, RS, Brazil

³ Department of Morphological Sciences, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

⁴ Postgraduation in Neuroscience Program, Instituto de Ciências Básicas da Saúde, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

⁵ Department of Basic Health Sciences, Universidade Federal de Ciências da Saúde de Porto Alegre, Porto Alegre, RS, Brazil

⁶ Department of Physical Education and Health, Postgraduate Program in Health Promotion, Universidade de Santa Cruz do Sul, Santa Cruz do Sul, RS, Brazil

Address for correspondence Deivis de Campos, PhD, Departamento de Ciências Básicas da Saúde, Universidade Federal de Ciências da Saúde de Porto Alegre – UFCSPA, Avenida Sarmiento Leite 245, 90050-170, Porto Alegre, RS, Brazil (e-mail: dcampos@ufcspa.edu.br).

J Morphol Sci 2018;35:180–182.

Abstract

Introduction The knowledge of the organization of the masseter muscle (MM) and the temporal muscle (TM) is extremely important when related to the study of the stomatognathic system. Moreover, some authors have shown that mastication is of great importance, not only for the intake of food but also for the systemic, mental and physical functions of the body.

Materials and Methods We have decided to analyze the biomechanical potential (length of the force arm, muscular work and mechanical advantage) of the MM and TM in the mandibles of mesofacial subjects ($n = 34$).

Results Our results show that the MM exhibits a better biomechanical potential than the TM ($p = 0.0001$).

Conclusion With these data, orthodontists may develop a specific treatment plan and get better results, especially in cases of patients in whom the biomechanical pattern of the temporomandibular joint is unfavorable.

Keywords

- masseter muscle
- temporal muscle
- mesofacial
- mandibles

Introduction

Some studies describe the type of fiber, the thickness and the neuromuscular pattern of the masticatory muscles.^{1–5} However, the descriptions regarding the biomechanical organization of the masticatory muscles in specific craniofacial standards are still inadequate, especially those that aim to elucidate the biomechanical differences between the human masseter muscle (MM) and the temporal muscle (TM).

Knowledge of the organization of the MM and of the TM is extremely important when related to the study of the stomatognathic system. This knowledge of the masticatory

muscle organization assists in researches associated with occlusion, facial growth and temporomandibular disorders.^{6–8} Thus, we have decided to analyze the biomechanical potential (length of the force arm, muscular work and mechanical advantage) of the MM and of the TM in the mandibles of mesofacial (ME) subjects.

For the present study, we have used 34 mandibles of adult subjects (mean age = 45.5 ± 7.1 years old) of both genders, from the collection of the Laboratory of Human Anatomy of the Universidade de Santa Cruz do Sul (UNISC, in the Portuguese acronym) (ethics committee—protocol 141346). All of the quantification procedures of the length of the force arm,

received
September 16, 2017
accepted
September 3, 2018
published online
November 14, 2018

DOI <https://doi.org/10.1055/s-0038-1675569>.
ISSN 2177-0298.

Copyright © 2018 by Thieme Revinter Publicações Ltda, Rio de Janeiro, Brazil

License terms

muscular work and mechanical advantage were made according to previous protocols.⁹

The distance between the condylar process and the coronoid process (the insertion site of the temporal muscle) represents the length of the force arm (LFA) of the TM, whereas the distance between the condylar process and the mental protuberance is the length of the resistance arm (LRA) (►Fig. 1). Similarly, the distance between the condylar process and the anterior border of the masseteric tuberosity (the insertion site of the masseter muscle) represents the LFA of the MM, whereas the distance between the condylar process and the mental protuberance is the LRA (►Fig. 1). Thus, the mechanical advantage of the TM and of the MM can be obtained using the following ratio: LFA/LRA. The inverse of this ratio represents the muscular work (LRA/LFA) of both muscles.

All of the measurements were performed on both sides (right and left) of all mandibles. However, we have decided to use data from one side (left) only because, using the paired t-test, we have concluded that there is no statistical difference between the sides and no apparent tendency toward any such difference ($p = 0.3574$). Only mandibles without any deformity or evidence of bone pathology were used. A p -value ≤ 0.05 was considered significant in all statistical tests performed. Data were analyzed with the aid of the GraphPad Prism 5.01 software (GraphPad Software, Inc., San Diego, CA, USA).

In the present study, the comparison (paired t-test) of the biomechanical potential between the MM and the TM in the mandibles of ME subjects showed that the results were significantly higher in MM for LFA and mechanical advantage

Table 1 Comparison of all morphometric parameters estimated

Morphometric Parameters	Masseter Muscle Mean $\pm$ SD	Temporal Muscle Mean $\pm$ SD	p -value
Length of the resistance arm – L _{RA} (mm)	106.14 $\pm$ 5.82	106.14 $\pm$ 5.82	-
Length of the force arm – L _{FA} (mm)	44.6 $\pm$ 4.7	36.9 $\pm$ 3.0	0.0001
Muscular work (L _{RA} /L _{FA})	2.39 $\pm$ 0.21	2.88 $\pm$ 0.23	0.0001
Mechanical advantage (L _{FA} /L _{RA})	0.42 $\pm$ 0.03	0.34 $\pm$ 0.02	0.0001

Abbreviations: SD, standard deviation.

(17.2% and 19.0%, respectively). Thus, the muscular work of the TM proved to be 17.0% higher than that of the MM (►Table 1).

Finally, the present work aims to provide a simple but multidisciplinary synthesis of the current knowledge concerning the morphogenesis of the biomechanical organization of the main muscles of mastication and to help to promote future studies in this area. With these data, orthodontists may develop a specific treatment plan and get better results, especially in cases of patients whose biomechanical pattern of the temporomandibular joint is unfavorable.

Fig. 1 Schematic drawing of the mandible showing the specific points that were used to measure the length of the force arm (LFA) and the length of the resistance arm (LRA) of the masseter muscle (MM) and temporal muscle (TM). a, condylar process; b, masseteric tuberosity; *, anterior border of the masseteric tuberosity; c, mental protuberance.

Therefore, the purpose of the present study is to compile the latest scientific information concerning the relationship between mastication and general health.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- 1 Boer NP, Batigália F, Alonso LG, Ribeiro EC. Classification methods for human masticatory muscle fibers: critical literature review. *Braz J Morphol Sci* 2007;24(03):164–169
- 2 Emshoff R, Emshoff I, Rudisch A, Bertram S. Reliability and temporal variation of masseter muscle thickness measurements utilizing ultrasonography. *J Oral Rehabil* 2003;30(12):1168–1172
- 3 Farella M, Bakke M, Michelotti A, Rapuano A, Martina R. Masseter thickness, endurance and exercise-induced pain in subjects with different vertical craniofacial morphology. *Eur J Oral Sci* 2003;111(03):183–188
- 4 Korfage JA, Brugman P, Van Eijden TM. Intermuscular and intramuscular differences in myosin heavy chain composition of the human masticatory muscles. *J Neurol Sci* 2000;178(02):95–106
- 5 Palinkas M, Nassar MS, Cecilio FA, et al. Age and gender influence on maximal bite force and masticatory muscles thickness. *Arch Oral Biol* 2010;55(10):797–802
- 6 Arijji Y, Sakuma S, Izumi M, et al. Ultrasonographic features of the masseter muscle in female patients with temporomandibular disorder associated with myofascial pain. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004;98(03):337–341
- 7 Bakke M, Thomsen CE, Vilmann A, Soneda K, Farella M, Møller E. Ultrasonographic assessment of the swelling of the human masseter muscle after static and dynamic activity. *Arch Oral Biol* 1996;41(02):133–140
- 8 Raadsheer MC, Kiliaridis S, Van Eijden TM, Van Ginkel FC, Prahl-Andersen B. Masseter muscle thickness in growing individuals and its relation to facial morphology. *Arch Oral Biol* 1996;41(04):323–332
- 9 Marques HB, Richter FF, Heck L, Xavier LL, de Campos D. Biomechanical potential of the temporal muscle in brachyfacial and dolichofacial subjects: a study on dry mandibles. *Orthod Craniofac Res* 2016;19(03):162–168