Serial Reduction of an Extremely Large Gastroschisis using Vacuum-Assisted Closure

Marilyn W. Butler¹ Julie Fuchs² Matias Bruzoni²

¹ Division of Pediatric Surgery, Department of Surgery, Oregon Health and Science University, Portland, Oregon, United States

² Division of Pediatric Surgery, Department of Surgery, Stanford University, Stanford, California, United States

Address for correspondence Marilyn W. Butler, MD, MPH, Division of Pediatric Surgery, Department of Surgery, Oregon Health and Science University, 501 N Graham Street, Suite 300, Portland, OR 97227, United States (e-mail: mwbutler@gmail.com).

Eur J Pediatr Surg Rep 2018;6:e97–e99.

Abstract

Keywords

- gastroschisis
- omphalocele
- vacuum-assisted closure
- VAC
- negative-pressure wound therapy

We herein describe a case of serial reduction of an extremely large and complex gastroschisis using vacuum-assisted closure (VAC) therapy in a boy born at 35^{5/7} weeks' gestation. A spring-loaded silicone silo was placed at birth. By day of life (DOL) 22, minimal visceral contents had been reduced, and the silo was difficult to maintain due to the size of the fascial defect and loss of abdominal domain. A bespoke VAC dressing was constructed, and biweekly dressing changes allowed gradual reduction of the gastroschisis until the viscera were consolidated. By DOL 50, the viscera were completely reduced, and VAC therapy was discontinued. Feeds were commenced on DOL 57 and increased to goal by DOL 86. The baby was discharged home on DOL 115. We conclude that VAC dressings can be used to aid gradual reduction of an extremely large gastroschisis, particularly in medical fragile infants.

New Insights and the Importance for the Pediatric Surgeon

When silo closure fails, an extremely large gastroschisis can be gradually reduced with VAC dressings, thereby avoiding the use of synthetic or biological mesh.

Introduction

While mortality of gastroschisis in some low- and middle-income countries remains as high as 100%, in developed countries, gastroschisis mortality is an extremely rare occurrence given access to skilled neonatal and surgical care, including ventilator support and total parenteral nutrition. However, complex gastroschisis, whether it is defined by gastrointestinal issues or abdominal wall factors, behaves as a disease process with much worse prognosis.¹

Despite improvements in survival, gastroschisis closure remains challenging, particularly when the fascial defect is large, eviscerated contents are substantial, medical status is precarious, or abdominal domain is lost.

Negative-pressure wound therapy (NPWT), including vacuum-assisted closure (VAC), has been used in children

since the 1990s for other challenging wounds to promote the formation of granulation tissue, reduce edema, remove infection, and control fluid losses.^{2,3} Previous series have described the application of NPWT for gastroschisis following silo placement after the eviscerated bowel is consolidated.^{2,4,5} We herein report the first case where VAC therapy was used to reduce a significantly matted mass of eviscerated bowel in an infant with gastroschisis.

Case Report

A boy with gastroschisis was born at 35^{5/7} weeks' gestation weighing 2.120 kg, to a G2P1 mother through vaginal delivery. Through a defect measuring approximately 4 cm in diameter, a substantial portion of matted abdominal contents was eviscerated, including the stomach, small intestines, and colon. A

received
July 4, 2018
accepted after revision
October 11, 2018

DOI <https://doi.org/10.1055/s-0038-1676045>.
ISSN 2194-7619.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms

Fig. 1 Initial vacuum-assisted closure dressing on day of life 22.

spring-loaded 5-cm Silicone Silo Bag was placed at birth (Bentec Medical, Woodland, California, United States) and was eventually upsized to a 7.5-cm Silicone Silo Bag. By day of life (DOL) 22, minimal visceral contents had been reduced, and the silo was difficult to maintain due to the large size of the fascial defect and loss of abdominal domain. A bespoke VAC dressing was constructed as follows: Whitefoam and Granu-Foam (KCI Medical, San Antonio, Texas, United States) dressings were cut to half their thickness, fashioned in the shape of a cup by sewing strips together, and placed over the eviscerated bowel. Strips of adhesive drapes were used to secure the dressing, circumferentially wrapping the infant. After puncturing the drape, a SensaT.R.A.C. Pad (KCI Medical) was placed over the dressing, connected to a VAC therapy unit, and placed to negative pressure that ranged between 25 and 75 mm Hg (→Fig. 1).

Endotracheal intubation was necessary for biweekly dressing changes until the viscera were consolidated (→Fig. 2). Then Mepitel (Mölnlycke Health Care, Gothenburg, Sweden) was placed over the viscera prior to Whitefoam and Granu-Foam dressings (→Fig. 3). By DOL 50, the abdominal contents

Fig. 2 Gastroschisis on day of life 29, 7 days after vacuum-assisted closure dressing was placed, showing consolidation of herniated abdominal contents.

Fig. 3 Gastroschisis on day of life 39, 17 days after vacuum-assisted closure dressing was placed, showing dressing materials and nearly complete reduction of herniated abdominal contents.

were completely reduced, and VAC therapy was discontinued. Mepitel was placed until the wound was closed (→Fig. 4).

Feeds were commenced on DOL 57 and increased to goal by DOL 86. The baby was discharged home on DOL 115. While initially anemic with cholestasis, by follow-up at 13 months, both had resolved, and his weight was at the 50th percentile for his age. At that point, he was discharged from the clinic and lost to further follow-up. At follow-up clinic visits, the skin had completely epithelialized. An incisional hernia was not noted, although it is presumed that one was present.

Discussion

Gastroschisis closure has challenged pediatric surgeons for decades. Standard of care for many years entailed immediate primary closure at birth, which resulted in respiratory compromise, abdominal compartment syndrome, bowel ischemia, decreased lower extremity perfusion, anuria, and need for vigorous fluid resuscitation to maintain blood pressure.⁶ Occasionally, in challenging cases, silicone sheeting or other plastic

Fig. 4 Gastroschisis on day of life 78, showing wound contraction.

materials such as intravenous bags were sewn to the edges of the fascia or the skin to create temporary silos until the viscera could be gradually reduced.^{7–10} Some used pneumatic pressure or wringer clamps to reduce the viscera within the silos.^{11,12} With time, however, the fascial defect typically grew larger, and sutures caused the fascia to lose its integrity, making subsequent closure more difficult. In 1995, the introduction of spring-loaded Silicone Silo Bags that could be easily placed at the bedside made delayed closure more convenient, and primary reduction was reserved for newborns whose abdominal contents could be reduced without significant physiological consequences.^{13–15} In some resource-poor countries, locally sourced female condoms have been used when spring-loaded silicone silos are not available.¹⁶

Despite these advances, in patients with large fascial defects and loss of abdominal domain, use of spring-loaded silos has been unsuccessful in reducing the gastroschisis. At various times, surgeons have used muscle or skin flaps,^{17,18} skin grafts,¹⁹ synthetic mesh,²⁰ biological mesh,²¹ amniotic grafts,²² and pericardial patches²³ to achieve visceral coverage and reduce fluid and heat losses. Skin does not readily cover many of these materials, however, and often mesh needs to be removed^{7,17,20,24} or leads to recurrence.²¹

More recently, sutureless closure of gastroschisis has incorporated the umbilical cord as a biological dressing for primary repair or delayed closure after silo placement.^{25,26} After the cord sloughs, VAC dressings are often used to cover the exposed viscera until the wound is epithelialized.^{2,4,5} Until the successful closure in our patient, however, there have been no reports of the use of VAC dressings for primary coverage in gastroschisis that has not yet been reduced. Although our patient did need intubation for the first few dressing changes, once the bowel was consolidated, the VAC dressing could be applied without sedation. Feeds were commenced once the abdomen was closed, and cholestatic liver changes resolved after 13 months.

We do note that our patient's initial management with the silo bag might have made eventual reduction more difficult. Although we generally use the smallest silo bag possible in order not to enlarge the fascial defect, our patient's defect was initially relatively large, and therefore we had no choice than to use a 5-cm silo bag. Upsizing the silo bag made the fascial defect even larger, and in retrospect, were we to encounter the same problem again, we would opt to commence VAC dressings sooner.

We conclude that VAC dressings can be used to aid gradual reduction of extremely large gastroschisis anomalies, particularly in medically fragile infants, without the need for synthetic or biological mesh.

Conflict of Interest

None.

References

- Laje P, Fraga MV, Peranteau WH, et al. Complex gastroschisis: clinical spectrum and neonatal outcomes at a referral center. *J Pediatr Surg* 2018;53(10):1904–1907
- Stoffan AP, Ricca R, Lien C, Quigley S, Linden BC. Use of negative pressure wound therapy for abdominal wounds in neonates and infants. *J Pediatr Surg* 2012;47(08):1555–1559
- Lopez G, Clifton-Koeppel R, Emil S. Vacuum-assisted closure for complicated neonatal abdominal wounds. *J Pediatr Surg* 2008;43(12):2202–2207
- McBride CA, Stockton K, Storey K, Kimble RM. Negative pressure wound therapy facilitates closure of large congenital abdominal wall defects. *Pediatr Surg Int* 2014;30(11):1163–1168
- Choi WW, McBride CA, Kimble RM. Negative pressure wound therapy in the management of neonates with complex gastroschisis. *Pediatr Surg Int* 2011;27(08):907–911
- Yaster M, Buck JR, Dudgeon DL, et al. Hemodynamic effects of primary closure of omphalocele/gastroschisis in human newborns. *Anesthesiology* 1988;69(01):84–88
- Cordero L, Touloukian RJ, Pickett LK. Staged repair of gastroschisis with Silastic Sheeting. *Surgery* 1969;65(04):676–682
- Schuster SR. A new method for the staged repair of large omphaloceles. *Surg Gynecol Obstet* 1967;125(04):837–850
- Gilbert MG, Mencia LF, Puranik SR, Litt RE, Altman DH. Management of gastroschisis and short bowel: report of 17 cases. *J Pediatr Surg* 1972;7(05):598–607
- Kidd JN Jr, Jackson RJ, Smith SD, Wagner CW. Evolution of staged versus primary closure of gastroschisis. *Ann Surg* 2003;237(06):759–764, discussion 764–765
- Sawin R, Glick P, Schaller R, Hatch E, Hall D, Hicks L. Gastroschisis wringer clamp: a safe, simplified method for delayed primary closure. *J Pediatr Surg* 1992;27(10):1346–1348
- Othersen HB Jr, Smith CD. Pneumatic reduction bag for treatment of gastroschisis and omphalocele. A 10-year experience. *Ann Surg* 1986;203(05):512–516
- Bianchi A, Dickson AP. Elective delayed reduction and no anesthesia: 'minimal intervention management' for gastrochisis. *J Pediatr Surg* 1998;33(09):1338–1340
- Lansdale N, Hill R, Gull-Zamir S, et al. Staged reduction of gastroschisis using preformed silos: practicalities and problems. *J Pediatr Surg* 2009;44(11):2126–2129
- Fischer JD, Chun K, Moores DC, Andrews HG. Gastroschisis: a simple technique for staged silo closure. *J Pediatr Surg* 1995;30(08):1169–1171
- Bustorff-Silva JM, Schmidt AF, Gonçalves A, Marba S, Sbragia L. The female condom as a temporary silo: a simple and inexpensive tool in the initial management of the newborn with gastroschisis. *J Matern Fetal Neonatal Med* 2008;21(09):648–651
- Fonkalsrud EW, Smith MD, Shaw KS, Borick JM, Shaw A. Selective management of gastroschisis according to the degree of viscer- oabdominal disproportion. *Ann Surg* 1993;218(06):742–747
- Aoyama K. A new operation for repair of large ventral hernias following giant omphalocele and gastroschisis. *J Pediatr Surg* 1979;14(02):172–176
- Anderl H, Menardi G, Hager J. Closure of gastroschisis by mesh skin grafts in problem cases. *J Pediatr Surg* 1986;21(10):870–872
- Stringel G. Large gastroschisis: primary repair with Gore-Tex patch. *J Pediatr Surg* 1993;28(05):653–655
- Beres A, Christison-Lagay ER, Romao RL, Langer JC. Evaluation of Surgisis for patch repair of abdominal wall defects in children. *J Pediatr Surg* 2012;47(05):917–919
- Gharib M, Ure BM, Klose M. Use of amniotic grafts in the repair of gastroschisis. *Pediatr Surg Int* 1996;11(2-3):96–99
- van Tuil C, Saxena AK, Willital GH. Experience with management of anterior abdominal wall defects using bovine pericard. *Hernia* 2006;10(01):41–47
- T Adikibi B, O'Toole S. Reverse tissue expansion in gastroschisis: what to do if the defect is too large to close after silo removal? *J Neonatal Surg* 2014;3(04):47
- Zivković SM. Repair of gastroschisis using umbilical cord as a patch. *J Pediatr Surg* 1991;26(10):1179–1180
- Sandler A, Lawrence J, Meehan J, Phearman L, Soper R. A "plastic" sutureless abdominal wall closure in gastroschisis. *J Pediatr Surg* 2004;39(05):738–741