PCSK9 in Haemostasis and Thrombosis: Possible Pleiotropic Effects of PCSK9 Inhibitors in Cardiovascular Prevention

Francesco Paciullo¹ Stefania Momi¹ Paolo Gresele¹

¹ Division of Internal and Cardiovascular Medicine, Department of Medicine, University of Perugia, Perugia, Italy

Thromb Haemost 2019;119:359–367.

Address for correspondence Paolo Gresele, MD, PhD, Division of Internal and Cardiovascular Medicine, Department of Medicine, University of Perugia, Strada Vicinale Via delle Corse, 06126 Perugia, Italy (e-mail: paolo.gresele@unipg.it).

Abstract

Since increased cholesterol levels are crucial in determining the development of atheroma, their reduction represents a mainstay in primary and secondary cardiovascular prevention. The most recent spectacular advancement in cholesterol-lowering therapy is represented by proprotein convertase subtilisin/kexin type-9 (PCSK9) inhibitors. Although their benefit over currently available treatments has been ascribed primarily to their strong low-density lipoprotein (LDL)-cholesterol reducing action, several clues suggest that PCSK9 inhibitors may also influence platelet function and blood coagulation. PCSK9 knockout mice develop less venous and arterial thrombosis and show reduced in vivo platelet activation upon arterial injury. In patients with acute coronary syndromes (ACSs) treated with P2Y₁₂ inhibitors, a direct association between PCSK9 serum levels and residual platelet reactivity was found. A direct correlation between urinary excretion of 11-dehydro-thromboxane-B₂, a marker of in vivo platelet activation, and circulating PCSK9 levels was reported in patients with atrial fibrillation. Moreover, recombinant human PCSK9 added in vitro to human platelets potentiated activation induced by weak agonists. Finally, blood clotting factor VIII (FVIII), which is associated with stroke and ACS risk, is cleared from the circulation by members of the LDL receptor (LDLR) family. Given that PCSK9 degrades LDLR, it is conceivable that PCSK9 inhibitors by enhancing the expression of LDLR may slightly decrease circulating FVIII, in this way contributing to the prevention of cardiovascular events. This review aims to discuss the possible and hypothetical interactions between PCSK9 and the haemostatic system and to examine the possible pleiotropic effects of PCSK9 inhibitors in cardiovascular prevention.

Keywords

- ▶ atherothrombosis
- ▶ CD36
- ▶ dyslipidaemia
- ▶ FVIII
- ▶ LOX-1

PCSK9, Lipid Metabolism and Cardiovascular Risk

Cholesterol is transported in the bloodstream by lipoprotein particles. The two major cholesterol-carrying lipoproteins are low-density lipoproteins (LDLs) and high-density lipoproteins (HDLs). LDLs, whose main protein fraction is apolipoprotein B100, transport cholesterol from the liver to

peripheral tissues, including the arterial walls.¹ LDLs are a major determinant of atherosclerosis, and both American and European guidelines recommend specific LDL threshold reductions to prevent cardiovascular events.² On the other hand, HDLs, whose main protein fraction is apolipoprotein A-I, transport cholesterol from peripheral tissues to the liver, facilitating its clearance.³ Circulating cholesterol levels are regulated by the balance between their biosynthesis and

received
March 19, 2018
accepted after revision
November 19, 2018

© 2019 Georg Thieme Verlag KG
Stuttgart · New York

DOI <https://doi.org/10.1055/s-0038-1676863>.
ISSN 0340-6245.

clearance. The rate-limiting step in cholesterol biosynthesis is the conversion of 3-hydroxy-3-methyl glutaryl coenzyme A (HMG-CoA) into mevalonate catalyzed by HMG-CoA reductase, which represents the pharmacologic target of statins, the most widely used cholesterol-lowering agents.⁴ LDL clearance takes place via specific LDL receptors (LDLRs) in the liver which bind apolipoprotein B100 forming a complex which is internalized.⁵ The LDL component of the complex is then degraded by lysosomal enzymes, while LDLR are recycled to the cell membrane to bind other LDL particles.⁶ The expression of LDLR on the hepatocyte surface is regulated by intracellular cholesterol levels, with lower intracellular cholesterol leading to increased surface LDLR, and vice versa.⁷ LDLRs are also regulated by proprotein convertase subtilisin/kexin type-9 (PCSK9), originally called Nrc-1, an enzyme which fosters their degradation. PCSK9 is mainly synthesized by the liver, but also by the brain, kidney, intestine, pancreas and steroidogenic tissue, and is found both in the intra- and extracellular space. Extracellularly, PCSK9 binds the first epidermal growth factor-like repeat of LDLR forming a tri-molecular complex (LDLR-LDL-PCSK9) which is then internalized. Once inside the hepatocyte, PCSK9 prevents LDLR from escaping lysosomal degradation, and therefore to recycle to the cell surface, thus reducing their expression. Intracellularly, PCSK9 binds nascent LDLR and targets them to lysosomes where they are degraded.⁸ The interest in PCSK9 has sharply risen after the observation that gain of function variants of its gene are associated with enhanced levels of circulating LDL cholesterol (LDL-c), and thus with increased cardiovascular risk.⁹ Conversely, carriers of loss-of-function variants have low LDL-c and a reduced risk of ischaemic cardiovascular disease.¹⁰ Moreover, in a seminal study, the adenoviral-mediated over-expression of PCSK9 in mice caused a striking increase in plasma cholesterol.¹¹ Since the discovery of the important role of PCSK9 in lipid metabolism, numerous approaches to its inhibition have been attempted, leading to the development of monoclonal antibodies (MoAbs) blocking its extracellular action, probably the most relevant for the regulation of circulating cholesterol levels.¹² Several phase III trials with anti-PCSK9 MoAb have been performed in patients with hypercholesterolemia and high cardiovascular risk and, based on their positive results in terms of safety and efficacy, two fully humanized anti-PCSK9 MoAbs, alirocumab and evolocumab, have received Food and Drug Administration (FDA) and European Medicines Agency approval in 2015 as a second line treatment, in addition to diet and maximally tolerated statin therapy, to lower LDL-c in adults with heterozygous familial hypercholesterolaemia or clinical atherosclerotic cardiovascular disease who require additional LDL-c lowering. Recently, based on the results of the landmark FOURIER trial,¹³ the FDA has approved evolocumab also to prevent heart attacks, strokes and coronary revascularization in adults with established cardiovascular disease. In this trial, including more than 27,000 high-risk patients with atherosclerotic cardiovascular disease with a LDL-c of ≥ 70 mg/dL who were receiving statin therapy, subjects randomized to evolocumab showed a significantly lower incidence of major

adverse cardiovascular events compared with those randomized to placebo.¹³ Moreover, a meta-analysis of 24 phase II and phase III trials involving 10,159 adults with hypercholesterolaemia, on statin treatment or not, has shown that anti-PCSK9 MoAbs significantly reduce all-cause mortality compared with no anti-PCSK9 treatment (odds ratio [OR], 0.45 [95% confidence interval CI, 0.23–0.86], $p = 0.015$).¹⁴ Thus, PCSK9 inhibitors appear to be a very effective strategy to prevent cardiovascular events.

Pleiotropic Effects of Lipid-Lowering Interventions

Since the detrimental effect of LDL-c on the cardiovascular system is concentration-dependent, the reduction of ischaemic cardiovascular events produced by PCSK9 inhibitors when added to statins has been ascribed primarily to their further, strong LDL-c-lowering action.¹⁵

However, statins may exert cardiovascular protective actions independent from LDL-c lowering which have been called pleiotropic effects.¹⁶ The reduction of cardiovascular events attained with statins in some clinical trials, like the JUPITER trial,¹⁷ has been indeed greater than that expected solely from LDL-c reduction.^{18,19} In fact, cholesterol-independent beneficial effects of statins on the cardiovascular system, such as the stabilization of atherosclerotic plaques, reversal of endothelial dysfunction, blunting of inflammation, enhancement of fibrinolysis and inhibition of platelet activation and blood coagulation, have been well documented.^{20–23}

On the other hand, in a recent meta-analysis of trials with lipid-lowering interventions including more than 300,000 patients, the relative risk reduction of major vascular events associated with PCSK9 inhibitors use was higher, even if not significantly, than that observed with statins for the same LDL-c reduction (OR, 0.49 [95% CI, 0.34–0.71] vs. 0.61 [95% CI, 0.58–0.65]).²⁴ Reasons advocated for this observation are the raising activity on anti-atherogenic HDL and the ability to reduce lipoprotein (a) [Lp(a)] of PCSK9 inhibitors.^{25,26} However, pleiotropic effects of PCSK9 inhibitors independent from lipid metabolism can also be considered.²⁷

Indeed, many pre-clinical and clinical data support the hypothesis that the cardiovascular protective effect of PCSK9 inhibitors may be more complex, involving mechanisms which go beyond their lipid-lowering action, several of which may affect the haemostatic system.

Hints for Pleiotropic Effects of PCSK9 on Haemostasis and Thrombosis

Haemostasis begins at a vascular injury site with platelet adhesion and aggregation (primary haemostasis), followed by the activation cascade of clotting factors (secondary haemostasis). Thrombosis, which can be considered an excessive extension of a haemostatic reaction, occurs in the arterial and venous vascular beds by mechanisms which differ in relation to the anatomical and rheological characteristics of these two systems. In arteries, thrombosis originates from the rupture of atherosclerotic lesions and is mainly generated by platelet

activation,²⁸ while in veins, thrombi are mainly the consequence of clotting activation favoured by stasis, hypercoagulability and endothelial damage.²⁹ However, several recent observations suggest that these two pathways contribute to both arterial and venous thrombosis. In fact, risk factors for ischaemic cardiovascular disease, among which hyperlipidaemia, induce endothelial dysfunction in both the arterial and venous vascular beds, leading to the development of either arterial or venous thrombosis depending on the concomitant inciting conditions.^{30,31}

Studies in Animals

The discovery of the central role played by PCSK9 in lipid metabolism has also been grounded on studies performed in gene-modified mice. The adenovirus-induced over-expression of PCSK9 in mice resulted in decreased hepatic LDLR expression with associated hypercholesterolaemia, whereas the deletion of the PCSK9 gene increased hepatic LDLR expression and reduced LDL-c circulation,¹¹ recapitulating the phenotypes of gain- or loss-of-function PCSK9 variants in humans.

Interestingly, PCSK9^{-/-} mice showed a reduction of FeCl₃ injury-induced carotid artery thrombosis, with the formation of unstable non-occlusive thrombi,³² suggesting an impaired platelet function. Indeed, in these mice the activation of circulating platelets provoked by arterial injury, shown by increased glycoprotein (GP) IIb/IIIa activation, P-selectin expression and circulating platelet-leukocyte aggregates, was strikingly reduced as compared with control mice.³² It cannot be disregarded, however, that the assessment of *in vivo* platelet activation in mice by the measurement of these biomarkers may be prone to errors.³³

Moreover, the hypercoagulable state induced by sepsis was exacerbated in PCSK9 over-expressing mice, as shown by enhanced thrombin-antithrombin complexes and reduced protein C plasma levels,³⁴ suggesting that changes in PCSK9 may also have an impact on blood coagulation.

Indeed, PCSK9^{-/-} mice developed significantly smaller venous thrombi as compared with wild-type mice after inferior vena cava ligation.³⁵ Plasma factor VIII (FVIII) levels are known to modulate venous thrombosis in mice,^{36,37} and there are theoretical reasons to suppose that PCSK9 may modulate circulating FVIII levels (see later); thus, although no blood clotting factor measurements were made in this study,³⁵ it is conceivable that an effect on FVIII may have contributed to reduce venous thrombosis. On the other hand, plasma levels of soluble P-selectin (sP-selectin), a platelet and endothelial activation biomarker,³⁸ were significantly lower in PCSK9-deficient than in control mice after the induction of vena cava thrombosis,³⁵ further suggesting that PCSK9^{-/-} mice have impaired platelet function, but also that PCSK9 deletion may reduce endothelial activation. In this regard, it is interesting that increased circulating sP-selectin was observed in humans with unprovoked deep vein thrombosis who have endothelial dysfunction.^{30,31}

Studies in Humans

The possible role of PCSK9 in modulating platelet function has been assessed also in humans. In a prospective, observational

study in patients with a recent acute coronary syndrome (ACS) undergoing percutaneous coronary intervention and receiving P2Y₁₂ inhibitors, the PCSK9-REACT study, a direct correlation between PCSK9 plasma levels and high-on-treatment platelet reactivity was observed, suggesting that PCSK9 enhances platelet activation.³⁹ Indeed, recent observations have shown that human recombinant PCSK9 pre-incubated *in vitro* with platelets potentiates aggregation, P-selectin expression and GPIIb/IIIa activation induced by a weak agonist,³² acting therefore as a primer of platelet activation.⁴⁰

In this regard, it is interesting that twice as much PCSK9 was found to be contained in platelets from type 2 diabetes mellitus (T2DM) patients with coronary artery disease (CAD) than in platelets from healthy controls,⁴¹ suggesting that PCSK9 may be released during platelet activation and contribute to the well-established T2DM-associated platelet hyper-reactivity and impaired responsiveness to anti-platelet agents.^{42,43}

Human megakaryocytes express messenger ribonucleic acid (mRNA) for PCSK9, although at low levels (A.S. Weyrich and R.A. Campbell, University of Utah, personal communication), but this is not found in platelets.⁴⁴ It can therefore be hypothesized that mRNA for PCSK9 is handled by megakaryocytes similarly to mRNA for matrix metalloproteinase-2, with mRNA not sorted into platelets but the protein present,⁴⁵ even if only in a platelet sub-population,⁴¹ as already reported for tissue factor (TF)⁴⁶ and endothelial nitric oxide synthase.⁴⁷

It was found that PCSK9 plasma levels positively correlate with the platelet count and plateletcrit in patients with stable CAD, further suggesting a link between PCSK9 and platelets in patients with coronary disease.⁴⁸ Furthermore, PCSK9 plasma levels have been recently reported to increase during an acute coronary event,⁴⁹ a condition associated with a striking bout of *in vivo* platelet activation,^{50,51} and to positively correlate with the severity of coronary artery lesions evaluated by the SYNTAX score.⁴⁹ On the other hand, platelet reactivity in ACS correlates positively with the SYNTAX score,⁵² further suggesting a role of enhanced circulating PCSK9 in platelet hyper-reactivity. Moreover, activated platelets of patients with CAD release soluble sortilin⁵³ which is known to facilitate PCSK9 secretion,⁵⁴ making even more plausible the hypothesis of a positive feedback activation of platelets through PCSK9 during ACS.

A direct relationship between *in vivo* platelet activation and PCSK9 plasma levels has been reported also in atrial fibrillation. In this study, circulating PCSK9 correlated with urinary 11-dehydro-thromboxane B₂ excretion,⁵⁵ an unbiased marker of *in vivo* platelet activation.⁵⁶

Several clues suggest that PCSK9 may also influence secondary haemostasis in humans. In fact, plasma levels of TF, a pro-coagulant glycoprotein triggering thrombin formation and playing a central role in atherothrombosis,⁵⁷ positively correlated with plasma PCSK9 in patients with CAD and T2DM.⁵⁸ Moreover, single nucleotide polymorphisms of the PCSK9 gene were shown to be associated with the development of thrombosis in carriers of anti-phospholipid antibodies,⁵⁹ subjects characterized by a hypercoagulable state and *in vivo* platelet activation,^{60,61} confirming a link between PCSK9, platelets and blood coagulation.

Altogether, these data represent a strong hint that the association of PCSK9 with cardiovascular risk involves mechanisms that go beyond the mere regulation of circulating LDL-c and which include effects on the haemostatic system.

Modulation of Platelet Activation by Lipoproteins and the Possible Influence of PCSK9 Inhibitors

Dyslipidaemia may influence platelet reactivity and haemostasis by several mechanisms. Enhanced oxidative stress associated with high circulating LDL-c leads to the formation of oxidized-LDL (ox-LDL), which strongly contribute to inflammation-driven thrombosis⁶² by activating CD36 and LOX-1 receptors on platelets^{63,64} (→ Fig. 1). CD36 is a member of the SRB-1 family which binds native- and ox-LDL and plays a role in thrombus formation,⁶⁵ while LOX-1 (also called oLR-1) is a multi-ligand scavenger receptor whose expression is triggered by pro-inflammatory stimuli.^{66,67} In addition, activated platelets themselves oxidize LDL generating ox-LDL, in this way propagating platelet activation.⁶⁸ Another mechanism by which dyslipidaemia may affect platelet reactivity is through the generation of lipid peroxide-modified phospholipids, which activate platelets by acting on Toll-like receptor 2.⁶⁹

Oxidized phospholipids are transported by Lp(a),⁷⁰ and the latter may also directly activate platelets by not yet unravelled mechanisms.⁷¹

Finally, HDL attenuate platelet function by interacting with the ApoER2 and SRB1 receptors, and also promoting cholesterol efflux from platelet membranes.⁷² Indeed, cholesterol incorporation in plasma membranes induces platelet hypersensitivity to stimuli, whereas its depletion strikingly reduces platelet reactivity.⁷³ Moreover, HDL inhibit platelet fibrinogen binding and aggregation in response to thrombin via decreased formation of the second messengers diacylglycerol and inositol trisphosphate,⁷⁴ and enhance platelet NO generation, thus increasing cyclic guanosine monophosphate, by acting on the apoER2 receptor.⁷⁵ Finally, HDL also down-regulate the coagulation cascade and stimulate fibrinolysis.⁷⁶

Considering the above summarized mechanisms, PCSK9 inhibition may modulate the effects of lipoproteins on platelets at various levels, thus reducing platelet activation. By strikingly decreasing plasma LDL-c, PCSK9 inhibitors may deplete platelet membranes of cholesterol, thus reducing platelet reactivity and pro-coagulant activity. In this regard, it is interesting that treatment of hypercholesterolaemic subjects with rosuvastatin, a powerful cholesterol-lowering statin, reduced platelet membrane cholesterol, TF expression and generation of FXa.⁷⁷

Fig. 1 Hypothetical effects of proprotein convertase subtilisin/kexin type-9 (PCSK9) and its inhibition on platelets. (1) PCSK9 inhibition strikingly reduces low-density lipoprotein cholesterol (LDL-c) levels, thus potentially depleting platelet membranes of cholesterol, a mechanism reducing platelet reactivity.⁷³ Moreover, the inhibition of platelet activation by PCSK9 inhibitors (2), a result of the various effects shown in the figure, may also reduce the ability of platelets to oxidize LDL thus decreasing the platelet stimulating activity of the latter through CD36 and LOX-1 receptors (3). Furthermore, PCSK9 inhibition reduces lipoprotein (a) [Lp(a)] levels, the main carriers of ox-phospholipids, thus potentially blunting their ability to activate platelets either through the Toll-like receptor 2 (TRL2) receptor (4) or directly (5). Finally, PCSK9 inhibition increases high-density lipoprotein (HDL) which reduces platelet activation acting on apoER2 and SRB1 receptors (6), and scavenging cholesterol from platelet membranes (7). +, stimulation; –, inhibition.

In addition, given that PCSK9 and LOX-1 positively influence the expression of each other,⁷⁸ it can be envisaged that PCSK9 inhibitors may reduce platelet LOX-1.

Moreover, the blunting of platelet activation by PCSK9 inhibitors may decrease ox-LDL generation,⁶⁸ interrupting the vicious circle that propagates platelet activation. Indeed, treatment of hypercholesterolaemic patients with alirocumab or evolocumab was shown to reduce platelet activation.⁷⁹ PCSK9 inhibitors, differently from statins, lower Lp(a) levels,^{26,80} and thus they may reduce its direct and oxidized phospholipid-mediated stimulatory effect on platelets. In addition, PCSK9 inhibitors also enhance HDL levels,²⁵ in this way potentially inhibiting platelet aggregation directly or by depleting platelet membrane cholesterol⁷²⁻⁷⁵ (► Fig. 1).

Finally, cell apoptosis is known to favour thrombosis, in part through the release of pro-coagulant microparticles,⁸¹ and PCSK9 was found to enhance apoptosis in vascular smooth muscle and endothelial cells.^{82,83} In this context, PCSK9 inhibitors, by mitigating apoptosis, might indirectly prevent thrombosis.

Regulation of FVIII Levels by LDLR and the Hypothetical Effect of PCSK9 Inhibitors

Blood clotting FVIII is a key plasma protein, encoded by a gene located on chromosome X, which plays a central role in coagulation. FVIII acts as a co-factor for FIX, thus favouring thrombin generation. FVIII is transported in the circulation by von Willebrand factor (VWF) which stabilizes it and reduces its clearance.⁸⁴

Epidemiologic studies have shown an association between increased FVIII plasma levels and arterial thrombosis, and elevated FVIII levels were found to correlate with a higher recurrence rate in patients with a prior myocardial infarction or ischaemic stroke.⁸⁵⁻⁹⁰ In contrast, haemophilia patients appear to be protected from ischaemic heart disease.⁹¹

Experimental studies in animal models, although often involving the use of supra-physiologic concentrations, provide further evidence in support of the role of FVIII in arterial thrombosis.^{36,92,93} The increased risk of arterial thrombosis associated with enhanced FVIII levels is thought to be due to the combination of increased thrombin generation and enhanced platelet adhesion/aggregation, the latter being induced by the concomitant increase of VWF.^{94,95}

Circulating levels of FVIII are regulated by its biosynthesis and by its clearance through hepatic LDLR and lipoprotein receptor-related protein 1 (LRP1), both members of the LDLR family (► Fig. 2). FVIII is composed of a heavy and a light chain, the latter containing the binding site for LRP1, which is normally covered by VWF.⁹⁶ Therefore, circulating FVIII not bound to VWF (~5% of total) is quickly recognized by hepatocyte LRP1, endocytosed and degraded.⁹⁷ The important role of LRP1 in FVIII clearance in vivo has been confirmed by the strong elevation of circulating FVIII levels in LRP-deficient mice as well as in mice with adenovirus-mediated over-expression of receptor-associated protein, a chaperone for LRP1 which blocks the binding of all ligands to the receptor.⁹⁸ On the other hand, LDLR plays a role in FVIII clearance too because the simultaneous deletion of the LDLR and LRP1 genes in mice (double LRP1/LDLR^{-/-}) further enhanced FVIII levels by 4.2-fold, while the

Fig. 2 Hypothetical effects of proprotein convertase subtilisin/kexin type-9 (PCSK9) inhibitors on factor VIII (FVIII), von Willebrand factor (VWF) and tissue factor levels. Hepatic low-density lipoprotein receptor (LDLR) and lipoprotein receptor-related protein 1 (LRP1) are both involved in the clearance of FVIII and VWF by the liver. PCSK9 inhibitors enhance the expression of hepatic LDLR, and possibly also of LRP1 (1), by both intra- and extracellular mechanisms (2), thus potentially enhancing FVIII internalization and degradation leading to a decrease in FVIII plasma levels. Moreover, the possible enhancement of LRP1 by PCSK9 inhibitors in monocytes (3), might reduce circulating tissue factor (TF) through its accelerated clearance. +, stimulation; -, inhibition.

Table 1 Hypothetical pleiotropic effects of PCSK9 inhibitors in atherothrombosis

Cell/tissue targets	Reported PCSK9 effect	Possible impact of PCSK9 inhibition	Hypothetical mechanisms
Platelets	Platelet activation and secretion ³² Impaired response to antiplatelet agents ³⁹	Platelet inhibition Enhanced response to antiplatelet agents ³⁹	Inhibition of ox-LDL, Lp(a)-mediated activation Inhibition of CD36 and LOX1 receptor-mediated activation Enhancement of HDL-mediated inhibition
Blood coagulation	No data	Reduction of circulating FVIII	Enhanced LDLR and LRP1-mediated FVIII clearance
Tissue factor	Positive correlation between PCSK9 and TF plasma levels ⁵⁸	Reduced blood clotting activation	Enhanced monocyte LRP1-mediated TF-TFPI complex clearance

Abbreviations: FVIII, factor VIII; HDL, high-density lipoprotein; LDL, low-density lipoprotein; LDLR, low-density lipoprotein receptor; Lp(a), lipoprotein (a); LRP1, lipoprotein receptor-related protein 1; TF, tissue factor; TFPI, tissue factor pathway inhibitor; PCSK9, proprotein convertase subtilisin/kexin type-9.

adenovirus-induced over-expression of hepatic LDLR accelerated the clearance of FVIII.⁹⁹ These data show that LDLR cooperates with LRP1 in reducing FVIII levels, and indeed polymorphisms of the LDLR gene were found to affect CAD risk in humans by modulating FVIII:C levels and independently from the lipid profile.¹⁰⁰ Monocyte LRP1 favours also the clearance of TF by mediating the internalization and degradation of the TF-TFPI complex.¹⁰¹

Currently, there are no data showing that FVIII levels may be influenced by PCSK9 inhibitors. However, it is conceivable that PCSK9 inhibitors, by strikingly increasing LDLR expression, may enhance the clearance of FVIII, thus reducing its plasma levels, a mechanism potentially contributing to the lowering of major adverse cardiovascular events (►Table 1). In this regard, it is relevant that data from the Multi-Ethnic Study of Atherosclerosis, a cohort study of healthy subjects free of clinical cardiovascular disease, showed that statin users had significantly lower FVIII levels.¹⁰² Moreover, a recent multi-centre, randomized, controlled, open-label study in patients with prior deep vein thrombosis, showed that a short-term course of high-dose rosuvastatin significantly reduced FVIII.¹⁰³ Statins besides inhibiting of HMG-CoA reductase, also increase LDLR¹⁰⁴ and, at least atorvastatin, also LRP1.¹⁰⁵ Therefore, FVIII reduction by statins is likely explained by the increased expression of LDLR and LRP1 and the resulting accelerated hepatic clearance of FVIII.

Although the hypothesis that PCSK9 inhibition may reduce FVIII levels by increasing its clearance seems plausible, experimental studies confirming it are required.

Conclusion

Several clues suggest that PCSK9 represents a major actor in cardiovascular disease, in part independently from its effects on lipid metabolism. Data from observational studies in humans and from experimental research in animals imply that PCSK9 may modulate both primary and secondary haemostasis either indirectly, through its effect on LDL-c, or directly by influencing platelet activation and plasma levels of FVIII. Previous observations, showing that the benefits of statins on cardiovascular events occur before any significant changes in lipid profile have taken place,¹⁹

have opened the way to a series of studies which unravelled several cholesterol-independent actions of this class of drugs, including the stabilization of atherosclerotic plaques, the improvement of endothelial function, the modulation of immune responses, the inhibition of oxidative stress and inflammation and the prevention of thrombosis. These studies not only have led to an advancement in our understanding of the role of inflammation in atherothrombosis, but also to the development of innovative therapeutic approaches targeting inflammation, like the anti-interleukin-1 β MoAb canakinumab.¹⁰⁶ A complete unravelling of the possible pleiotropic activities of PCSK9 inhibitors, and in particular of their possible anti-thrombotic effects, may potentially widen the indications for this new therapeutic class and clarify their potential role in the treatment of the acute phase of ischaemic cardiovascular disease. The recent publication of the ODISSEY Trial shows that alirocumab reduced cardiovascular events and mortality in patients with a recent acute coronary syndrome and not on target for LDL-c on statin therapy, confirming that PCSK9 inhibitor therapy has an important role in secondary cardiovascular prevention in patients at high risk.¹⁰⁷

Funding

This work was supported in part by a grant from Regione Umbria (Progetto di ricerca finalizzata - BANDO 2013) to P.G.

Conflict of Interest

None declared.

Acknowledgements

We thank Dr. A.S. Weyrich and Dr. R.A. Campbell (Department of Internal Medicine, Utah University, Salt Lake City, United States) for sharing unpublished data on megakaryocyte transcriptome.

References

- Goldstein JL, Brown MS. Regulation of low-density lipoprotein receptors: implications for pathogenesis and therapy of hypercholesterolemia and atherosclerosis. *Circulation* 1987;76 (03):504-507

- 2 Grundy SM, Stone NJ, Bailey AL, et al. AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA Guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. *J Am Coll Cardiol* 2018. Doi: 10.1016/j.jacc.2018.11.002
- 3 Santos-Gallego CG, Badimon JJ, Rosenson RS. Beginning to understand high-density lipoproteins. *Endocrinol Metab Clin North Am* 2014;43(04):913–947
- 4 Pedersen TR. The success story of LDL cholesterol lowering. *Circ Res* 2016;118(04):721–731
- 5 Segrest JP, Jones MK, De Loof H, Dashti N. Structure of apolipoprotein B-100 in low density lipoproteins. *J Lipid Res* 2001;42(09):1346–1367
- 6 Brown MS, Anderson RG, Basu SK, Goldstein JL. Recycling of cell-surface receptors: observations from the LDL receptor system. *Cold Spring Harb Symp Quant Biol* 1982;46(Pt 2):713–721
- 7 Brown MS, Goldstein JL. A receptor-mediated pathway for cholesterol homeostasis. *Science* 1986;232(4746):34–47
- 8 Zhang DW, Garuti R, Tang WJ, Cohen JC, Hobbs HH. Structural requirements for PCSK9-mediated degradation of the low-density lipoprotein receptor. *Proc Natl Acad Sci U S A* 2008;105(35):13045–13050
- 9 Abifadel M, Varret M, Rabès JP, et al. Mutations in PCSK9 cause autosomal dominant hypercholesterolemia. *Nat Genet* 2003;34(02):154–156
- 10 Kent ST, Rosenson RS, Avery CL, et al. PCSK9 loss-of-function variants, low-density lipoprotein cholesterol, and risk of coronary heart disease and stroke: data from 9 studies of blacks and whites. *Circ Cardiovasc Genet* 2017;10(04):e001632
- 11 Maxwell KN, Breslow JL. Adenoviral-mediated expression of Pcsk9 in mice results in a low-density lipoprotein receptor knockout phenotype. *Proc Natl Acad Sci U S A* 2004;101(18):7100–7105
- 12 Lagace TA. PCSK9 and LDLR degradation: regulatory mechanisms in circulation and in cells. *Curr Opin Lipidol* 2014;25(05):387–393
- 13 Sabatine MS, Giugliano RP, Keech AC, et al; FOURIER Steering Committee and Investigators. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med* 2017;376(18):1713–1722
- 14 Navarese EP, Kolodziejczak M, Schulze V, et al. Effects of proprotein convertase subtilisin/kexin type 9 antibodies in adults with hypercholesterolemia: a systematic review and meta-analysis. *Ann Intern Med* 2015;163(01):40–51
- 15 Ridker PM. Mortality differences associated with treatment responses in CANTOS and FOURIER: insight and implications. *Circulation* 2018;137(17):1763–1766
- 16 Liao JK, Laufs U. Pleiotropic effects of statins. *Annu Rev Pharmacol Toxicol* 2005;45:89–118
- 17 Ridker PM, Danielson E, Fonseca FA, et al; JUPITER Study Group. Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N Engl J Med* 2008;359(21):2195–2207
- 18 Baigent C, Keech A, Kearney PM, et al; Cholesterol Treatment Trialists' (CTT) Collaborators. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins. *Lancet* 2005;366(9493):1267–1278
- 19 Ludman A, Venugopal V, Yellon DM, Hausenloy DJ. Statins and cardioprotection—more than just lipid lowering? *Pharmacol Ther* 2009;122(01):30–43
- 20 Blum A, Shamburek R. The pleiotropic effects of statins on endothelial function, vascular inflammation, immunomodulation and thrombogenesis. *Atherosclerosis* 2009;203(02):325–330
- 21 Krysiak R, Okopień B, Herman Z. Effects of HMG-CoA reductase inhibitors on coagulation and fibrinolysis processes. *Drugs* 2003;63(17):1821–1854
- 22 Antonopoulos AS, Margaritis M, Shirodaria C, Antoniadou C. Translating the effects of statins: from redox regulation to suppression of vascular wall inflammation. *Thromb Haemost* 2012;108(05):840–848
- 23 Oesterle A, Laufs U, Liao JK. Pleiotropic effects of statins on the cardiovascular system. *Circ Res* 2017;120(01):229–243
- 24 Silverman MG, Ference BA, Im K, et al. Association between lowering LDL-C and cardiovascular risk reduction among different therapeutic interventions: a systematic review and meta-analysis. *JAMA* 2016;316(12):1289–1297
- 25 Filippatos TD, Kei A, Rizos CV, Elisaf MS. Effects of PCSK9 inhibitors on other than low-density lipoprotein cholesterol lipid variables. *J Cardiovasc Pharmacol Ther* 2018;23(01):3–12
- 26 Gaudet D, Kereiakes DJ, McKenney JM, et al. Effect of alirocumab, a monoclonal proprotein convertase subtilisin/kexin 9 antibody, on lipoprotein(a) concentrations (a pooled analysis of 150 mg every two weeks dosing from phase 2 trials). *Am J Cardiol* 2014;114(05):711–715
- 27 Bittner V. Pleiotropic effects of PCSK9 (proprotein convertase subtilisin/kexin type 9) inhibitors? *Circulation* 2016;134(22):1695–1696
- 28 Santos-Gallego CG, Picatoste B, Badimón JJ. Pathophysiology of acute coronary syndrome. *Curr Atheroscler Rep* 2014;16(04):401–409
- 29 Santos-Gallego CG, Bayón J, Badimón JJ. Thrombi of different pathologies: implications for diagnosis and treatment. *Curr Treat Options Cardiovasc Med* 2010;12(03):274–291
- 30 Gresle P, Momi S, Miglisci R. Endothelium, venous thromboembolism and ischaemic cardiovascular events. *Thromb Haemost* 2010;103(01):56–61
- 31 Miglisci R, Becattini C, Pesavento R, et al. Endothelial dysfunction in patients with spontaneous venous thromboembolism. *Haematologica* 2007;92(06):812–818
- 32 Camera M, Rossetti L, Barbieri SS, et al. PCSK9 as a positive modulator of platelet activation. *J Am Coll Cardiol* 2018;71(08):952–954
- 33 Kassassir H, Siewiera K, Sychowski R, Watała C. Can the antiplatelet effects of cangrelor be reliably studied in mice under in vivo and in vitro conditions using flow cytometry? *Pharmacol Rep* 2013;65(04):870–883
- 34 Dwivedi DJ, Grin PM, Khan M, et al. Differential expression of PCSK9 modulates infection, inflammation, and coagulation in a murine model of sepsis. *Shock* 2016;46(06):672–680
- 35 Wang H, Wang Q, Wang J, et al. Proprotein convertase subtilisin/kexin type 9 (PCSK9) deficiency is protective against venous thrombosis in mice. *Sci Rep* 2017;7(01):14360
- 36 Emmerechts J, Vanassche T, Løyen S, et al. Partial versus complete factor VIII inhibition in a mouse model of venous thrombosis. *Thromb Res* 2012;129(04):514–519
- 37 Golder M, Mewburn J, Lillicrap D. In vitro and in vivo evaluation of the effect of elevated factor VIII on the thrombogenic process. *Thromb Haemost* 2013;109(01):53–60
- 38 Falcinelli E, Francisci D, Belfiori B, et al. In vivo platelet activation and platelet hyperreactivity in abacavir-treated HIV-infected patients. *Thromb Haemost* 2013;110(02):349–357
- 39 Navarese EP, Kolodziejczak M, Winter MP, et al. Association of PCSK9 with platelet reactivity in patients with acute coronary syndrome treated with prasugrel or ticagrelor: the PCSK9-REACT study. *Int J Cardiol* 2017;227:644–649
- 40 Gresle P, Falcinelli E, Momi S. Potentiation and priming of platelet activation: a potential target for antiplatelet therapy. *Trends Pharmacol Sci* 2008;29(07):352–360
- 41 Rossetti L, Ferri N, Marchiano S, et al. PCSK9 beyond its role in cholesterol homeostasis: co-activator of platelet function. *Eur Heart J* 2017;38:446–447
- 42 Gresle P, Guglielmini G, De Angelis M, et al. Acute, short-term hyperglycemia enhances shear stress-induced platelet activation in patients with type II diabetes mellitus. *J Am Coll Cardiol* 2003;41(06):1013–1020

- 43 Gresele P, Marzotti S, Guglielmini G, et al. Hyperglycemia-induced platelet activation in type 2 diabetes is resistant to aspirin but not to a nitric oxide-donating agent. *Diabetes Care* 2010;33(06):1262–1268
- 44 Rowley JW, Oler AJ, Tolley ND, et al. Genome-wide RNA-seq analysis of human and mouse platelet transcriptomes. *Blood* 2011;118(14):e101–e111
- 45 Cecchetti L, Tolley ND, Michetti N, Bury L, Weyrich AS, Gresele P. Megakaryocytes differentially sort mRNAs for matrix metalloproteinases and their inhibitors into platelets: a mechanism for regulating synthetic events. *Blood* 2011;118(07):1903–1911
- 46 Brambilla M, Facchinetti L, Canzano P, et al. Human megakaryocytes confer tissue factor to a subset of shed platelets to stimulate thrombin generation. *Thromb Haemost* 2015;114(03):579–592
- 47 Radziwon-Balicka A, Lesyk G, Back V, et al. Differential eNOS-signalling by platelet subpopulations regulates adhesion and aggregation. *Cardiovasc Res* 2017;113(14):1719–1731
- 48 Li S, Zhu CG, Guo YL, et al. The relationship between the plasma PCSK9 levels and platelet indices in patients with stable coronary artery disease. *J Atheroscler Thromb* 2015;22(01):76–84
- 49 Cariou B, Guérin P, Le May C, et al. Circulating PCSK9 levels in acute coronary syndrome: results from the PCSK9 prospective study. *Diabetes Metab* 2017;43(06):529–535
- 50 Fitzgerald DJ, Roy L, Catella F, Fitzgerald GA. Platelet activation in unstable coronary disease. *N Engl J Med* 1986;315(16):983–989
- 51 Gresele P, Falcinelli E, Loffredo F, et al. Platelets release matrix metalloproteinase-2 in the coronary circulation of patients with acute coronary syndromes: possible role in sustained platelet activation. *Eur Heart J* 2011;32(03):316–325
- 52 De Servi S, Crimi G, Calabrò P, et al. Relationship between diabetes, platelet reactivity, and the SYNTAX score to one-year clinical outcome in patients with non-ST-segment elevation acute coronary syndrome undergoing percutaneous coronary intervention. *EuroIntervention* 2016;12(03):312–318
- 53 Hu D, Yang Y, Peng DQ. Increased sortilin and its independent effect on circulating proprotein convertase subtilisin/kexin type 9 (PCSK9) in statin-naïve patients with coronary artery disease. *Int J Cardiol* 2017;227:61–65
- 54 Gustafsen C, Kjolby M, Nyegaard M, et al. The hypercholesterolemia-risk gene SORT1 facilitates PCSK9 secretion. *Cell Metab* 2014;19(02):310–318
- 55 Pastori D, Nocella C, Farcomeni A, et al; ATHERO-AF Study Group. Relationship of PCSK9 and urinary thromboxane excretion to cardiovascular events in patients with atrial fibrillation. *J Am Coll Cardiol* 2017;70(12):1455–1462
- 56 Catella F, Fitzgerald GA. Paired analysis of urinary thromboxane B2 metabolites in humans. *Thromb Res* 1987;47(06):647–656
- 57 Tatsumi K, Mackman N. Tissue factor and atherothrombosis. *J Atheroscler Thromb* 2015;22(06):543–549
- 58 Wang M, Li YF, Guo YG, Chen MM, Jiang ZL, Song JY. Positive correlation between plasma PCSK9 and tissue factors levels in patients with angiographically diagnosed coronary artery disease and diabetes mellitus. *J Geriatr Cardiol* 2016;13(04):312–315
- 59 Ochoa E, Iriando M, Manzano C, et al. LDLR and PCSK9 are associated with the presence of antiphospholipid antibodies and the development of thrombosis in aPLA carriers. *PLoS One* 2016;11(01):e0146990
- 60 Urbanus RT, de Laat B. Antiphospholipid antibodies and the protein C pathway. *Lupus* 2010;19(04):394–399
- 61 Bontadi A, Ruffatti A, Falcinelli E, et al. Platelet and endothelial activation in catastrophic and quiescent antiphospholipid syndrome. *Thromb Haemost* 2013;109(05):901–908
- 62 Obermayer G, Afonyushkin T, Binder CJ. Oxidized low-density lipoprotein in inflammation-driven thrombosis. *J Thromb Haemost* 2018;16(03):418–428
- 63 Chen K, Febbraio M, Li W, Silverstein RL. A specific CD36-dependent signaling pathway is required for platelet activation by oxidized low-density lipoprotein. *Circ Res* 2008;102(12):1512–1519
- 64 Hofmann A, Brunssen C, Morawietz H. Contribution of lectin-like oxidized low-density lipoprotein receptor-1 and LOX-1 modulating compounds to vascular diseases. *Vascul Pharmacol* 2017. Doi: 10.1016/j.vph.2017.10.002
- 65 Magwenzi S, Woodward C, Wraith KS, et al. Oxidized LDL activates blood platelets through CD36/NOX2-mediated inhibition of the cGMP/protein kinase G signaling cascade. *Blood* 2015;125(17):2693–2703
- 66 Yoshimoto R, Fujita Y, Kakino A, Iwamoto S, Takaya T, Sawamura T. The discovery of LOX-1, its ligands and clinical significance. *Cardiovasc Drugs Ther* 2011;25(05):379–391
- 67 Marwali MR, Hu CP, Mohandas B, et al. Modulation of ADP-induced platelet activation by aspirin and pravastatin: role of lectin-like oxidized low-density lipoprotein receptor-1, nitric oxide, oxidative stress, and inside-out integrin signaling. *J Pharmacol Exp Ther* 2007;322(03):1324–1332
- 68 Carnevale R, Bartimoccia S, Nocella C, et al. LDL oxidation by platelets propagates platelet activation via an oxidative stress-mediated mechanism. *Atherosclerosis* 2014;237(01):108–116
- 69 Biswas S, Xin L, Panigrahi S, et al. Novel phosphatidylethanolamine derivatives accumulate in circulation in hyperlipidemic ApoE^{-/-} mice and activate platelets via TLR2. *Blood* 2016;127(21):2618–2629
- 70 Leibundgut G, Scipione C, Yin H, et al. Determinants of binding of oxidized phospholipids on apolipoprotein (a) and lipoprotein (a). *J Lipid Res* 2013;54(10):2815–2830
- 71 Riches K, Porter KE. Lipoprotein(a): cellular effects and molecular mechanisms. *Cholesterol* 2012;2012:923289
- 72 Calkin AC, Drew BG, Ono A, et al. Reconstituted high-density lipoprotein attenuates platelet function in individuals with type 2 diabetes mellitus by promoting cholesterol efflux. *Circulation* 2009;120(21):2095–2104
- 73 Shattil SJ, Anaya-Galindo R, Bennett J, Colman RW, Cooper RA. Platelet hypersensitivity induced by cholesterol incorporation. *J Clin Invest* 1975;55(03):636–643
- 74 Nofer JR, Walter M, Kehrel B, et al. HDL3-mediated inhibition of thrombin-induced platelet aggregation and fibrinogen binding occurs via decreased production of phosphoinositide-derived second messengers 1,2-diacylglycerol and inositol 1,4,5-trisphosphate. *Arterioscler Thromb Vasc Biol* 1998;18(06):861–869
- 75 Riddell DR, Graham A, Owen JS. Apolipoprotein E inhibits platelet aggregation through the L-arginine:nitric oxide pathway. Implications for vascular disease. *J Biol Chem* 1997;272(01):89–95
- 76 van der Stoep M, Korpelaar SJ, Van Eck M. High-density lipoprotein as a modulator of platelet and coagulation responses. *Cardiovasc Res* 2014;103(03):362–371
- 77 Panes O, González C, Hidalgo P, et al. Platelet tissue factor activity and membrane cholesterol are increased in hypercholesterolemia and normalized by rosuvastatin, but not by atorvastatin. *Atherosclerosis* 2017;257:164–171
- 78 Ding Z, Liu S, Wang X, et al. Cross-talk between LOX-1 and PCSK9 in vascular tissues. *Cardiovasc Res* 2015;107(04):556–567
- 79 Barale C, Bonomo K, Noto F, et al. Effects of PCSK9 inhibitors on platelet function in adults with hypercholesterolemia. *Atherosclerosis* 2017;263:e30–e31
- 80 Kotani K, Banach M. Lipoprotein(a) and inhibitors of proprotein convertase subtilisin/kexin type 9. *J Thorac Dis* 2017;9(01):E78–E82
- 81 Geddings JE, Mackman N. New players in haemostasis and thrombosis. *Thromb Haemost* 2014;111(04):570–574
- 82 Ding Z, Liu S, Wang X, et al. Cross-talk between PCSK9 and damaged mtDNA in vascular smooth muscle cells: role in apoptosis. *Antioxid Redox Signal* 2016;25(18):997–1008
- 83 Li J, Liang X, Wang Y, Xu Z, Li G. Investigation of highly expressed PCSK9 in atherosclerotic plaques and ox-LDL-induced endothelial cell apoptosis. *Mol Med Rep* 2017;16(02):1817–1825

- 84 Kaufman RJ, Pipe SW, Tagliavacca L, Swaroop M, Moussalli M. Biosynthesis, assembly and secretion of coagulation factor VIII. *Blood Coagul Fibrinolysis* 1997;8(02, Suppl 2):S3–S14
- 85 Bank I, Libourel EJ, Middeldorp S, et al. Elevated levels of FVIII:C within families are associated with an increased risk for venous and arterial thrombosis. *J Thromb Haemost* 2005;3(01):79–84
- 86 Siegler JE, Samai A, Albright KC, Boehme AK, Martin-Schild S. Factoring in factor VIII with acute ischemic stroke. *Clin Appl Thromb Hemost* 2015;21(07):597–602
- 87 Rumley A, Lowe GD, Sweetnam PM, Yarnell JW, Ford RP. Factor VIII, von Willebrand factor and the risk of major ischaemic heart disease in the Caerphilly Heart Study. *Br J Haematol* 1999;105(01):110–116
- 88 Tracy RP, Arnold AM, Ettinger W, Fried L, Meilahn E, Savage P. The relationship of fibrinogen and factors VII and VIII to incident cardiovascular disease and death in the elderly: results from the cardiovascular health study. *Arterioscler Thromb Vasc Biol* 1999;19(07):1776–1783
- 89 Folsom AR, Peacock JM, Boerwinkle E; Atherosclerosis Risk in Communities (ARIC) Study Investigators. Variation in PCSK9, low LDL cholesterol, and risk of peripheral arterial disease. *Atherosclerosis* 2009;202(01):211–215
- 90 Milgrom A, Lee K, Rothschild M, et al. Thrombophilia in 153 patients with premature cardiovascular disease $\leq$ age 45. *Clin Appl Thromb Hemost* 2018;24(02):295–302
- 91 Rosendaal FR, Briët E, Stibbe J, et al. Haemophilia protects against ischaemic heart disease: a study of risk factors. *Br J Haematol* 1990;75(04):525–530
- 92 Machlus KR, Lin FC, Wolberg AS. Procoagulant activity induced by vascular injury determines contribution of elevated factor VIII to thrombosis and thrombus stability in mice. *Blood* 2011;118(14):3960–3968
- 93 Kawasaki T, Kaida T, Arnout J, Vermynen J, Hoylaerts MF. A new animal model of thrombophilia confirms that high plasma factor VIII levels are thrombogenic. *Thromb Haemost* 1999;81(02):306–311
- 94 Brummel-Ziedins K, Undas A, Orfeo T, et al. Thrombin generation in acute coronary syndrome and stable coronary artery disease: dependence on plasma factor composition. *J Thromb Haemost* 2008;6(01):104–110
- 95 Rice GI, Grant PJ. FVIII coagulant activity and antigen in subjects with ischaemic heart disease. *Thromb Haemost* 1998;80(05):757–762
- 96 Lenting PJ, Neels JG, van den Berg BM, et al. The light chain of factor VIII comprises a binding site for low density lipoprotein receptor-related protein. *J Biol Chem* 1999;274(34):23734–23739
- 97 Saenko EL, Yakhyayev AV, Mikhailenko I, Strickland DK, Sarafanov AG. Role of the low density lipoprotein-related protein receptor in mediation of factor VIII catabolism. *J Biol Chem* 1999;274(53):37685–37692
- 98 Bovenschen N, Herz J, Grimbergen JM, et al. Elevated plasma factor VIII in a mouse model of low-density lipoprotein receptor-related protein deficiency. *Blood* 2003;101(10):3933–3939
- 99 Bovenschen N, Mertens K, Hu L, Havekes LM, van Vlijmen BJ. LDL receptor cooperates with LDL receptor-related protein in regulating plasma levels of coagulation factor VIII in vivo. *Blood* 2005;106(03):906–912
- 100 Martinelli N, Girelli D, Lunghi B, et al. Polymorphisms at LDLR locus may be associated with coronary artery disease through modulation of coagulation factor VIII activity and independently from lipid profile. *Blood* 2010;116(25):5688–5697
- 101 Hamik A, Setiadi H, Bu G, McEver RP, Morrissey JH. Down-regulation of monocyte tissue factor mediated by tissue factor pathway inhibitor and the low density lipoprotein receptor-related protein. *J Biol Chem* 1999;274(08):4962–4969
- 102 Adams NB, Lutsey PL, Folsom AR, et al. Statin therapy and levels of hemostatic factors in a healthy population: the Multi-Ethnic Study of Atherosclerosis. *J Thromb Haemost* 2013;11(06):1078–1084
- 103 Biedermann JS, Kruij MJHA, van der Meer FJ, et al. Rosuvastatin use improves measures of coagulation in patients with venous thrombosis. *Eur Heart J* 2018;39(19):1740–1747
- 104 Tiwari V, Khokhar M. Mechanism of action of anti-hypercholesterolemia drugs and their resistance. *Eur J Pharmacol* 2014;741:156–170
- 105 Moon JH, Kang SB, Park JS, et al. Up-regulation of hepatic low-density lipoprotein receptor-related protein 1: a possible novel mechanism of antiatherogenic activity of hydroxymethylglutaryl-coenzyme A reductase inhibitor Atorvastatin and hepatic LRP1 expression. *Metabolism* 2011;60(07):930–940
- 106 Ridker PM, Everett BM, Thuren T, et al; CANTOS Trial Group. Antiinflammatory therapy with canakinumab for atherosclerotic disease; CANTOS Trial Group. *N Engl J Med* 2017;377(12):1119–1131
- 107 Schwartz GG, Steg PG, Szarek M, et al. Alirocumab and cardiovascular outcomes after acute coronary syndrome. *N Engl J Med* 2018;379:2097–2107