Histomorphometric Comparative Analysis between Oral Dysplastic Potentially Malignant Disorders and Oral Squamous Cell Carcinoma

Natalia Da Silva Pereira¹ Tiago Novaes Pinheiro²

¹School of Health Sciences, Department of Dentistry, Amazonas State University, Manaus, Amazonas, Brazil

²Department of Pathology, Amazonas State University, Manaus, Amazonas, Brazil

Address for correspondence Tiago Novaes Pinheiro, DDS, MSc, PhD. Department of Pathology, Amazonas State University, Manaus, Amazonas, Brazil (e-mail: tpinheiro@uea.edu.br).

Eur J Dent 2019;13:1–4

Abstract

Objective This study aimed to determine the correlation between the vascular mast cells and inflammatory changes in the potentially malignant disorders, compared to the oral squamous cell carcinoma (OSCC) in varying degrees of dysplasia.

Materials and Methods Thirty samples were selected: 10 of OSCC, 10 of oral leukoplakia, and 10 of actinic keratosis. The toluidine blue technique was used on the mast cells, and hematoxylin and eosin were used for analyzing the lymphocytes, mitosis, and vessels. The quantification was performed using the ImageJ software after obtaining the images by light microscopy with a $\times 40$ objective. Analysis of variance with $p < 0.05$ was considered for statistical significance.

Results Mast cells ($p < 0.0158$), vessels ($p < 0.9431$), lymphocytes ($p < 0.0001$), and mitoses ($p < 0.0009$) were found in OSCC. In potentially malignant disorders, a lower density of these structures and a higher concentration of mitosis in the actinic keratosis were observed.

Conclusion The results showed evidence of a positive correlation between mast cells and vascularization in the OSCC, suggesting the aggression of the disease. Intense inflammatory infiltrate indicates that other molecular events are involved in the carcinogenesis process, and further studies are necessary for a better understanding of it.

Keywords

- carcinogenesis
- lymphocytes
- mast cells
- mitosis
- vascularization

Introduction

Precancerous condition is defined as a biological condition that does not necessarily alter the local tissue clinical appearance, but is associated with a significantly increased risk of cancer. Oral squamous cell carcinoma is the most prevalent malignancy in the oral region worldwide.¹ Recent studies have investigated the role of mast cells in the infiltrate of these disorders contributing to their progression or to inhibit tumor growth.² This study aims to verify the inflammatory behavior of potentially malignant oral disorders such as oral leukoplakia (OL) and actinic keratosis (AK) as compared with oral squamous cell carcinoma (OSCC) in order to detect when these lesions may evolve in a malignant way, and to consequently identify early oral cancer.

Materials and Methods

The study was submitted to and approved by the Research Ethics Committee (2.253.859).

The developed search was an experimental comparative analysis, based on the material analysis of cases diagnosed in the pathology service of the authors' educational institution.

Thirty cases were analyzed, including 10 OSCC cases (group 1), 10 cases of OL (group 2), and 10 cases of AK (group 3). For inclusion in this research, the diagnosis of the lesions had to be conclusive, and the report had to contain the microscopic variables and local findings of the lesions, regardless of age and gender. In addition, specimens associated with other lesions or with insufficient amounts of tissue for analysis were excluded from the study. For

microscopic analysis, biopsied slides with 4- μ m sections that were previously prepared and stained with the hematoxylin were used. Each sample stained with HE was analyzed in 3 microscopic fields, and for mast cells quantification, the slides were stained with toluidine blue (0.1%) and analyzed in 10 microscopic fields, as shown in ►Fig. 1. Images of the histological samples were obtained by means of light microscopy (BIOVAL) with a $\times 40$ objective and optovar of $\times 1.6$, recorded by a Zeiss Axiocam 503 color camera, $\times 10$ objective, resulting in a total increase of $\times 640$. The images were saved on a computer and weighted with ImageJ Java software version 1.48 v [ImageJ software developer Java (open source), NIHUSA] through the Cell Counter plugin. The microscopic variables studied included the following:

- Dysplastic alterations
- Typical and atypical mitosis count
- Inflammatory infiltrate intensity
- Blood vessel count
- Mast cell count in the areas with higher concentrations

Dysplastic alterations were observed according to the cellular modifications, such as increased nucleoli, nuclear hyperchromatism, nuclear pleomorphism, increased nucleus/cytoplasm ratio, increased mitotic activity, and cellular pleomorphism. In addition, epithelial changes were observed, such as basal layer hyperplasia, hypercellularity, and abnormal stratification of the epithelium. The intensity of the inflammatory infiltrate was classified quantitatively

Fig. 1 Histopathological image showing mast cells in connective tissue (blue staining toluidine, $\times 400$).

and considered intense when the total cell count was above 121 and moderate with a count between 71 and 120, with discrete counts between 10 and 70.³

The analysis of variance was performed with the software StatCalc version 8.2.2 of AcaStat (AcaStat Software developer, Poinciana, Florida) and assigned $p < 0.05$ for statistical significance.

Results

An important statistical significance was observed between the groups. Higher concentrations of mast cells, vessels, and lymphocytes were seen in the OSCC group (group 1) with a statistical significance of $p < 0.0158$, $p < 0.9431$, and $p < 0.0001$, respectively, when compared to the OL (group 2) and AK (group 3) lesions. While in the mitotic figure quantification, the AK lesions showed higher counts, with $p < 0.0009$, as shown in ►Table 1 and represented in ►Fig. 2.

Discussion

For many years, mast cells have been considered the main agents of reactions such as allergic reactions through the release of bioactive cytokines, chemokines, proteases, leukotrienes, and polyamines. Emerging roles for mast cells have recently been identified, which highlight their relevance not only in innate immunity but also in adaptive immunity. Morphologically, they are oval with a central nucleus and a

Fig. 2 Graphical representation of the averages of mast cells, lymphocytes, blood vessels, and mitosis of group 1 (oral squamous cell carcinoma), group 2 (oral leukoplakia), and group 3 (actinic keratosis); SD: Standard deviation.

Table 1 Comparison of the means and standard deviation of mast cells, lymphocytes, blood vessels, and mitosis between oral squamous cell carcinoma, oral leukoplakia, and actinic keratosis

Groups	Mast Cells		Lymphocytes		Blood Vessels		Mitosis	
Group 1	35.5	22.2	189.3	137.8	12.2	15.9	2.2	2.1
Group 2	34.4	16.9	29	19.6	11.6	9.9	1.5	1.9
Group 3	15	6.8	19.8	18.2	10.5	5.3	8.6	6.5
ANOVA	$p < 0.0158$		$p < 0.0001$		$p < 0.9431$		$p < 0.0009$	

Abbreviations: OSCC: Oral squamous cell carcinoma, OL: Oral leukoplakia, AK: Actinic keratosis.

Group 1 (OSCC), group 2 (OL), group 3 (AK), P (statistical significance).

large amount of granules in the cytoplasm and are usually located close to the blood vessels.⁴

These cells have been associated with resistance to tumor growth-producing molecules such as IL-1, 4, and 6, and tumor necrosis factor alpha, as well as with benefits, by promoting expanded vascular delivery.⁵

The potential effects of mast cells on tumor growth according to Maltby et al's study⁶ can be categorized as direct effects on tumor cells, such as mast cell-mediated cytotoxicity, or indirect effects, such as mast cell-directed angiogenesis, tissue remodeling of the surrounding environment, and recruitment of immune cells. In vitro studies have shown that the addition of whole mast cells may lead to neovascularization, which requires granular components of mast cells to perform, whereas the presence of mast cell membranes alone is insufficient. In addition, stem cell factor (SCF) produced by excess tumor cells has been implicated in the accumulation of mast cells in the periphery of developing tumors and in the increased production of VEGF.

Neto et al⁷ observed that mast cell deficiency in the inflammatory infiltrate increased the incidence of tumors, suggesting a very important positive role of these cells in immunological defense against tumors.

In the present study, we found higher densities of mast cells, blood vessels, and intense inflammatory infiltrate and a relatively small number of mitoses in OSCC than in AK and OL. Recent studies have shown similar results. Ingaleswar et al⁸ carried out a study by analyzing vessel and mast cell density in OSCC compared to the healthy mucosa and found a linear increase of vessels as the density of mast cells increased in malignant lesions, suggesting a positive correlation. Another similar study by Iamaroon et al⁹ compared the correlation between the mast cells and microvessel quantity in OSCC, cancerous dysplasia, hyperkeratotic lesions, and clinically healthy mucosa and revealed increased densities of these histopathological structures based on the progression of the disease, that is, the count was higher in carcinoma when compared to the other lesions and the normal oral mucosa. This is due to the production of several mediators. In addition, heparin, which is the dominant proteoglycan in mast cells, has many properties, one of which is mitogenic power on endothelial cells to stimulate the migration of endothelial cells.⁸ A study carried out by Pittoni et al¹⁰ supported the idea that the high density of the inflammatory infiltrate found in this research may also contribute to tumor growth through the release of MMP-9, a proteinase secreted by inflammatory cells, including mast cells that cleave the SCF receptor on the tumor cell membrane, which increases the attraction between these cells through the interaction between this receptor and the c-kit found in mast cells. Another study by Lopamudra das Roy et al¹¹ confirmed the above results by showing the increased recruitment of mast cells within tumors from mice with arthritis, in which metastasis in the lungs and bones was facilitated.

Other studies have shown divergent results. StaSikOWS-ka-Kanicka et al¹² also associated the density of infiltrative cells and microvessels in OSCC and observed a significantly lower number of mast cells in lesions with unfavorable

prognoses when compared to those with favorable prognoses and proposed that the deficiency of these cells in malignant lesions is associated with poor prognosis.

Sathyakumar et al¹³ observed a direct relationship between the density of mast cells and blood vessels when comparing normal mucosa to leukoplakia with and without dysplasia, showing progressive increases in density, being lowering in normal oral mucosa and higher in lesions with dysplasia. Despite the fact that there is a positive correlation between these histological components, Ali Tahir et al¹⁴ believed that it is not direct since it is not only the mast cells that release proinflammatory enzymes. Similar results were observed by Michailidou et al¹⁵ and Rojas et al¹⁶. The first suggested a positive association between mast cell concentration and vascularization in the normal buccal mucosa, leukoplakia with and without dysplasia, and OSCC, with a linear increase in concentrations, that is, a lower concentration in the normal oral mucosa and higher concentrations in leukoplakias with dysplasia and carcinoma. Rojas et al evaluated clinically healthy lips and AK and observed an increase in mast cells in keratosis, especially in the areas close to solar elastosis, compared to the normal oral mucosa.

The reduced number of mast cells observed in the lesions of OL by Neto et al⁷ is equivalent to the results found in this study, but diverges regarding reduced mast cell count in OSCC. Lachter et al¹⁷ found similar results when observing fewer mast cells in cancerous alterations and malignant neoplasms in colorectal lesions when compared with normal colorectal tissue.

Thus, in this research, significantly high mast cell density, lymphocyte vascularization, and a low mitosis count were observed in the carcinoma, whereas decreases in these factors were observed in OL and AK, with the exception of AK, which showed a significant increase of mitosis.

The decrease in the number of mast cells in these lesions may be related to the migration failure of these cells, reflecting an important modification in the microenvironment during the initiation and progression of a tumor. Furthermore, the lesion may be in the second phase of the carcinogenesis process, characterized by self-regulation of tumor cell growth.¹⁸ Other studies have shown that there is research that further reveals that the reduction of the expression of E-cad, tumor suppressor gene that is expressed in the epithelial tissues, may increase tumor progression.¹⁹

Conclusion

In this study, mast cell and microvessel densities were significantly increased in OSCC lesions, suggesting tumor progression and aggression through the positive regulation of angiogenesis by these cells. However, mast cells are only part of this complex process of carcinogenesis, together with other factors secreted in the tumor environment that can modulate angiogenesis. Further studies should be conducted to contribute to a better understanding of these events.

Financial Support and Sponsorship

None.

Conflicts of Interest

None.

References

- Vigneswaran N, Williams MD. Epidemiologic trends in head and neck cancer and aids in diagnosis. *Oral Maxillofac Surg Clin North Am* 2014;26(2):123–141
- Gangwar RS, Friedman S, Seaf M, Levi-Schaffer F. Mast cells and eosinophils in allergy: close friends or just neighbors. *Eur J Pharmacol* 2016;778:77–83
- Sharma B, Sriram G, Saraswathi TR, Sivapathasundharam B. Immunohistochemical evaluation of mast cells and angiogenesis in oral squamous cell carcinoma. *Indian J Dent Res* 2010;21(2):260–265
- Shea-Donohue T, Stiltz J, Zhao A, Notari L. Mast cells. *Curr Gastroenterol Rep* 2010;12(5):349–357
- Abbas AK, Lichtman AH, Pillai S. *Cellular and Molecular Immunology*. 8th ed. Philadelphia, PA: Elsevier; 2014:544
- Maltby S, Khazaie K, McNagny KM. Mast cells in tumor growth: angiogenesis, tissue remodelling and immune-modulation. *Biochim Biophys Acta* 2009;1796(1):19–26
- Oliveira-Neto HH, Leite AF, Costa NL, et al. Decrease in mast cells in oral squamous cell carcinoma: possible failure in the migration of these cells. *Oral Oncol* 2007;43(5):484–490
- Ingaleswar PS, Pandit S, Desai D, Redder CP, Shetty AS, Mithun KM. Immunohistochemical analysis of angiogenesis by CD34 and mast cells by toluidine blue in different grades of oral squamous cell carcinoma. *J Oral Maxillofac Pathol* 2016;20(3):467–473
- Iamaroon A, Pongsiriwet S, Jittidecharaks S, Pattanaporn K, Prapayasatok S, Wanachantararak S. Increase of mast cells and tumor angiogenesis in oral squamous cell carcinoma. *J Oral Pathol Med* 2003;32(4):195–199
- Pittoni P, Piconese S, Tripodo C, Colombo MP. Tumor-intrinsic and -extrinsic roles of c-Kit: mast cells as the primary off-target of tyrosine kinase inhibitors. *Oncogene* 2011;30(7):757–769
- Das Roy L, Curry JM, Sahraei M, et al. Arthritis augments breast cancer metastasis: role of mast cells and SCF/c-Kit signaling. *Breast Cancer Res* 2013;15(2):R32
- Stasikowska-Kanicka O, Wągrowaska-Danilewicz M, Danilewicz M. Association of infiltrating cells with microvessel density in oral squamous cell carcinoma. *Pol J Pathol* 2017;68(1):40–48
- Sathyakumar M, Sriram G, Saraswathi T, Sivapathasundharam B. Immunohistochemical evaluation of mast cells and vascular endothelial proliferation in oral precancerous lesion-leukoplakia. *J Oral Maxillofac Pathol* 2012;16(3):343–348
- Tahir A, Nagi AH, Ullah E, Janjua OS. The role of mast cells and angiogenesis in well-differentiated oral squamous cell carcinoma. *J Cancer Res Ther* 2013;9(3):387–391
- Michailidou EZ, Markopoulos AK, Antoniadis DZ. Mast cells and angiogenesis in oral malignant and premalignant lesions. *Open Dent J* 2008;2:126–132
- Rojas IG, Martínez A, Pineda A, Spencer ML, Jiménez M, Rudolph MI. Increased mast cell density and protease content in actinic cheilitis. *J Oral Pathol Med* 2004;33(9):567–573
- Lachter J, Stein M, Lichtig C, Eidelman S, Munichor M. Mast cells in colorectal neoplasias and premalignant disorders. *Dis Colon Rectum* 1995;38(3):290–293
- Anuradha A, Kiran Kumar Naik B, Vijay Srinivas G, Devi RS, Puneet HK. Incidence of mast cells in oral squamous cell carcinoma: a short study. *J Oncol* 2014;2014:614291
- Sridevi U, Jain A, Nagalaxmi V, Kumar UV, Goyal S. Expression of E-cadherin in normal oral mucosa, in oral precancerous lesions and in oral carcinomas. *Eur J Dent* 2015;9(3):364–372