D. A. EVANS*, S. J. MILLER, T. LECTKA (HARVARD UNIVERSITY, CAMBRIDGE, USA)
 Bis(oxazoline)copper(II) Complexes as Chiral Catalysts for the Enantioselective Diels–Alder Reaction
J. Am. Chem. Soc. **1993**, *115*, 6460–6461.

Enantioselective Copper-Catalyzed Diels–Alder Cycloaddition

Examples:

Double stereodifferentiating experiments:

Significance: Following Narasaka's report on titanium-catalyzed enantioselective Diels–Alder reactions (*J. Am. Chem. Soc.* **1989**, *111*, 5340) and Corey's variants using chiral magnesium(II) and iron(II) catalysts (*J. Am. Chem. Soc.* **1991**, *113*, 728; *Tetrahedron Lett.* **1992**, *33*, 6807), Evans disclosed a highly enantioselective Diels–Alder reaction employing copper(II)-catalysts and BOX ligands.

Comment: Products were obtained in high yields and excellent enantioselectivities. This class of chiral Lewis acid catalysts has since been applied to asymmetric aldol reactions, Michael additions, and carbonyl ene reactions by the same group (J. S. Johnson, D. A. Evans *Acc. Chem. Res.* **2000**, *33*, 325).

SYNFACTS Contributors: Mark Lautens, José F. Rodríguez
 Synfacts 2019, 15(07), 0753 Published online: 17.06.2019
 DOI: 10.1055/s-0039-1689962; Reg-No.: L07019SF

© 2019, Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Metals in Synthesis

Key words

copper catalysis

Diels–Alder
cycloaddition

Lewis acids

Synfact
Classic