J. M. FRANKE, B. K. RALISKI, S. C. BOGGESE, D. V. NATESAN, E. T. KORETSKY, P. ZHANG, R. U. KULKARNI, P. E. DEAL, E. W. MILLER* (UNIVERSITY OF CALIFORNIA, BERKELEY, USA)

BODIPY Fluorophores for Membrane Potential Imaging

J. Am. Chem. Soc. **2019**, *141*, 12824–12831.

Water-Soluble BODIPY Dyes for Voltage Imaging

Category

Chemistry in
Medicine and
Biology

Key words

voltage dyes

fluorophores

BODIPY

Synfact
of the
Month

Synthesis of water-soluble BODIPY VF dyes:

BODIPY VF series:

Significance: Previously, water solubility of BODIPY dyes was achieved through direct functionalization of the BODIPY fluorophore. Miller and co-workers report an elegant and versatile solution that does not significantly alter the dye electronics through introduction of a sulfonate on the *meso*-aromatic ring. These fluorophores are used to generate a series of new VoltageFluor (VF) dyes that enable the recording of action potential dynamics in neurons and cardiomyocytes.

Comment: The TFA-catalyzed condensation of aldehydes with pyrroles in the synthesis of BODIPY dyes is commonly performed in CH_2Cl_2 and often reported to be low-yielding. Here, the authors optimized this reaction with their sulfonated aldehyde, which was found to be insoluble in CH_2Cl_2 and toluene. Comparably good yields were obtained in DMF (e.g. 49% over three steps, see above). Therefore, the reported strategy for water-soluble BODIPY synthesis could be very useful for the wider community.

SYNFACTS Contributors: Dirk Trauner, Johannes Morstein
Synfacts 2019, 15(11), 1309 Published online: 18.10.2019
DOI: 10.1055/s-0039-1690291; Reg-No.: T10519SF

© 2019, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.