A. B. CHARETTE*, H. JUTEAU (UNIVERSITÉ DE MONTRÉAL, CANADA)

Design of Amphoteric Bifunctional Ligands: Application to the Enantioselective Simmons–Smith Cyclopropanation of Allylic Alcohols

J. Am. Chem. Soc. **1994**, *116*, 2651–2652.

Enantioselective Simmons–Smith Cyclopropanation: The Charette Modification

Demonstrated examples:

Postulated mechanism:

Significance: In the paper, Charette and Juteau described a route towards enantioenriched cyclopropanes using a chiral amphoteric bifunctional ligand. This was the first report that enabled efficient chiral cyclopropane synthesis without reliance on covalently bonded chiral auxiliaries.

Comment: This modification tolerates various functional groups and is highly enantioselective. The chiral ligand could also be recovered through an aqueous extraction at the end of the reaction.

Reviews: A. B. Charette, J.-F. Marcoux *Synlett* **1995**, 1197–1207; H. Lebel, J.-F. Marcoux, C. Molinaro, A. B. Charette *Chem. Rev.* **2003**, *103*, 977–1050.

SYNFACTS Contributors: Mark Lautens, Egor M. Larin
Synfacts 2019, 15(09), 1013 Published online: 20.08.2019
DOI: 10.1055/s-0039-1690581; Reg-No.: L10019SF

© 2019, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Metals in Synthesis

Key words

zinc

cyclopropanation

Simmons–Smith
cyclopropanation

Synfact
Classic

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.