S. SITTIHAN, T. F. JAMISON* (MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, USA)

Total Synthesis of the Marine Ladder Polyether Gymnocin B

J. Am. Chem. Soc. **2019**, *141*, 11239–11244.

Total Synthesis of Gymnocin B

Epoxide-opening cascades:

Elaborated coupling fragments:

Tetrahedron Lett. **2010**, *51*, 4673

Target compound:

Significance: Sittihan and Jamison report the total synthesis of gymnocin B, the second largest contiguous marine ladder polyether isolated to date. This complex target features 15 cyclic ether rings and 33 stereogenic centers, posing an exceptional synthetic challenge. The authors succeeded in the preparation of gymnocin B by implementation of bioinspired epoxide-opening cascades and a series of fragment couplings and cyclizations.

Comment: Key components **D**, **E**, and **F** were prepared from easily accessible compounds by exquisitely controlled epoxide-opening cascades. Further elaboration of these building blocks allowed combination of subunits **G–J** and construction of the remaining rings to afford the exceedingly challenging target gymnocin B.

SYNFACTS Contributors: Erick M. Carreira, Niels Sievertsen
Synfacts 2019, 15(10), 1089 Published online: 17.09.2019
DOI: 10.1055/s-0039-1690646; Reg-No.: C05219SF

© 2019, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Synthesis of Natural
Products and
Potential Drugs

Key words

gymnocin B

marine ladder
polyether

epoxide-opening
cascade

modular synthesis

Synfact
of the
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.