D. P. CURRAN*, D. M. RAKIEWICZ (UNIVERSITY OF PITTSBURGH, USA)

Tandem Radical Approach to Linear Condensed Cyclopentanoids. Total Synthesis of (±)-Hirsutene

J. Am. Chem. Soc. **1985**, *107*, 1448–1449.

D. P. CURRAN*, D. M. RAKIEWICZ (UNIVERSITY OF PITTSBURGH, USA)

Radical-Initiated Polyolefinic Cyclizations in Linear Triquinane Synthesis. Model Studies and Total Synthesis of (±)-Hirsutene
Tetrahedron **1985**, *41*, 3943–3958.

Synthesis of (±)-Hirsutene

Significance: Hirsutene is a linear triquinane that was isolated from the fungi *Coriolus consors* in 1976. Its lack of heteroatomic functional groups along with the presence of four contiguous stereocenters and two quaternary carbon atoms render this condensed cyclopentanoid natural product challenging. In 1985, Curran and Rakiewicz reported an elegant synthesis of (±)-hirsutene employing a late-stage radical cyclization.

Comment: Accessed in two steps from the corresponding enone, acetate **A** was converted into lactone **B** by Ireland–Claisen rearrangement followed by phenylselenolactonization. Elimination, S_N2' reaction, and deprotection furnished carboxylic acid **E**. Further transformation led to iodoenone **H**, which underwent 5-*exo*-trig/5-*exo*-dig radical cyclization cascade in the presence of AIBN/Bu₃SnH to afford (±)-hirsutene.

SYNFACTS Contributors: Erick M. Carreira, Matthieu J. R. Richter

Synfacts 2019, 15(10), 1090 Published online: 17.09.2019

DOI: 10.1055/s-0039-1690649; Reg-No.: C05619SF