Long Noncoding RNAs of the Arterial Wall as Therapeutic Agents and Targets in Atherosclerosis

Lesca M. Holdt¹ Alexander Kohlmaier¹ Daniel Teupser¹

¹Institute of Laboratory Medicine, University Hospital, LMU Munich, Munich, Germany

Thromb Haemost 2019;119:1222–1236.

Address for correspondence Daniel Teupser, MD, PhD, Institute of Laboratory Medicine, University Hospital, LMU Munich, Marchioninstr. 15, 81377 Munich, Germany (e-mail: daniel.teupser@med.uni-muenchen.de).

Abstract

Long noncoding ribonucleic acids (lncRNAs) have been defined as transcripts which are > 200 ribonucleotides in size and are not translated into protein. Recent work has shown that many lncRNAs do have specific molecular functions and biological effects, and are involved in a growing number of diseases, including atherosclerosis. As a consequence, lncRNAs are also becoming interesting targets for therapeutic intervention. Here, we focus on lncRNAs which are expressed in the arterial wall, and describe potential RNA therapeutic approaches of atherosclerosis by manipulating lncRNAs without affecting genome deoxyribonucleic acid content: Starting out with an overview of all lncRNAs that have so far been implicated in atherosclerosis by *in vivo* studies, we describe methodologies for their activation, inactivation, and RNA sequence manipulation. We continue by addressing how artificial (nonnative) therapeutic lncRNAs may be designed, and which molecular functions these designer lncRNAs may exploit. We conclude with an outlook on approaches for chemical lncRNA modification, RNA mass production, and site-specific therapeutic delivery.

Keywords

- transcription
- splicing
- lncRNAs
- circRNAs
- gene expression

Introduction

It is well known today that a large portion of the human genome (~70%) is transcribed and that the majority of the produced transcripts are noncoding (ENCODE or FANTOM consortia).^{1–4} Among the 200,000 known transcripts, around 28,000 stem from long noncoding ribonucleic acid (lncRNA) loci (GENCODE).⁵ Genome-wide association studies (GWAS) recurrently find disease-linked genetic variation in the non-protein-coding sequence space,⁶ often overlapping gene regulatory elements like enhancers, which are actively transcribed and give rise to noncoding RNAs.^{7,8} Concurrently, transcriptomic analyses in patient cohorts reveal numerous lncRNAs which are differentially expressed in diseased tissues. Recent work has shown that many lncRNAs are functional, leading to the notion that lncRNAs may represent a large class of potential therapeutic agents and targets. In this minireview, we describe lncRNAs linked to atherosclerosis in humans which are expressed in cells of the arterial wall (wall endothelial cells [ECs], vascular smooth muscle cells

[VSMCs], and circulating and resident immune cells). We do not cover the roles of lncRNAs regulating atherosclerosis risk factors, such as lipid metabolism, diabetes, or hypertension, or adaptation to ischemic stress. We describe molecular roles of relevant disease-linked lncRNAs, and techniques to therapeutically manipulate them at the RNA level, referred to as RNA therapeutics, an approach that focuses on controlling RNA form and sequence without affecting the deoxyribonucleic acid (DNA) in our genomes.

lncRNA Classes and their Suggested Therapeutic Potential

lncRNAs come in two major flavors, with tens of thousands of cases in each class: (1) linear lncRNAs from dedicated lncRNA genes with their own promoter and terminator and (2) covalently closed circular lncRNAs. The latter are produced through splicing from existing primary transcripts of any type of gene (► **Fig. 1A**, ► **Table 1**) (see Refs. ⁹ and ¹⁰ for review). Linear and circular lncRNA production and

received
November 15, 2018
accepted after revision
April 23, 2019

DOI <https://doi.org/10.1055/s-0039-1692680>.
ISSN 0340-6245.

© 2019 Georg Thieme Verlag KG
Stuttgart · New York

License terms

Fig. 1 Classes and functions of long noncoding ribonucleic acids (lncRNAs). (A) The positioning of lncRNA genes relative to protein-coding genes is a classification criterion. (1) Large intergenic noncoding RNAs (lincRNAs). (2) Natural antisense transcripts (as-lincRNAs). (3) Promoter upstream antisense transcripts (PROMPTs). (4) Enhancer RNAs (eRNAs). lncRNAs transcripts can be linear or circular. Circular lncRNAs are produced by backsplicing of a downstream exon to an upstream exon and do not carry 5'Cap or 3'polyA tail. (B) Functions of lncRNA with atherosclerotic relevance. Functions can be classified in *cis*- and *trans*-acting roles in transcriptional regulation in the nucleus, and in standalone effects (others) in the cytoplasm (bottom). During transcription control, either the lncRNA transcript is functional, or the progression of the RNA polymerase over the lncRNA gene body is the functional determinant (coinciding with chromatin-decompaction of deoxyribonucleic acid [DNA] elements at the lncRNA locus, or with splicing of nascent lncRNA still attached to its DNA template). Abbreviations: IRES, internal ribosome entry site; ORF, open reading frame; RIG-I, retinoic acid inducible gene I; TF, transcription factor.

Table 1 Dichotomy of linear and circular long noncoding RNAs

Linear lncRNAs	circRNAs (3'-5'-linked exon-containing circular RNAs)
> 200 nucleotides (nts) in length	
Non-protein-coding (lacking open reading frames > 100 codons)	
Tens of thousands of cases	
Many isoforms per case	
Produced with cell type- and context-specificity	
Mostly lower abundance per cell than average mRNAs	
Considered to be mostly functional	
Average length ~1,000 nts	Average length ~550 nts
Nuclear enrichment (not all)	Cytoplasmic enrichment (not all)
Mostly carrying 5'Cap and 3' polyA tails (depending on lncRNA class)	Generically lacking 5'Cap and 3' polyA tails due to covalent 3'-5' backbone linkage after backsplicing
Produced by transcription from unique transcriptional units with their own promoter	Produced by spliceosome from primary transcripts (mRNAs or lncRNAs)

Abbreviation: lncRNA, long noncoding ribonucleic acid.

abundance are regulated and are cell-type specific. Both linear and circular lncRNAs have been shown to be overall functional in one pathway or another. lncRNAs effector mechanisms are diverse and complex (►Fig. 1B), making it nontrivial to decide which one to interfere with for therapy:

- (1) lncRNAs guide, scaffold, and control transcription-regulating protein complexes:

Many lncRNAs are known to affect transcription of RNAP I and II target genes, often by impacting chromatin readers and writers in gene promoter control.¹¹ Examples with relevance for atherosclerosis are *ANRIL*, *H19*, *lincRNA-p21*, *MALAT1*, *MEG3*, *NEAT1*, or *TUG1* (►Table 2). lncRNAs affect transcription *in cis*, or also *in trans*, especially if their steady-state abundance is large and allows diffusion of the lncRNA throughout the nucleoplasm. Mechanistically, lncRNAs can recruit chromatin remodelers to target genes via hybridization to DNA, or control their enzymatic activity, or function as negative decoys. lncRNAs can also regulate the transcription of microRNA and other noncoding RNAs (a process distinct from RNA sponging, see 5).

- (2) Enhancer RNAs/eRNAs:

Active enhancers for gene promoters have recently been found to be transcribed and to give rise to enhancer lncRNAs (eRNAs). These can confer enhancer activity by capturing the promoter-contacting Mediator protein complex.¹² Examples of eRNAs important for atherosclerosis are *HOTTIP*, *LEENE*, and *SMILR* (►Table 2). Depending on enhancer, both, the eRNA and the chromatin-opening during transcription of the eRNA locus can be therapeutically relevant.¹¹

- (3) Antisense lncRNAs/asRNAs:

Some lncRNA genes reside within protein-coding gene units, even overlapping coding exons in antisense. Effects on host genes can be positive and negative. Examples with relevance for atherosclerosis are *ANRIL*, *HOXC-AS1*, *MALAT1*,

and *SENCR* (►Table 2). Globally, antisense transcription dampens transcriptional noise and is not part of signal-dependent expression control.¹³ Therapeutic programming through asRNAs is complex, because it may require engineering the genomic locus.

- (4) lncRNAs in subnuclear bodies:

Some lncRNAs can affect other genes through their architectural role in assembling eu- and heterochromatin subnuclear territories. Examples are *MALAT1* in Polycomb bodies,¹⁴ or *NEAT1* in paraspeckles.¹⁵ These indirectly affect gene expression and pre-messenger RNA (mRNA) processing, respectively, depending on the vicinity of genes to these subnuclear bodies. The broadness of the effect and the complexity of the process make it difficult to achieve specificity in therapy.

- (5) lncRNAs as microRNA sponge:

Although many publications implicate endogenous lncRNAs as microRNA sponges (and as inhibitors of microRNA availability and function, therein), many of these reports are met with criticism because evidence often bases on uncontrolled lncRNA overexpression. Few lncRNAs pass the stoichiometric requirements for sponging, though, as most are endogenously not sufficiently highly expressed compared with the number of corresponding microRNA targets and copy numbers of microRNAs per cell (see Ref. ¹⁶ for overview).

- (6) lncRNAs which bind and regulate proteins:

Mass spectrometric analyses showed that a single lncRNA can bind dozens of different proteins in the nucleus and in the cytoplasm, and thereby affect multiple molecular mechanisms at once¹⁷; This concept is best seen for well-studied lncRNAs like *XIST*, which was found to bind > 80 proteins, indicating that it participated in DNA and histone modification and RNA remodeling machineries.¹⁷ With relevance to atherosclerosis, circular *ANRIL* (*circANRIL*) binds to the rRNA processing PeBoW complex for protein translation

control,¹⁸ whereas linear *ANRIL* interacts with members of the PRC1¹⁹ as well as with the PRC2 Polycomb-repressive complexes^{19–21} during presumptive transcription control of target genes (► **Table 2**). In another case, *TUG1* can serve as competing endogenous RNA, and also promote gene activity by chromatin fiber positioning,¹⁴ and was even suggested to regulate cytoskeletal contractility by enhancing the cytoplasmic activity of Ezh2 toward methylating α -actin^{22,23} (► **Table 2**). Conceptually, a lncRNA may also hierarchically regulate a single transcriptional regulator, thereby influencing, in one step, a range of downstream genes and cellular processes. For example, *NRON* participates in scaffolding and restraining the nuclear factor of activated T cell (NFAT) transcription factor in a latently active form in the cytoplasm in unstimulated resting T cells.^{24,25} Transcription factor control is recurrently ascribed to many lncRNAs (► **Table 2**). Together, there is potential for lncRNAs in tuning protein complex activity as “RNA-drugs,” akin to small-molecule drugs, but lncRNA-dependent activity changes in lncRNA:protein complexes are difficult to study, and so far not understood in any case in mechanistic and structural detail.

(7) Protein translation from noncoding RNA:

Despite being nonprotein-coding by definition, some lncRNAs do contain small open reading frames (sORFs), some of which can be translated. Atherosclerosis-specific functions of sORFs are so far unknown. Still, protein expression is, in principle, possible from, both, linear and circular lncRNAs, if the required translation-initiating signals are artificially incorporated into synthetic constructs²⁶ (see chapter on “Disease Therapy by Artificial (Nonnative) Designer lncRNAs” below).

(8) Bifunctional noncoding mRNAs:

The clear separation between coding and noncoding RNA is blurring and some mRNAs carry functions also as untranslated RNAs. For example, the *steroid receptor RNA*, which is in principle protein-coding, also functions as lncRNA in chromatin regulation at specific target genes in the nucleus.^{27,28} And p53, as noncoding RNA, binds and affects the activity of the MDM2 enzyme through structural RNA motifs.²⁹ In another more indirect case, under stress cohorts of several hundred mRNAs become transcribed from alternative upstream transcription start sites, a process that blocks their transcription from the actual promoters. This shift leads to a novel longer RNA isoforms that include a short upstream ORFs with a new stop codon, whose translation blocks the translation of the actual functional protein encoded in a given locus.³⁰ Thereby, functionality lies less in an active role as *cis/trans*-acting noncoding RNA, but more in indirect effects of translation control on the proteome.^{31,32} These findings expand the operational space for noncoding RNA therapeutics (► **Table 1**).

Known Roles of lncRNAs in Atherosclerosis

Limiting our review to those lncRNAs expressed in the vascular wall and functioning in atherosclerosis, 31 lncRNAs

have so far been implicated in vascular cell types (► **Table 2**). Most were initially found because of being differentially expressed in patient cohorts. Only a few (*ANRIL*, *circANRIL*, *MIAT*, *H19*, *LINC00305*), were identified by unbiased GWAS. Fifteen of the listed lncRNAs were studied in immune cell types (such as peripheral blood monocytes, circulating and vascular wall macrophages, or foam cells), 17 in ECs, and 13 in VSMCs.

In the following, we highlight lncRNAs from ► **Table 2** where *in vivo* evidence for therapeutic potential exists. Four groups may be distinguished: (1) lncRNAs with a documented therapeutic benefit for atherosclerosis, (2) lncRNAs with a benefit for other vascular diseases, (3) lncRNAs essential for normal vascular biology, and (4) lncRNAs generally involved in inflammatory signaling (with expected relevance for atherosclerosis).

1. *In vivo* evidence for therapeutic a potential in atherosclerosis has been determined only for one lncRNA: *Neat1*, a well-known lncRNA,⁷⁶ is upregulated in plaques, and knocking-out *Neat1* in mice decreased neointimal lesions in an atherosclerosis model.⁶⁴ Since a full-body mouse knockout was analyzed, it remained unclear in which cell type *Neat1* functioned.⁶⁴ A function in VSMCs was tested *in vitro*: During carotid artery injury, VSMCs usually dedifferentiate from a quiescent to a proliferative/synthetic phenotype, and *Neat1* promoted this atherogenic switch by repressing the function of the chromatin activator WDR5/MLL on serum response factor (SRF) target genes.⁶⁴ Consequently, therapeutically reducing *NEAT1* in VSMCs in lesions might be useful for antagonizing the proatherogenic myocardin-SRF-dependent phenotypic switching of VSMCs,^{64,77} or the proatherogenic oxidized low-density lipoprotein (ox-LDL)-dependent inflammatory signaling in macrophages.⁷⁸ However, given that VSMC proliferation and matrix synthesis are in other contexts also considered beneficial (for example, for plaque repair or for fibrous cap stability) (see Ref.⁷⁹ for review), more work is needed before *NEAT1* can be considered for cell-type-specific therapy.
2. Three lncRNAs showed therapeutic potential at least regarding other vascular diseases: Downregulation of *H19* ameliorates aneurysms,³⁸ and downregulation of *Miat*,⁶² *circHipk3*,³³ or *cZNF609*³⁴ ameliorates diabetic retinopathy.
3. Two lncRNAs, *MeXis*⁶¹ and *Malat1*,^{57,80,81} have been studied by knockouts in mice. Nevertheless, the therapeutic potential of these lncRNAs remains untested: In the first case, *MeXis* levels were found to increase by ox-LDL stimulation of macrophages, upon which this lncRNA induced the *Abca1* transporter and cholesterol efflux.⁶¹ Since knockout of *MeXis* led to increased plaque growth in bone marrow reconstitution experiments of *ldlr*^{-/-} mice, therapeutically increasing *MeXis* (human *TCONS00016111*) expression, especially in patients with single-nucleotide polymorphisms (SNPs) in this gene,⁶¹ might potentially be therapeutically relevant. Care is advised, however, when interpreting data for *Malat1*, which has opposing roles in different cardiovascular conditions: On the one hand, *MALAT1* was found to be downregulated in the plaque,⁵⁵ and knocking-

Table 2 Mammalian lncRNAs implicated in atherogenesis by affecting cells of the vasculature

lncRNA	Regulation in atherosclerotic vascular wall	Immune cells	Endothelial cells	Smooth muscle cells	Molecular function	Effector mechanism	References
<i>ANRIL</i>	Up	x	x	x	Transcription Guiding chromatin regulators (cis/trans-regulation) Enhancer elements in locus	<ul style="list-style-type: none"> • PBMCs, VSMCs proproliferative, antiapoptotic • PBMCs, ECs proadhesive, proinflammatory • ECs proinflammatory, proapoptotic 	19,21
<i>circANRIL</i>	Down	x		x	Protein regulation: Inhibiting rRNA processing PeBoW complex	VSMCs, MΦ antiproliferative, proapoptotic	18
<i>circHIPK3</i>	n.d.		x		ceRNA/microRNA sponge	EC proproliferative, promigrative, antiapoptotic	33
<i>cZNF609</i>	Down		x		ceRNA/microRNA sponge	ECs antiproliferative, proapoptotic, antimigrative	34
<i>GAS5</i>	Up	x	x	x	Parent molecule for small RNA SnoRNAs ceRNA/microRNA sponge Transcription Decoy for TFs	<ul style="list-style-type: none"> • EC, MΦ antiproliferative, proapoptotic • VSMC antidiifferentiative 	35,36
<i>H19</i>	Down (in plaques) Up (in aneurysm)	x	x	x	Transcription Antagonizing microRNAs Tethering chromatin modifiers Parent molecule for microRNAs Binding mRNAs mRNA decay	<ul style="list-style-type: none"> • MΦ proinflammatory • VSMC proapoptotic, invasive • ECs proproliferative, anti-inflammatory, proangiogenic 	37–39
<i>HAS2-AS1</i>	Up			x	Transcription Activating sense transcript	VSMC proproliferative	40,41
<i>HOTAIR</i>	Down		x		Transcription	<ul style="list-style-type: none"> • MΦ proinflammatory, proapoptotic • EC proproliferative, promigrative, antiapoptotic 	42,43
<i>HOTTIP</i>	Up		x		Transcription Tethering chromatin modifiers eRNA-like	EC proproliferative, promigrative	44
<i>HOXC-AS1</i>	Down	x			Transcription Activating sense transcript	MΦ anticholesterol-loading	45
<i>LEENE</i>	n.d.		x		Transcription eRNA	EC anti-inflammatory, antiadhesive	46
<i>Lethe</i>	n.d.	x			Transcription Decoy for TF	MΦ anti-inflammatory	47
<i>LINC00305</i>	Up	x	x		Protein regulation Scaffolding receptors ceRNA/microRNA sponge	<ul style="list-style-type: none"> • PBMC proinflammatory • EC antiproliferative; proapoptotic 	48,49
<i>LincRNA-Cox2</i>	n.d.	x			Transcription Scaffolding chromatin modifier	MΦ immunomodulatory, proinvasive	50–52

Table 2 (Continued)

lncRNA	Regulation in atherosclerotic vascular wall	Immune cells	Endothelial cells	Smooth muscle cells	Molecular function	Effector mechanism	References
<i>lincRNA-Dynlrb2-2</i>	n.d.	x			n.d.	MΦ anticholesterol	53
<i>lincRNA-p21</i>	Down	x	x	x	Transcription Tether for hnRNP, corepressor for p53 Protein regulation Inhibiting E3 ligase Binding to mRNAs Suppression of translation	• VSMC, MΦ antiproliferative, proapoptotic • EC proapoptotic	54
<i>Malat1</i>	Down	x	x	x	Transcription Binding chromatin remodelers 3D chromatin positioning Splicing control Parent molecule for small RNA Cytoplasmic masRNAs	• PBMC, MΦ anti-inflammatory • EC antimigrative, proproliferative • SMC prodifferentiative	55–57
<i>Mantis</i>	n.d.		x		Transcription Scaffold of chromatin modifier	• EC proangiogenic	58
<i>Meg3</i>	Down	x	x	x	Transcription Tether of chromatin modifier Protein regulation Interacting with TF CeRNA/microRNA sponge	• VSMC antiproliferative, antimigrative, proapoptotic • EC antiproliferative	59,60
<i>MeXis</i>	Down	x			Transcription	MΦ cholesterol efflux	61
<i>MIAT/Gomafu</i>	Up		x		Splicing control ceRNA/microRNA sponge	• EC proproliferative, promigrative, proangiogenic	55,62
<i>Myoslid</i>	n.d.			x	n.d.	VSMC antiproliferative, antimigrative	63
<i>Neat1</i>	Up			x	Transcription Decoy for chromatin regulator	VSMC proproliferative, promigrative; antidiifferentiative	64,65
<i>NRON</i>	n.d.	x			Protein regulation Inhibiting TF	T cells immunomodulatory	24
<i>Pacer</i>	n.d.	x			Transcription Inhibiting TF	MΦ anti-inflammatory	66
<i>RNCR3/LINC00599</i>	Up		x	x	ceRNA/microRNA sponge	• VMC proproliferative and promigrative ECs antiapoptotic	67
<i>SENCR</i>	Down		x	x	n.d.	• VSMC proproliferative, promigrative • EC proproliferative promigrative, proangiogenic	68,69
<i>SMILR</i>	Up			x	n.d.	• VSMC proproliferative;	70
<i>STEEL/HOXD-AS1</i>	n.d. (Up)		x		Transcription Binding chromatin modifier	EC proangiogenic, angiogenic patterning	71

(Continued)

Table 2 (Continued)

lncRNA	Regulation in atherosclerotic vascular wall	Immune cells	Endothelial cells	Smooth muscle cells	Molecular function	Effector mechanism	References
<i>THRIL/Linc1992</i>	n.d.	x			Transcription: Binding chromatin factor	MΦ immunomodulatory	72
<i>Tug1</i>	Up	x	x	x	Transcription: 3D chromatin positioning Protein regulation ceRNA/microRNA sponge	<ul style="list-style-type: none"> VSMC, MΦ proliferative, antiapoptotic, proinflammatory VSMCs actin polymerization ECs proapoptotic, antiadhesive 	22,73,74
<i>Uca1</i>	n.d.		x		Transcription	<ul style="list-style-type: none"> EC proliferative, promigrative, antiapoptotic 	75

Abbreviations: 3D, three-dimensional; EC, endothelial cell; lncRNA, long noncoding ribonucleic acid; PBMC, peripheral blood mononuclear cell; TF, transcription factor; VSMC, vascular smooth muscle cell.

out *Malat1* in mice was recently found to trigger immune dysregulation and atherosclerosis in an *apoe*^{-/-} mutant background, remarkably even without the challenge by a coronary artery disease-triggering fat-rich diet.⁸¹ With a similar direction of effect, *Malat1* knockouts developed larger infarcts in brain ischemic mouse models.^{82,83} These two studies suggested that it might be worthwhile to normalize *MALAT1* through overexpression during therapy. Yet, in other contexts, such as aortic thoracic aneurysms or vascular diseases of the Marfan syndrome, beneficial effects seemed to lie rather in *MALAT1* inhibition, and not in its induction.⁵⁷ Also, bluntly overexpressing *Malat1* may have limitations because it is known from other studies that increases in *MALAT1* would promote cancer by effects on cell migration, metastasis, and angiogenesis in hypoxic conditions.^{84,85} Summarizing, different studies showed context- and cell type-dependent therapeutic requirements for *MALAT1*. Consequently, further studies and tools for achieving tissue tropism in delivering therapeutic lncRNA only to specific cells, or for conditional activation/inactivation of lncRNAs in specific conditions and cells, will be required.

- Twelve lncRNAs have been indirectly implicated in atherogenesis through *in vitro* studies in cultured vascular cell types, or through *in vivo* insight in their effectors. A major group in this class comprises 6 lncRNAs involved in modulating inflammatory signaling *in vivo* (*lincRNA-Cox2*, *PACER*, *Lethe*, *THRIL*, *NRON*, *STEEL*) (► **Table 2**). These have considerable therapeutic potential because of the intimate contribution of inflammation to atherogenesis (see following chapter on “Disease Therapy by Manipulating Endogenous lncRNAs”). The rest of lncRNAs in this group control ECs (*Miat*), VSMCs (*Myoslid*), or both (*Gas5*, *lincRNA-p21*, *Meg3*), through functioning in diverse processes.

Translating Molecular Function to Therapeutic Value

In the following, we summarize key therapeutic principles centered on lncRNAs (► **Fig. 2A**). Thereby, we distinguish (1) therapeutic approaches exploiting endogenous lncRNA functionality, and (2) approaches based on artificial (nonnative) designer lncRNAs.

Disease Therapy by Manipulating Endogenous lncRNAs

During the onset of disease, lncRNAs expression levels change in cell- and context-dependent modes. For therapy, lncRNAs that are overactive in disease can be normalized by knockdown approaches. Gain-of-function approaches or RNA sequence correction can be used to antagonize disease-linked changes at the RNA level in other lncRNAs.

- Transcriptional control* is, so far, the major known function of noncoding RNAs. Twenty-one of 31 atherosclerosis-linked lncRNAs function as guides, scaffolds, and regulators of chromatin factors (► **Table 2**). In contrast to

Fig. 2 Concepts and tools for using long noncoding ribonucleic acids (lncRNAs) of the arterial wall in antiatherogenic therapy. (A) Concepts for lncRNA therapy. lncRNA genotype and the ratio of linear and circular lncRNA transcripts must be considered during therapy (left panel). lncRNAs can be therapeutically used to correct the primary molecular causes of disease or to block the pathologically induced cellular effector pathways that confer plaque growth (right panel). Another option would be to, more generally, kill-off disease-causing cells in plaques (right panel). The healthy cell state is denoted in yellow, the diseased state in red. (B) Tools for manipulating lncRNA levels in cells, classified under LOF (loss-of-function) and GOF (gain-of-function) approaches. Target RNAs are labeled in green. Targeting RNAs or ribozymes are marked in red, targeting deoxyribonucleic acid (DNA)-like antisense oligonucleotides (ASOs) in blue. Scissors denote target RNA cleavage by nucleases. Abbreviations: CRISPRa, CRISPR activation for promoter activation; dCas9^{VP16}, inactive Cas9 fused to the generic herpes simplex viral protein 16 (VP16) activation domain; dCas9^{Tet1}, inactive Cas9 fusion with catalytic domain of Tet1 (DNA 5-methylcytosine demethylase); dCas13^{ADAR2}, catalytically inactive Cas13 RNA-guided RNase fused to the ADAR2 (adenosine deaminases acting on RNA 2); gRNA, CRISPR guide RNA; pA – poly (A) tail; REPAIR, RNA Editing for Programmable A to I Replacement; RNAi, RNA interference.

small-molecule drugs which inhibit broadly-acting chromatin factors (such as DNA/histone methyltransferases, demethylases, acetylases, and chromatin readers),⁸⁶ epigenetic therapy through manipulating lncRNA levels and sequence may have the advantage that it allows, in principle, to sequence-specifically control specific subsets of target genes. Separately, it becomes a therapeutic option to guide generic repressor proteins (KRAB- and Chromo-domains) or activators (VP16) to promoters of disease-linked genes via noncoding RNAs.^{87,88}

(2) *Controlling splicing* is equally important for therapy as transcription control. Splice-regulating antisense oligonucleotides (ASOs) can be used to tune splicing in mRNAs and lncRNAs. Additionally, a new therapeutic front was opened when it was found that cotranscriptional circRNA biogenesis interfered with linear RNA biogenesis.^{89–91} A well-studied case is *ANRIL*, transcribed from Chr9p21 in humans, the most prominent atherosclerosis risk loci known so far.^{18,92} Linear *ANRIL* levels positively correlate

with atherosclerosis severity, while *circANRIL* levels anti-correlate with atherosclerosis.^{18,21} While linear *ANRIL* expression causes transitions reminiscent of atherogenesis (inflammation, overproliferation, adhesion in macrophages and VSMCs),^{19–21,93–95} *circANRIL* mediates opposite effects by blocking proliferation and enhancing cell death rates.¹⁸ Thus, one way to decrease atherosclerosis risk would be to coincidentally decrease linear *ANRIL* and increase *circANRIL*, because linear splice forms are known to be increased and circular isoforms decreased in atherosclerosis. How to therapeutically re-engineer the wild-type splicing pattern in a locus is highly complex, as learned from diseases like muscular dystrophies that arise from splicing defects (see Ref.⁹⁶ for a recent review), and it has not yet been practically achieved to simultaneously correct levels of linear and circular *ANRIL*. Theoretically, and confined to RNA-centric approaches, it would be possible to deliver *in vitro* produced synthetic *circANRIL* together with ASOs that target exonic *ANRIL* sequences

that are not contained in the circular isoforms. Other options would be to use splice-changing ASOs, that inhibit forward splicing, and promote intron-backfolding for assisting backsplicing and circularization. Finally, deeper insight into how specific atherosclerosis-linked SNPs in *ANRIL* affect splice site choice⁹² (e.g., through RNA-binding proteins), may allow to get a grip on deliberately affecting circularization.

(3) *Proinflammatory signaling* drives atherogenesis and various lncRNAs regulate inflammation-dependent gene expression in different cell types. For example, during nuclear factor kappa B signaling in macrophages, *LincRNA-Cox2* represses numerous genes by scaffolding RelA-p50 into SWI/SNF complexes.⁵¹ In contrast, *PACER* upregulates *COX-2*, a proatherogenic cyclooxygenase, by interfering with the formation of repressive p50:p50 homodimers,⁶⁶ and *Lethe* inhibits RelA binding to DNA.⁴⁷ Other lncRNAs are active in T cells or ECs: *NRON* scaffolds the cytoplasmic IQGAP1/NFAT1 protein complex, thereby restraining calcineurin from activating the proinflammatory NFAT1 transcription factor in T cells.^{24,97} In ECs, *STEEL* recruits the activating poly-adenosine diphosphate ribosylase PARP1 to *KLF2* and *eNOS*.⁷¹ Depleting *PARP1* is known to limit atherosclerotic plaque growth,⁹⁸ suggesting that inhibiting *STEEL* could be interesting. Together, upregulating *Lethe* and downregulating *lincRNA-Cox2*, *PACER*, and *THRIL* may be therapeutically useful (see Refs.⁹⁹ and ¹⁰⁰ for review on atherogenic inflammation).

Disease Therapy by Artificial (Nonnative) Designer lncRNAs

To achieve lncRNAs with novel functionality, endogenous RNA sequence can be altered, or synthetic constructs are overexpressed. These can contain sequence combinations not existing in endogenous RNA, or stem from *in vitro* evolution routines:

- (1) Although *microRNA sponging* is not considered a common endogenous function for lncRNAs, therapeutically overexpressed lncRNAs are more abundant than endogenous RNAs and may very well become sponges. Such artificial sponges can be optimized by increasing microRNA-binding sites, and by optimally spacing them.¹⁰¹ Furthermore, if target-matched sites are shortened from 8 to 6 nt, microRNA degradation happens instead of sponging, opening therapeutic possibilities even further.¹⁶ circRNAs may become the superior type of sponges because circRNAs are more stable against cellular exoribonucleases than linear RNAs.^{9,10}
- (2) *Artificial RNA aptamers*, as high-affinity binders of biomolecules, constitute a large therapeutic class. Endogenous lncRNAs fold into secondary structures, and more so than mRNAs.¹⁰² Notable are conserved stem-loop structures in protein-interaction interfaces. With a novel *in vitro* sequence evolution methods in development (e.g., SELEX with Pol θ CS13 ribonucleotidyl transferases that deliver random RNA libraries and tolerate 2'-functionalized ribonucleotides¹⁰³), it becomes tangible to engineer

protein- or metabolite-binding RNAs, much like small molecules are classically used as protein-targeting "drugs." An alternative application of RNA aptamers is to bind surface receptors in diseased cells and confer cell entry of drugs and effectors fused to RNA.¹⁰⁴

(3) *Therapeutically relevant peptides* can be expressed from designer lncRNA: Although the vast majority of linear and circular lncRNAs are endogenously not translated to polypeptides by ribosomes, there are exceptions: small ORFs encoding micropeptides are known in lncRNAs, and some have cardiovascular relevance (*LINC0094* 8→myoregulin; *SMIM6*→endoregulin; *LOC100507537*→DWORF).¹⁰⁵ Also, a tiny fraction of native circRNAs can be translated if non-coding RNA segments in the circle fold into internal ribosome entry sites (IRES) to drive translation initiation (*circZNF609*).²⁶ Therefore, one future therapeutic option would be to circularize mRNA to obtain stable expression of therapeutic proteins from a designer circRNA containing an artificial IRES sequence.

(4) *Immunotherapy of atherosclerosis by boosting innate immune signaling*: One future option for atherosclerosis therapy is to boost specific branches in innate-adaptive immune system cross-talk, a concept stemming from research on antitumor strategies (see Ref.¹⁰⁶ for overview). This may potentially supplement the more classical anti-inflammatory strategies to fight atherosclerosis.¹⁰⁷ Recent insight shows that also noncoding RNA may have a place in immunotherapy: First, externally provided synthetic non-coding RNA or RNA analogs are already used as adjuvants to increase the immunogenicity of peptide-based vaccination, by virtue of their ability to stimulate cytoplasmic innate immunity receptors RIG-I and MDA5 as "nonself."^{108–110} Vaccinations, such as with the tolerizing apoB100 epitope,¹¹¹ might in the future benefit. Controlling RIG-I signaling, as far as known from the cancer field, may well also be therapeutically useful for atherosclerosis therapy. The aim here may be to induce programmed cell death by natural killer cells, to enhance phagocytosis by dendritic cell subtypes, to leverage the contribution of specific T cell subtypes in resolving lesions, and to promote neoantigen presentation to lymphocytes.¹¹² A broad MDA-5 and RIG-I activation by RNA is, however, certainly not the goal, because this is known to promote proinflammatory signaling in ECs and macrophages¹¹³ or osteogenic calcification in aortic VSMCs.¹¹⁴ As of yet, too little is known about the cross-talk between innate¹¹⁵ and adaptive immunity¹¹⁶ in atherosclerosis. Therefore, it is open whether it would be beneficial to transfect synthetic noncoding (uncapped or circular) RNAs as therapeutic triggers of RIG-I^{117,118} into specific protective immune cell types, or whether, oppositely, it is the reduction of endogenous RIG-I signaling that may bear protective effects.

Molecular Techniques for lncRNA-Based Therapy

In the following, we briefly describe different modern technologies to knockdown, overexpress, and study aberrantly

expressed or spliced lncRNAs or to change their sequence content for the purpose of therapy (►Fig. 2). On a general note, drug development in atherosclerosis often starts out with disease modeling in animals because not all relevant disease-initiating cell–cell interactions can be recapitulated in cell culture models.

Therapeutic Modulation of lncRNA Levels In Vivo

In the first place, identifying lncRNAs with potential therapeutic relevance necessitates finding animal orthologs of human disease-linked lncRNAs. In atherosclerosis, this can be straightforward (*Malat1*, *Miat*, *Rncr3*, *H19*), or more complicated (*ANRIL*¹¹⁹), because evolutionary selective constraint for the majority of lncRNA sequences is modest at most, and exon–intron structure changes accordingly faster than for protein-coding genes.¹²⁰ If the genomic structure of an atherosclerosis-relevant locus is overall conserved but yields non-coding RNA with only limited conservation, humanizing synthetic disease-linked regions in mice with relevant human lncRNAs through knock-ins may be a viable approach.

If mouse lncRNA orthologs exist, knockdown techniques can be directly executed, involving lncRNA depletion via the RNAi machinery (mostly in the cytoplasm), RNase-H1-type enzymes (also in the cell nucleus), or RNA-cutting ribozymes (►Fig. 2B). Conversely, lncRNA overexpression occurs from plasmids or viral vectors. For circRNA biogenesis from DNA vectors, in many studies, reverse complementary intronic repeats are routinely placed adjacent to circularizing exons, which support backsplicing through backfolding (see Refs. ⁹ and ¹⁰ for review). An interesting new concept are self-amplifying RNA “replicons,” derived from disarmed, cytoplasmic, self-replicating RNA-alphaviruses, which can make RNA therapy permanent without the need for genome integration of RNA-generating vectors.¹²¹ Another novel approach employs guide RNAs to target heterologous activator or repressor domains to lncRNA promoters, which causes up-/downregulation of transcription efficiency.¹²²

Apart from expression via vectors, lncRNAs can also be locally provided to the target tissue, which is the arterial lesion-containing vessel wall, by transfection of *in vitro*-produced RNA molecules: One problem of this approach is that routine solid-phase chemical RNA synthesis is still size-limited (currently 100 nts). Thus, lncRNAs, due to their size, are often transcribed by T7 RNA polymerase *in vitro*. Also, RNA circularization is possible *in vitro*. Thereby, linear T7 transcripts are circularized, either chemically through artificial linkers (e.g., via phosphotriester or click chemistry) or enzymatically through 2'-5' or 3'-5' backbone linkage (via T4 DNA/RNA ligases, tRNA ligases, or ribozymes such as group II intron derivatives) (see Ref. ¹²³ for review). Recombinant expression in heterologous hosts (*Escherichia coli*, yeast) would allow higher linear or circular lncRNAs yield than achieved by *in vitro* transcription, but so far suffered from stability issues and heterogeneity of RNA ends. Novel unconventional bacterial hosts, such as the marine *Rhodovulum sulfidophilum*, circumvent some problems, as they secrete nucleic acids but do not contain RNases in their extracellular space.¹²⁴ Together, novel expression hosts, RNA affinity tags,

and methods like exponential *in vitro* synthesis of RNA through polymerase chain transcription¹²⁵ allow producing sufficient amounts of high-quality RNA for therapy (see Ref. ¹²⁶ for an overview).

Designing lncRNA Function by Modulating RNA Sequence

In addition to modifying lncRNA levels, a second therapeutic approach is to create designer lncRNAs, whose RNA sequence, structural motifs, or posttranscriptional modifications are purposely engineered. Enabling sequence modifications *in vivo*, the technical evolution of Cas enzymes has recently made a significant step forward. Instead of modifying DNA via the classical Cas9 enzymes, Cas13 nucleases were found to target RNA. In one application, Cas13 derivatives allow to purposefully destruct RNAs.^{127,128} But Cas13 derivatives also serve to modify RNA sequence when fusing Cas13 to the ADAR2 enzyme: The latter confers adenosine deamination to inosine in a target RNA, with inosine being functionally equivalent to guanosine in translation and splicing (termed REPAIR tool in ►Fig. 2B).¹²⁸ Linking other RNA-modifying enzymes to Cas13 has potential to modify target RNAs in different ways.

A range of artificial chemical RNA modifications (both at bases and in the phosphodiester backbone) have been chemically introduced in synthetic therapeutic nucleic acids, and benefits for therapy have been determined, mostly from experience with ASOs. Some modifications improve resistance against nucleases, increase potency, or improve pharmacokinetic properties and cellular uptake (see Ref. ¹²⁹ for review). Covalent modifications, as used in ASOs, can theoretically be applied equally to *in vitro* synthesized lncRNAs, such as links to the ribose 2' position (2'-fluoro, 2'-O-methoxyethyl, or cEt-constrained 2'-O-Ethyl). But to date, chemically modified lncRNAs have not yet been used for therapeutic purposes, in part because any modifications may negatively affect interactions with proteins or client RNAs. On the other hand, insight into the roles of some endogenously occurring posttranscriptional chemical modifications of RNAs (both coding and noncoding) are emerging, paving the field of “epitranscriptomics,” as allusion to the so important concept of epigenomic control. For example, methylation (m⁵C, m⁶A), pseudouridylation (Ψ), and editing (deamination of A-to-I), known since more than 50 years, have more recently been functionally related to stability (also) of noncoding RNA,¹³⁰ to the formation of higher-order structure necessary for contact with proteins,^{131,132} to splicing,^{133,134} to RNA backbone rigidity¹³⁵ and base-pairing features,¹³⁶ and functional recognition of microRNA binding sites.^{137,138} But this knowledge has not yet been exploited for engineering lncRNA therapy.

Structure–Function Studies in Designer lncRNAs

In comparison to modulating RNA sequence and covalent modifications on RNA, designing RNA function through engineering secondary and tertiary RNA folds is even more

complex. Although several native RNAs have evolved to regulate proteins, engineering a protein-activity-regulating lncRNA by bioinformatically designing RNA structure *de novo* is not yet possible. However, exploring native RNA: protein complexes with new structural methods (e.g. cryoEM), and through chemical probing, now even possibly inside living cells (PARIS, SHAPE-MaP),¹³⁹ will help to control how lncRNAs bind, scaffold, and regulate their molecular targets.

Hand-in-hand goes the development of novel technologies for profiling the molecular detail of lncRNA interactions with chromatin factors in DNA complexes, a major role of lncRNAs. Techniques for this structure–function analysis at a higher level include ChIRP, CHART, RAP, and ChOP (see Ref.¹⁴⁰ for an overview). Insight from these technologies is central because knowing the chromosomal lncRNA targets in disease-relevant cell types is necessary to achieve specificity in lncRNA therapy.^{127,128}

Potential Side Effects during Systemic lncRNA Therapy and Therapeutic Implications

If thoroughly studied, virtually every lncRNA becomes known to engage multiple effector mechanisms in cell type-specific manners. Consequently, side effects are likely in systemic and long-lasting therapy, as applying to atherosclerosis. For example, *MALAT1* levels drop in plaques,⁵⁵ and *Malat1* knockouts developed atherosclerosis,⁸¹ suggesting that therapeutic upregulation of *MALAT1* might ameliorate disease. On the other hand, *MALAT1* is upregulated in cancer cells and is cancer-promoting.⁸⁴ Such dichotomy necessitates tools for conditional lncRNA delivery. Similar disastrous side effects and dangers regarding cancer development and metabolic syndrome apply to *circHIPK3* and *Gas5*.

To avoid these side effects, and also to counteract off-targeting, conditional delivery schemes are necessary: For guiding their expression in atherosclerotic lesions within the vascular wall, synthetic lncRNAs can be conjugated to plaque-homing peptides (e.g., Ac2–26/LyP-1),¹⁴¹ antibodies or lipid/lipoprotein carriers (high-density lipoprotein),¹⁴² or be packaged in lipid vesicles with targeting cues on their surface (e.g., CCR receptors). The most modern relevant approach for conditional expression in biomedicine is localized delivery through photo- and optoacoustic approaches.¹⁴³ An alternative is to elute RNA from coated stents or from perivascular hydrogels. Not last, lncRNA activity can, in principle, be controlled by laser light, when lncRNA are synthesized with optogenetically regulatable backbones or caging groups,¹⁴⁴ but optogenetic control of lncRNAs has not yet been performed in therapy *in vivo*, so far.

Conclusion

To date, around 70 clinical trials are known to center on RNA therapeutics, and these exclusively involve small interfering ASOs/siRNAs and therapeutic mRNAs, but not yet lncRNAs.¹⁴⁵ Despite the relatively slow translation of RNA-centered therapy into the clinics,¹⁴⁵ RNA therapeutics is

gaining renewed interest, not last through novel insights into lncRNA biology. Further, the ease by which candidate lncRNA can be screened and optimized in their interaction with disease targets surpasses the work with classical small molecule drugs whose targeting to proteins is complex to predict, control, and modify. Any future therapy with lncRNAs will benefit from insight into RNA mass production, chemical modifications, and cellular delivery schemes developed for ASOs/siRNAs over the last decades. As many of the previously unknown cell subtypes that contribute to plaque growth currently become molecularly characterized by novel methods like mass spectrometric cytometry,^{146–148} and as methods for RNA chromatin profiling at single-cell resolution from limited tissue sources emerge,¹⁴⁹ the vision of a highly specific lncRNA-centered therapy in atherosclerosis is materializing, possibly sooner than expected.

Funding

This work was in part funded by the German Research Foundation (DFG) as part of the Collaborative Research Center CRC1123 “Atherosclerosis - Mechanisms and Networks of Novel Therapeutic Targets” (project B1) and by the Leducq-foundation CADgenomics.

Conflict of Interest

None declared.

References

- Bertone P, Stolc V, Royce TE, et al. Global identification of human transcribed sequences with genome tiling arrays. *Science* 2004; 306(5705):2242–2246
- Carninci P, Kasukawa T, Katayama S, et al; FANTOM Consortium; RIKEN Genome Exploration Research Group and Genome Science Group (Genome Network Project Core Group). The transcriptional landscape of the mammalian genome. *Science* 2005;309(5740):1559–1563
- Birney E, Stamatoyannopoulos JA, Dutta A, et al; ENCODE Project Consortium; NISC Comparative Sequencing Program; Baylor College of Medicine Human Genome Sequencing Center; Washington University Genome Sequencing Center; Broad Institute; Children's Hospital Oakland Research Institute. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 2007;447(7146):799–816
- Djebali S, Davis CA, Merkel A, et al. Landscape of transcription in human cells. *Nature* 2012;489(7414):101–108
- Hon CC, Ramilowski JA, Harshbarger J, et al. An atlas of human long non-coding RNAs with accurate 5' ends. *Nature* 2017;543(7644):199–204
- Maurano MT, Humbert R, Rynes E, et al. Systematic localization of common disease-associated variation in regulatory DNA. *Science* 2012;337(6099):1190–1195
- Andersson R, Gebhard C, Miguel-Escalada I, et al. An atlas of active enhancers across human cell types and tissues. *Nature* 2014;507(7493):455–461
- Ørom UA, Derrien T, Beringer M, et al. Long noncoding RNAs with enhancer-like function in human cells. *Cell* 2010;143(01):46–58
- Barrett SP, Salzman J. Circular RNAs: analysis, expression and potential functions. *Development* 2016;143(11):1838–1847
- Holdt LM, Kohlmaier A, Teupser D. Molecular roles and function of circular RNAs in eukaryotic cells. *Cell Mol Life Sci* 2018;75(06): 1071–1098

- 11 Engreitz JM, Haines JE, Perez EM, et al. Local regulation of gene expression by lncRNA promoters, transcription and splicing. *Nature* 2016;539(7629):452–455
- 12 Lai F, Orom UA, Cesaroni M, et al. Activating RNAs associate with Mediator to enhance chromatin architecture and transcription. *Nature* 2013;494(7438):497–501
- 13 Huber F, Bunina D, Gupta I, et al. Protein abundance control by non-coding antisense transcription. *Cell Reports* 2016;15(12):2625–2636
- 14 Yang L, Lin C, Liu W, et al. ncRNA- and Pc2 methylation-dependent gene relocation between nuclear structures mediates gene activation programs. *Cell* 2011;147(04):773–788
- 15 Yamazaki T, Souquere S, Chujo T, et al. Functional domains of NEAT1 architectural lncRNA induce paraspeckle assembly through phase separation. *Mol Cell* 2018;70(06):1038–1053
- 16 Denzler R, McGeary SE, Title AC, Agarwal V, Bartel DP, Stoffel M. Impact of microRNA levels, target-site complementarity, and cooperativity on competing endogenous RNA-regulated gene expression. *Mol Cell* 2016;64(03):565–579
- 17 Chu C, Zhang QC, da Rocha ST, et al. Systematic discovery of Xist RNA binding proteins. *Cell* 2015;161(02):404–416
- 18 Holdt LM, Stahringer A, Sass K, et al. Circular non-coding RNA ANRIL modulates ribosomal RNA maturation and atherosclerosis in humans. *Nat Commun* 2016;7:12429
- 19 Yap KL, Li S, Muñoz-Cabello AM, et al. Molecular interplay of the noncoding RNA ANRIL and methylated histone H3 lysine 27 by polycomb CBX7 in transcriptional silencing of INK4a. *Mol Cell* 2010;38(05):662–674
- 20 Holdt LM, Hoffmann S, Sass K, et al. Alu elements in ANRIL non-coding RNA at chromosome 9p21 modulate atherogenic cell functions through trans-regulation of gene networks. *PLoS Genet* 2013;9(07):e1003588
- 21 Zhou X, Han X, Wittfeldt A, et al. Long non-coding RNA ANRIL regulates inflammatory responses as a novel component of NF- κ B pathway. *RNA Biol* 2016;13(01):98–108
- 22 Chen R, Kong P, Zhang F, et al. EZH2-mediated α -actin methylation needs lncRNA TUG1, and promotes the cortex cytoskeleton formation in VSMCs. *Gene* 2017;616:52–57
- 23 Su IH, Dobenecker MW, Dickinson E, et al. Polycomb group protein ezh2 controls actin polymerization and cell signaling. *Cell* 2005;121(03):425–436
- 24 Sharma S, Findlay GM, Bandukwala HS, et al. Dephosphorylation of the nuclear factor of activated T cells (NFAT) transcription factor is regulated by an RNA-protein scaffold complex. *Proc Natl Acad Sci U S A* 2011;108(28):11381–11386
- 25 Li J, Chen C, Ma X, et al. Long noncoding RNA NRON contributes to HIV-1 latency by specifically inducing tat protein degradation. *Nat Commun* 2016;7:11730
- 26 Wesselhoeft RA, Kowalski PS, Anderson DG. Engineering circular RNA for potent and stable translation in eukaryotic cells. *Nat Commun* 2018;9(01):2629
- 27 Sampath K, Ephrussi A. CncRNAs: RNAs with both coding and non-coding roles in development. *Development* 2016;143(08):1234–1241
- 28 Lanz RB, McKenna NJ, Onate SA, et al. A steroid receptor coactivator, SRA, functions as an RNA and is present in an SRC-1 complex. *Cell* 1999;97(01):17–27
- 29 Candeias MM, Malbert-Colas L, Powell DJ, et al. P53 mRNA controls p53 activity by managing Mdm2 functions. *Nat Cell Biol* 2008;10(09):1098–1105
- 30 Cheng Z, Otto GM, Powers EN, et al. Pervasive, coordinated protein-level changes driven by transcript isoform switching during meiosis. *Cell* 2018;172(05):910–923
- 31 Van Dalen KM, Hodapp S, Keskin A, et al. Global proteome remodeling during ER stress involves Hac1-driven expression of long undecoded transcript isoforms. *Dev Cell* 2018;46(02):219–235
- 32 Chia M, Tresenrider A, Chen J, et al. Transcription of a 5' extended mRNA isoform directs dynamic chromatin changes and interference of a downstream promoter. *eLife* 2017;6:6
- 33 Shan K, Liu C, Liu BH, et al. Circular noncoding RNA HIPK3 mediates retinal vascular dysfunction in diabetes mellitus. *Circulation* 2017;136(17):1629–1642
- 34 Liu C, Yao MD, Li CP, et al. Silencing of circular RNA-ZNF609 ameliorates vascular endothelial dysfunction. *Theranostics* 2017;7(11):2863–2877
- 35 Chen L, Yang W, Guo Y, et al. Exosomal lncRNA GAS5 regulates the apoptosis of macrophages and vascular endothelial cells in atherosclerosis. *PLoS One* 2017;12(09):e0185406
- 36 Tang R, Zhang G, Wang YC, Mei X, Chen SY. The long non-coding RNA GAS5 regulates transforming growth factor β (TGF- β)-induced smooth muscle cell differentiation via RNA Smad-binding elements. *J Biol Chem* 2017;292(34):14270–14278
- 37 Han Y, Ma J, Wang J, Wang L. Silencing of H19 inhibits the adipogenesis and inflammation response in ox-LDL-treated Raw264.7 cells by up-regulating miR-130b. *Mol Immunol* 2018;93:107–114
- 38 Li DY, Busch A, Jin H, et al. H19 induces abdominal aortic aneurysm development and progression. *Circulation* 2018;138(15):1551–1568
- 39 Hofmann P, Sommer J, Theodorou K, et al. Long non-coding RNA H19 regulates endothelial cell aging via inhibition of Stat3 signaling. *Cardiovasc Res* 2019;115(01):230–242
- 40 Vigetti D, Deleonibus S, Moretto P, et al. Natural antisense transcript for hyaluronan synthase 2 (HAS2-AS1) induces transcription of HAS2 via protein O-GlcNAcylation. *J Biol Chem* 2014;289(42):28816–28826
- 41 van den Boom M, Sarbia M, von Wnuck Lipinski K, et al. Differential regulation of hyaluronic acid synthase isoforms in human saphenous vein smooth muscle cells: possible implications for vein graft stenosis. *Circ Res* 2006;98(01):36–44
- 42 Peng Y, Meng K, Jiang L, et al. Thymic stromal lymphopoietin-induced HOTAIR activation promotes endothelial cell proliferation and migration in atherosclerosis. *Biosci Rep* 2017;37(04):BSR20170351
- 43 Liu J, Huang GQ, Ke ZP. Silence of long intergenic noncoding RNA HOTAIR ameliorates oxidative stress and inflammation response in ox-LDL-treated human macrophages by upregulating miR-330-5p. *J Cell Physiol* 2019;234(04):5134–5142
- 44 Liao B, Chen R, Lin F, et al. Long noncoding RNA HOTTIP promotes endothelial cell proliferation and migration via activation of the Wnt/ β -catenin pathway. *J Cell Biochem* 2018;119(03):2797–2805
- 45 Huang C, Hu YW, Zhao JJ, et al. Long noncoding RNA HOXC-AS1 suppresses ox-LDL-induced cholesterol accumulation through promoting HOXC6 expression in THP-1 macrophages. *DNA Cell Biol* 2016;35(11):722–729
- 46 Miao Y, Ajami NE, Huang TS, et al. Enhancer-associated long non-coding RNA LEENE regulates endothelial nitric oxide synthase and endothelial function. *Nat Commun* 2018;9(01):292
- 47 Rapicavoli NA, Qu K, Zhang J, Mikhail M, Laberge RM, Chang HY. A mammalian pseudogene lncRNA at the interface of inflammation and anti-inflammatory therapeutics. *eLife* 2013;2:e00762
- 48 Zhang DD, Wang WT, Xiong J, et al. Long noncoding RNA LINC00305 promotes inflammation by activating the AHRN-NF- κ B pathway in human monocytes. *Sci Rep* 2017;7:46204
- 49 Zhang BY, Jin Z, Zhao Z. Long intergenic noncoding RNA 00305 sponges miR-136 to regulate the hypoxia induced apoptosis of vascular endothelial cells. *Biomed Pharmacother* 2017;94:238–243
- 50 Carpenter S, Aiello D, Atianand MK, et al. A long noncoding RNA mediates both activation and repression of immune response genes. *Science* 2013;341(6147):789–792
- 51 Hu G, Gong AY, Wang Y, et al. LincRNA-Cox2 promotes late inflammatory gene transcription in macrophages through

- modulating SWI/SNF-mediated chromatin remodeling. *J Immunol* 2016;196(06):2799–2808
- 52 Covarrubias S, Robinson EK, Shapleigh B, et al. CRISPR/Cas-based screening of long non-coding RNAs (lncRNAs) in macrophages with an NF- κ B reporter. *J Biol Chem* 2017;292(51):20911–20920
 - 53 Hu YW, Yang JY, Ma X, et al. A lincRNA-DYNLRB2-2/GPR119/GLP-1R/ABCA1-dependent signal transduction pathway is essential for the regulation of cholesterol homeostasis. *J Lipid Res* 2014;55(04):681–697
 - 54 Wu G, Cai J, Han Y, et al. LincRNA-p21 regulates neointima formation, vascular smooth muscle cell proliferation, apoptosis, and atherosclerosis by enhancing p53 activity. *Circulation* 2014;130(17):1452–1465
 - 55 Arslan S, Berkan Ö, Lalem T, et al; Cardiolinc™ network. Long non-coding RNAs in the atherosclerotic plaque. *Atherosclerosis* 2017;266:176–181
 - 56 Gast M, Rauch BH, Nakagawa S, et al. Immune system-mediated atherosclerosis caused by deficiency of long noncoding RNA MALAT1 in ApoE^{−/−} mice. *Cardiovasc Res* 2019;115(02):302–314
 - 57 Lino Cardenas CL, Kessinger CW, Cheng Y, et al. An HDAC9-MALAT1-BRG1 complex mediates smooth muscle dysfunction in thoracic aortic aneurysm. *Nat Commun* 2018;9(01):1009
 - 58 Leisegang MS, Fork C, Josipovic I, et al. Long noncoding RNA MANTIS facilitates endothelial angiogenic function. *Circulation* 2017;136(01):65–79
 - 59 Wu Z, He Y, Li D, et al. Long noncoding RNA MEG3 suppressed endothelial cell proliferation and migration through regulating miR-21. *Am J Transl Res* 2017;9(07):3326–3335
 - 60 Zhou Y, Zhong Y, Wang Y, et al. Activation of p53 by MEG3 non-coding RNA. *J Biol Chem* 2007;282(34):24731–24742
 - 61 Sallam T, Jones M, Thomas BJ, et al. Transcriptional regulation of macrophage cholesterol efflux and atherogenesis by a long noncoding RNA. *Nat Med* 2018;24(03):304–312
 - 62 Yan B, Yao J, Liu JY, et al. lncRNA-MIAT regulates microvascular dysfunction by functioning as a competing endogenous RNA. *Circ Res* 2015;116(07):1143–1156
 - 63 Zhao J, Zhang W, Lin M, et al. MYSLID is a novel serum response factor-dependent long noncoding RNA that amplifies the vascular smooth muscle differentiation program. *Arterioscler Thromb Vasc Biol* 2016;36(10):2088–2099
 - 64 Ahmed ASI, Dong K, Liu J, et al. Long noncoding RNA *NEAT1* (nuclear paraspeckle assembly transcript 1) is critical for phenotypic switching of vascular smooth muscle cells. *Proc Natl Acad Sci U S A* 2018;115(37):E8660–E8667
 - 65 Sun J, Chen G, Jing Y, et al. LncRNA expression profile of human thoracic aortic dissection by high-throughput sequencing. *Cell Physiol Biochem* 2018;46(03):1027–1041
 - 66 Krawczyk M, Emerson BM. p50-associated COX-2 extragenic RNA (PACER) activates COX-2 gene expression by occluding repressive NF- κ B complexes. *eLife* 2014;3:e01776
 - 67 Shan K, Jiang Q, Wang XQ, et al. Role of long non-coding RNA-RNCR3 in atherosclerosis-related vascular dysfunction. *Cell Death Dis* 2016;7(06):e2248
 - 68 Shahmoradi N, Nasiri M, Kamfirooz H, Kheiry MA. Association of the rs55172 polymorphism in SENCER long non-coding RNA and atherosclerotic coronary artery disease. *J Cardiovasc Thorac Res* 2017;9(03):170–174
 - 69 Boulberdaa M, Scott E, Ballantyne M, et al. A role for the long noncoding RNA SENCER in commitment and function of endothelial cells. *Mol Ther* 2016;24(05):978–990
 - 70 Ballantyne MD, Pinel K, Dakin R, et al. Smooth muscle enriched long noncoding RNA (SMILR) regulates cell proliferation. *Circulation* 2016;133(21):2050–2065
 - 71 Man HSJ, Sukumar AN, Lam GC, et al. Angiogenic patterning by STEEL, an endothelial-enriched long noncoding RNA. *Proc Natl Acad Sci U S A* 2018;115(10):2401–2406
 - 72 Li Z, Chao TC, Chang KY, et al. The long noncoding RNA THRIL regulates TNF α expression through its interaction with hnRNPL. *Proc Natl Acad Sci U S A* 2014;111(03):1002–1007
 - 73 Zhang L, Cheng H, Yue Y, Li S, Zhang D, He R. TUG1 knockdown ameliorates atherosclerosis via up-regulating the expression of miR-133a target gene FGF1. *Cardiovasc Pathol* 2018;33:6–15
 - 74 Chen C, Cheng G, Yang X, Li C, Shi R, Zhao N. Tanshinol suppresses endothelial cells apoptosis in mice with atherosclerosis via lncRNA TUG1 up-regulating the expression of miR-26a. *Am J Transl Res* 2016;8(07):2981–2991
 - 75 Yin D, Fu C, Sun D. Silence of lncRNA UCA1 represses the growth and tube formation of human microvascular endothelial cells through miR-195. *Cell Physiol Biochem* 2018;49(04):1499–1511
 - 76 Guo J, Cai H, Zheng J, et al. Long non-coding RNA NEAT1 regulates permeability of the blood-tumor barrier via miR-181d-5p-mediated expression changes in ZO-1, occludin, and claudin-5. *Biochim Biophys Acta Mol Basis Dis* 2017;1863(09):2240–2254
 - 77 Ackers-Johnson M, Talasila A, Sage AP, et al. Myocardin regulates vascular smooth muscle cell inflammatory activation and disease. *Arterioscler Thromb Vasc Biol* 2015;35(04):817–828
 - 78 Chen DD, Hui LL, Zhang XC, Chang Q. NEAT1 contributes to ox-LDL-induced inflammation and oxidative stress in macrophages through inhibiting miR-128. *J Cell Biochem* 2018. Doi: 10.1002/jcb.27541
 - 79 Bennett MR, Sinha S, Owens GK. Vascular smooth muscle cells in atherosclerosis. *Circ Res* 2016;118(04):692–702
 - 80 Yu CK, Xu T, Assoian RK, Rader DJ. Mining the stiffness-sensitive transcriptome in human vascular smooth muscle cells identifies long noncoding RNA stiffness regulators. *Arterioscler Thromb Vasc Biol* 2018;38(01):164–173
 - 81 Gast M, Schroen B, Voigt A, et al. Long noncoding RNA MALAT1-derived mascRNA is involved in cardiovascular innate immunity. *J Mol Cell Biol* 2016;8(02):178–181
 - 82 Zhang X, Tang X, Liu K, Hamblin MH, Yin KJ. Long noncoding RNA Malat1 regulates cerebrovascular pathologies in ischemic stroke. *J Neurosci* 2017;37(07):1797–1806
 - 83 Xin JW, Jiang YG. Long noncoding RNA MALAT1 inhibits apoptosis induced by oxygen-glucose deprivation and reoxygenation in human brain microvascular endothelial cells. *Exp Ther Med* 2017;13(04):1225–1234
 - 84 Arun G, Diermeier S, Akerman M, et al. Differentiation of mammary tumors and reduction in metastasis upon Malat1 lncRNA loss. *Genes Dev* 2016;30(01):34–51
 - 85 Michalik KM, You X, Manavski Y, et al. Long noncoding RNA MALAT1 regulates endothelial cell function and vessel growth. *Circ Res* 2014;114(09):1389–1397
 - 86 Hauser AT, Robaa D, Jung M. Epigenetic small molecule modulators of histone and DNA methylation. *Curr Opin Chem Biol* 2018;45:73–85
 - 87 Gilbert LA, Larson MH, Morsut L, et al. CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* 2013;154(02):442–451
 - 88 Qi LS, Larson MH, Gilbert LA, et al. Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene expression. *Cell* 2013;152(05):1173–1183
 - 89 Ashwal-Fluss R, Meyer M, Pamudurti NR, et al. circRNA biogenesis competes with pre-mRNA splicing. *Mol Cell* 2014;56(01):55–66
 - 90 Zhang Y, Xue W, Li X, et al. The biogenesis of nascent circular RNAs. *Cell Reports* 2016;15(03):611–624
 - 91 Zhang XO, Wang HB, Zhang Y, Lu X, Chen LL, Yang L. Complementary sequence-mediated exon circularization. *Cell* 2014;159(01):134–147
 - 92 Burd CE, Jeck WR, Liu Y, Sanoff HK, Wang Z, Sharpless NE. Expression of linear and novel circular forms of an INK4/ARF-associated non-coding RNA correlates with atherosclerosis risk. *PLoS Genet* 2010;6(12):e1001233

- 93 Almontashiri NA, Antoine D, Zhou X, et al. 9p21.3 coronary artery disease risk variants disrupt TEAD transcription factor-dependent transforming growth factor β regulation of p16 expression in human aortic smooth muscle cells. *Circulation* 2015;132(21):1969–1978
- 94 Miller CL, Pjanic M, Wang T, et al. Integrative functional genomics identifies regulatory mechanisms at coronary artery disease loci. *Nat Commun* 2016;7:12092
- 95 Mumbach MR, Satpathy AT, Boyle EA, et al. Enhancer connectome in primary human cells identifies target genes of disease-associated DNA elements. *Nat Genet* 2017;49(11):1602–1612
- 96 Scotti MM, Swanson MS. RNA mis-splicing in disease. *Nat Rev Genet* 2016;17(01):19–32
- 97 Goettsch C, Rauner M, Hamann C, et al. Nuclear factor of activated T cells mediates oxidised LDL-induced calcification of vascular smooth muscle cells. *Diabetologia* 2011;54(10):2690–2701
- 98 von Lukowicz T, Hassa PO, Lohmann C, et al. PARP1 is required for adhesion molecule expression in atherogenesis. *Cardiovasc Res* 2008;78(01):158–166
- 99 Chen YG, Satpathy AT, Chang HY. Gene regulation in the immune system by long noncoding RNAs. *Nat Immunol* 2017;18(09):962–972
- 100 Tabas I, Lichtman AH. Monocyte-macrophages and T cells in atherosclerosis. *Immunity* 2017;47(04):621–634
- 101 Jost I, Shalamova LA, Gerresheim GK, Niepmann M, Bindereif A, Rossbach O. Functional sequestration of microRNA-122 from Hepatitis C Virus by circular RNA sponges. *RNA Biol* 2018;15(08):1032–1039
- 102 Spitale RC, Flynn RA, Zhang QC, et al. Structural imprints in vivo decode RNA regulatory mechanisms. *Nature* 2015;519(7544):486–490
- 103 Randrianjatovo-Gbalou I, Rosario S, Sismeiro O, et al. Enzymatic synthesis of random sequences of RNA and RNA analogues by DNA polymerase theta mutants for the generation of aptamer libraries. *Nucleic Acids Res* 2018;46(12):6271–6284
- 104 Pastor F, Kolonias D, Giangrande PH, Gilboa E. Induction of tumour immunity by targeted inhibition of nonsense-mediated mRNA decay. *Nature* 2010;465(7295):227–230
- 105 Nelson BR, Makarewich CA, Anderson DM, et al. A peptide encoded by a transcript annotated as long noncoding RNA enhances SERCA activity in muscle. *Science* 2016;351(6270):271–275
- 106 Gajewski TF, Schreiber H, Fu YX. Innate and adaptive immune cells in the tumor microenvironment. *Nat Immunol* 2013;14(10):1014–1022
- 107 Bäck M, Hansson GK. Anti-inflammatory therapies for atherosclerosis. *Nat Rev Cardiol* 2015;12(04):199–211
- 108 Chen YG, Kim MV, Chen X, et al. Sensing self and foreign circular RNAs by Intron identity. *Mol Cell* 2017;67(02):228–238
- 109 Coffman RL, Sher A, Seder RA. Vaccine adjuvants: putting innate immunity to work. *Immunity* 2010;33(04):492–503
- 110 Wang L, Smith D, Bot S, Dellamary L, Bloom A, Bot A. Noncoding RNA danger motifs bridge innate and adaptive immunity and are potent adjuvants for vaccination. *J Clin Invest* 2002;110(08):1175–1184
- 111 Kimura T, Tse K, McArdle S, et al. Atheroprotective vaccination with MHC-II-restricted ApoB peptides induces peritoneal IL-10-producing CD4 T cells. *Am J Physiol Heart Circ Physiol* 2017;312(04):H781–H790
- 112 Elion DL, Cook RS. Harnessing RIG-I and intrinsic immunity in the tumor microenvironment for therapeutic cancer treatment. *Oncotarget* 2018;9(48):29007–29017
- 113 Asdonk T, Steinmetz M, Krogmann A, et al. MDA-5 activation by cytoplasmic double-stranded RNA impairs endothelial function and aggravates atherosclerosis. *J Cell Mol Med* 2016;20(09):1696–1705
- 114 Ramachandran B, Stabley JN, Cheng SL, et al. A GTPase-activating protein-binding protein (G3BP1)/antiviral protein relay conveys arteriosclerotic Wnt signals in aortic smooth muscle cells. *J Biol Chem* 2018;293(21):7942–7968
- 115 Christ A, Bekkering S, Latz E, Riksen NP. Long-term activation of the innate immune system in atherosclerosis. *Semin Immunol* 2016;28(04):384–393
- 116 Wolf D, Ley K. Immunity and inflammation in atherosclerosis. *Circ Res* 2019;124(02):315–327
- 117 Li X, Liu CX, Xue W, et al. Coordinated circRNA biogenesis and function with NF90/NF110 in viral infection. *Mol Cell* 2017;67(02):214–227
- 118 Hochheiser K, Klein M, Gottschalk C, et al. Cutting edge: the RIG-I ligand 3pRNA potently improves CTL cross-priming and facilitates antiviral vaccination. *J Immunol* 2016;196(06):2439–2443
- 119 Visel A, Zhu Y, May D, et al. Targeted deletion of the 9p21 non-coding coronary artery disease risk interval in mice. *Nature* 2010;464(7287):409–412
- 120 Ulitsky I. Evolution to the rescue: using comparative genomics to understand long non-coding RNAs. *Nat Rev Genet* 2016;17(10):601–614
- 121 Kim DY, Atasheva S, McAuley AJ, et al. Enhancement of protein expression by alphavirus replicons by designing self-replicating subgenomic RNAs. *Proc Natl Acad Sci U S A* 2014;111(29):10708–10713
- 122 Bester AC, Lee JD, Chavez A, et al. An integrated genome-wide CRISPRa approach to functionalize lncRNAs in drug resistance. *Cell* 2018;173(03):649–664
- 123 Müller S, Appel B. In vitro circularization of RNA. *RNA Biol* 2017;14(08):1018–1027
- 124 Suzuki H, Ando T, Umekage S, Tanaka T, Kikuchi Y. Extracellular production of an RNA aptamer by ribonuclease-free marine bacteria harboring engineered plasmids: a proposal for industrial RNA drug production. *Appl Environ Microbiol* 2010;76(03):786–793
- 125 Chen T, Romesberg FE. Polymerase chain transcription: exponential synthesis of RNA and modified RNA. *J Am Chem Soc* 2017;139(29):9949–9954
- 126 Baronti L, Karlsson H, Marušič M, Petzold K. A guide to large-scale RNA sample preparation. *Anal Bioanal Chem* 2018;410(14):3239–3252
- 127 Abudayyeh OO, Gootenberg JS, Essletzbichler P, et al. RNA targeting with CRISPR-Cas13. *Nature* 2017;550(7675):280–284
- 128 Cox DBT, Gootenberg JS, Abudayyeh OO, et al. RNA editing with CRISPR-Cas13. *Science* 2017;358(6366):1019–1027
- 129 Khvorova A, Watts JK. The chemical evolution of oligonucleotide therapies of clinical utility. *Nat Biotechnol* 2017;35(03):238–248
- 130 Wang Y, Li Y, Toth JI, Petroski MD, Zhang Z, Zhao JC. N6-methyladenosine modification destabilizes developmental regulators in embryonic stem cells. *Nat Cell Biol* 2014;16(02):191–198
- 131 Liu N, Dai Q, Zheng G, He C, Parisien M, Pan T. N(6)-methyladenosine-dependent RNA structural switches regulate RNA-protein interactions. *Nature* 2015;518(7540):560–564
- 132 Patil DP, Chen CK, Pickering BF, et al. m(6)A RNA methylation promotes XIST-mediated transcriptional repression. *Nature* 2016;537(7620):369–373
- 133 Kasowitz SD, Ma J, Anderson SJ, et al. Nuclear m6A reader YTHDC1 regulates alternative polyadenylation and splicing during mouse oocyte development. *PLoS Genet* 2018;14(05):e1007412
- 134 Rueter SM, Dawson TR, Emeson RB. Regulation of alternative splicing by RNA editing. *Nature* 1999;399(6731):75–80
- 135 Davis DR. Stabilization of RNA stacking by pseudouridine. *Nucleic Acids Res* 1995;23(24):5020–5026
- 136 Karijolich J, Yu YT. Converting nonsense codons into sense codons by targeted pseudouridylation. *Nature* 2011;474(7351):395–398

- 137 Kawahara Y, Zinshteyn B, Sethupathy P, Iizasa H, Hatzigeorgiou AG, Nishikura K. Redirection of silencing targets by adenosine-to-inosine editing of miRNAs. *Science* 2007;315(5815):1137–1140
- 138 Goldstein B, Agranat-Tamir L, Light D, Ben-Naim Zgayer O, Fishman A, Lamm AT. A-to-I RNA editing promotes developmental stage-specific gene and lncRNA expression. *Genome Res* 2017;27(03):462–470
- 139 Smola MJ, Weeks KM. In-cell RNA structure probing with SHAPE-MaP. *Nat Protoc* 2018;13(06):1181–1195
- 140 Kashi K, Henderson L, Bonetti A, Carninci P. Discovery and functional analysis of lncRNAs: methodologies to investigate an uncharacterized transcriptome. *Biochim Biophys Acta* 2016;1859(01):3–15
- 141 Kamaly N, Fredman G, Subramanian M, et al. Development and in vivo efficacy of targeted polymeric inflammation-resolving nanoparticles. *Proc Natl Acad Sci U S A* 2013;110(16):6506–6511
- 142 Duivenvoorden R, Tang J, Cormode DP, et al. A statin-loaded reconstituted high-density lipoprotein nanoparticle inhibits atherosclerotic plaque inflammation. *Nat Commun* 2014;5:3065
- 143 Zhang Y, Yu J, Kahkoska AR, Gu Z. Photoacoustic drug delivery. *Sensors (Basel)* 2017;17(06):E1400
- 144 Lubbe AS, Szymanski W, Feringa BL. Recent developments in reversible photoregulation of oligonucleotide structure and function. *Chem Soc Rev* 2017;46(04):1052–1079
- 145 Kaczmarek JC, Kowalski PS, Anderson DG. Advances in the delivery of RNA therapeutics: from concept to clinical reality. *Genome Med* 2017;9(01):60
- 146 Cole JE, Park I, Ahern DJ, et al. Immune cell census in murine atherosclerosis: cytometry by time of flight illuminates vascular myeloid cell diversity. *Cardiovasc Res* 2018;114(10):1360–1371
- 147 Misra A, Feng Z, Chandran RR, et al. Integrin beta3 regulates clonality and fate of smooth muscle-derived atherosclerotic plaque cells. *Nat Commun* 2018;9(01):2073
- 148 Guimarães-Camboa N, Evans SM. Are perivascular adipocyte progenitors mural cells or adventitial fibroblasts? *Cell Stem Cell* 2017;20(05):587–589
- 149 Cusanovich DA, Hill AJ, Aghamirzaie D, et al. A single-cell atlas of in vivo mammalian chromatin accessibility. *Cell* 2018;174(05):1309–1324