Knowledge of Pregnant Women about Pregnancy Gingivitis and Children Oral Health

Rafi A. Togoo¹ Balsam Al-Almai¹ Fatimah Al-Hamdi¹ Salha H. Huaylah¹ Mashael Althobati¹ Safeyah Alqarni¹

¹Department of Pediatric Dentistry and Orthodontics King Khalid University College of Dentistry Abha, Kingdom of Saudi Arabia

Address for correspondence Salha H. Huaylah, BDS, Department of Pediatric Dentistry and Orthodontics King Khalid University College of Dentistry Abha, Kingdom of Saudi Arabia (e-mail: salhahuaylah@gmail.com).

Eur J Dent 2019;13:261–270

Abstract

Objective A cross-sectional questionnaire-based study was carried out among 251 pregnant women to assess their level of knowledge and perception regarding pregnancy gingivitis and the oral health of new born children.

Materials and Methods A self-administered validated questionnaire was distributed among pregnant women visiting the Child and Maternity Hospital, Ministry of Health, Aseer, Saudi Arabia. The questionnaire included the questions related to their knowledge/awareness about cause, treatment, preventive measures, and the adverse effects of gingivitis on oral health of a woman.

Statistical Analysis The data was entered into Microsoft excel sheet and put to statistical analysis using SPSS (version 17). Descriptive analysis and also chi-squared test was used for comparison.

Results Majority of pregnant women were found unaware about developing of pregnancy gingivitis as well as the cause, effects, treatment, and preventive measures for the same. Most of the subjects agreed that it was essentially required to have oral health education before their pregnancy term would start.

Conclusion Most of the pregnant women said that they were unaware about pregnancy gingivitis, though few reported of being aware about it. It is inferred that the collaborative efforts between general dentists, pediatric dentists, public health dentists, and medical professionals are needed to raise awareness among pregnant women regarding timely dental care to avoid the adverse pregnancy related outcomes due to gingivitis and periodontitis.

Keywords

- pregnancy
- gingivitis
- knowledge
- women

Introduction

Various physiological changes take place through the body of a pregnant woman that include both general and those related to oral cavity.¹ Due to hormonal fluctuations during pregnancy, alterations occur in the levels of estrogen and progesterone resulting in tortuous and dilated microvasculature, more permeability of oral blood vessels, and lowering of host immunity, which eventually cause the host to become more vulnerable to oral infections.^{2,3} Both reversible and irreversible changes take place in oral cavities of women during

pregnancy. High levels of estrogen have been found to be associated with occurrence of gingival hyperplasia, gingivitis, pyogenic granulomas, dental caries, and alterations in salivary flow.⁴ In the studies conducted previously, oral health of pregnant women, adverse pregnancy outcomes (such as preterm births, low birth weight), and oral health of infants have all been linked to each other.⁵

Pregnancy leads not only to the physiological changes in a woman but it also has a menacing impact on her behavioral as well as dietary habits. With the increased need of energy, a pregnant woman has to modify the quantity and content of

diet that ultimately increases the carbohydrate quantum of food, resulting in more cariogenic food. al-Kanhal and Bani, in their study on pregnant women in Saudi Arabia, reported that majority of women desired for more and more sweets, dates, milk, and salty and sour food, hence increasing their risk of dental caries and dental erosion.⁶

Oral health promotion and relevant oral health education, which would acquaint them regarding timely treatment and preventive measures for gaining optimal oral health, are imperative to be imparted to women especially during the child-bearing age. Women, during their pregnancy, face a number of problems in utilizing the available oral health services due to which their oral health gets deteriorated. They lack knowledge about the appropriate timing for getting their dental treatment done during or before pregnancy. Some of them suffer from negative attitude toward seeking dental care that might be related to their past experience and/or misconception about it. At times, even the dentists show reluctance to treat pregnant women that adds to the negligence of pregnant women toward oral health.⁷

Hence, a study was conducted in Aseer, Saudi Arabia among pregnant women to assess their knowledge and perception toward pregnancy gingivitis and oral health of their children.

Materials and Methods

A cross-sectional questionnaire-based study was carried out among 18 to 40 years old pregnant women, who visited the Child and Maternity Hospital, Aseer, Saudi Arabia. Prior to conducting of study, ethical approval was obtained from Scientific Research Committee, Aseer, Saudi Arabia. A total convenient sample of 251 pregnant women was included in the study. All the pregnant women who were permanent residents of Saudi Arabia, belonged to age group of 18 to 40 years and gave the consent of participation in the research, were included in the study, while all those who suffered from any systemic ailments such as diabetes and bleeding disorders were excluded from the study.

A close-ended questionnaire was formulated and its reliability was checked by conducting a pilot study on 20 pregnant women who were not a part of the final study sample. The questions were checked for ease in understanding and having clarity by repeating the study on the same respondents. Their responses were thereby checked for reliability using Kappa statistics that yielded satisfactory results (0.81).

The final questionnaire (**Appendix A**) included two sections: in the first section demographic data such as age and educational qualification of the participants were recorded and in the second section questions related to awareness and knowledge about oral health of women were enquired; as for instance those related to pregnancy gingivitis, its effect on oral health of women, its impact on their new born children, the preventive and treatment measures related to pregnancy gingivitis, and their frequency of visiting the dentist during pregnancy. Moreover, the questionnaire pertained to subject's knowledge regarding the effect of poor oral health on pregnancy outcomes such as preterm births and low birth weights.

An 11-item questionnaire, in both English and Arabic languages, was used and the subjects were interviewed face-to-face so that any difficulty that might creep up in understanding the questions could be explained to the subjects by interviewer. Confidentiality of the responses and identity of the patients were scrupulously maintained and assured so as to avoid nonresponse and social desirability bias.

The data was entered in Microsoft Excel sheet and put to statistical analysis using SPSS (version 17). Descriptive analysis and chi-squared tests were used for comparison.

Results

In the present study, the age of pregnant women, who participated, ranged between 27 and 39 years, with mean age of 31.80 years ($df=3.36$). Majority (42.62%) of the pregnant women had just heard about developing pregnancy gingivitis and they were not fully aware about it and 31.47% had not even heard about it, whereas 25.89% were aware about it. Majority of women (41.03%) did not get their dental checkup done during pregnancy even after many of them (28.68%) had developed reddened and swollen gums. The cause of pregnancy gingivitis was not known to 32.66% of the pregnant women. As many as 35.05% respondents felt that daily tooth brushing and flossing would prevent pregnancy gingivitis (**►Table 1**). Yet as many as 28.68% pregnant women thought that dentists would prefer medications as a treatment of pregnancy gingivitis, while 19.92% did not know about it. Around 27% of subjects considered tooth sensitivity to be the main consequence of pregnancy gingivitis. When enquired whether pregnancy gingivitis could affect the oral health of a new born, a majority of women (47.41%) were undecided about it, while 28.68% were unaware of any such effect on the oral health of the new born. Most of the pregnant women (55.37%) were unaware about the adverse pregnancy outcome, associated with pregnancy gingivitis (**►Table 2**). When the responses of pregnant women were compared, based on the level of education, a statistically significant difference was found between the responses of all the questions ($p<0.05$). Respondents to the tune of 49% reported that they used moist towels for cleaning oral cavity of their children who were less than 18 months of age (**►Fig. 1**). Majority of subjects believed positively to receive dental education compulsorily at the start of their pregnancy term (**►Fig. 2**).

Discussion

Inflammation of soft tissues surrounding the teeth, without loss of attachment, is known as gingivitis, while periodontitis involves damage of supporting tooth structures. The inflammation of gums that is exaggerated during the period of pregnancy is known as pregnancy gingivitis given rise due to changes in hormonal levels and in that of life style of a pregnant woman. In various previous studies, poor oral health such as periodontitis in a pregnant woman has been associated with various adverse pregnancy outcomes such as preterm births and low birth weights.⁸ Increase in the hormonal levels of estrogen and progesterone can cause

Table 1 Frequency distribution of responses according to education level of subjects

Q1) Being aware about developing pregnancy gingivitis during pregnancy	Total no. of respondents (%age)	Illiterate women	Primary school	High school	Higher secondary school	Graduate	Postgraduate
Yes, I am fully aware	65 (25.89%)	1 (1.53%)	2 (3.07%)	3 (4.61%)	6 (9.2%)	8 (12.30%)	45 (69.23%)
No, I'm not aware	79 (31.47%)	22 (27.84%)	9 (11.39%)	20 (25.31%)	12 (15.18%)	14 (17.72%)	2 (2.53%)
Not fully aware, but heard it	107 (42.62%)	22 (20.56%)	17 (15.88%)	26 (24.29%)	21 (19.62%)	20 (18.69%)	1 (0.93%)
Chi-squared test = 149.54, df = 12, p-Value = 0.0000							
Q2) Getting the dental checkup done during or before pregnancy	Total no. of respondents (%age)	Illiterate women	Primary school	High school	Higher secondary school	Graduate	Postgraduate
No, I didn't	103 (41.03%)	35 (33.98%)	18 (17.47%)	21 (20.38%)	21 (20.38%)	6 (5.82%)	8 (7.76%)
Yes, I did	62 (24.8%)	3 (4.83%)	4 (6.45%)	11 (17.74%)	11 (17.74%)	22 (35.48%)	24 (38.70%)
Yes I planned to but did not go.	49 (19.52)	6 (12.24%)	2 (4.08%)	8 (16.32%)	8 (16.32%)	8 (16.32%)	6 (12.24%)
Yes, due to dental condition	37 (14.74%)	1 (2.70%)	4 (10.81%)	9 (24.32%)	9 (24.32%)	6 (16.21%)	10 (27.02%)
Chi-squared test = 73.867, df = 15, p-Value = 0.0000							
Q3) Do you have any following signs of pregnancy gingivitis	Total no. of respondents (%age)	Illiterate women	Primary school	High school	Higher secondary school	Graduate	Postgraduate
Swollen & reddened gums	72 (28.68%)	14 (19.44%)	6 (8.33%)	14 (19.44%)	9 (12.5%)	12 (16.66%)	14 (19.44%)
Bad breath	65 (25.89%)	12 (18.46%)	8 (12.30%)	11 (16.92%)	8 (12.30%)	10 (15.38%)	10 (15.38%)
Gum bleeding	69 (27.49%)	14 (20.28%)	11 (15.94%)	6 (8.69%)	7 (10.14%)	8 (11.59%)	12 (17.39%)
Red spot over oral lining	10 (3.98)	4 (40%)	2 (20%)	5 (50%)	2 (20%)	0 (0%)	0 (0%)
None	35 (13.94%)	1 (2.85%)	1 (2.85%)	3 (8.57%)	13 (37.14%)	12 (34.28%)	12 (34.28%)
Chi-squared test = 38.631, df = 20, p-Value = 0.0071							
Q4) What may be the cause of gingivitis during pregnancy	Total no. of respondents (%age)	Illiterate women	Primary school	High school	Higher secondary school	Graduate	Postgraduate
Hormonal changes	78 (31.07%)	2 (2.56%)	2 (2.56%)	14 (17.94%)	12 (15.38%)	20 (25.64%)	22 (28.20%)
Traumatic tooth brushing	20 (7.96)	9 (45%)	2 (10%)	4 (20%)	2 (10%)	1 (5%)	2 (10%)
Poor oral hygiene	45 (17.92%)	2 (4.44%)	3 (6.66%)	9 (20%)	9 (20%)	14 (31.11%)	11 (24.44%)
Eating fried food	6 (2.39%)	2 (33.33%)	1 (16.66%)	2 (33.33%)	2 (33.33%)	1 (16.66%)	3 (50%)
Malnutrition	20 (7.96%)	8 (40%)	8 (40%)	2 (10%)	1 (5%)	0 (0%)	3 (15%)
Do not know	82 (32.66%)	22 (26.82%)	12 (14.63%)	18 (21.95%)	13 (15.85%)	6 (7.31%)	7 (8.53%)
Chi-squared test = 85.285, df = 25, p-Value = 0.0000							

continued

continued

Q5) Measures that you think can prevent Pregnancy gingivitis	Total no. of respondents (%age)	Illiterate women	Primary school	High schooling	Higher secondary school	Graduate	Postgraduate
Daily tooth brushing and flossing	88 (35.05%)	15 (17.04%)	11 (12.5%)	17 (19.31%)	11 (12.5%)	14 (15.90%)	18 (20.45%)
Balanced diet	55 (21.91%)	12 (21.81%)	3 (5.45%)	11 (20%)	12 (21.81%)	9 (16.36%)	10 (18.18%)
Dental checkup before pregnancy	58 (23.10%)	9 (15.51%)	10 (17.24%)	13 (22.41%)	7 (12.06%)	9 (15.51%)	13 (22.41%)
Monthly Scaling	10 (3.98%)	3 (30%)	0 (0%)	2 (20%)	2 (20%)	0 (0%)	3 (30%)
Dental checkup during second trimester	32 (12.74%)	2 (6.25%)	1 (3.12%)	6 (18.75%)	7 (21.87%)	10 (31.25%)	4 (12.5%)
Do not know	8 (3.18%)	4 (50%)	3 (37.5%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Chi-squared test = 39.0232, df = 25, p-Value = 0.0367							

Table 2 Frequency distribution of responses according to education level of subjects

Q6) What as per your knowledge, is the treatment dentists provide for pregnancy gingivitis	Total no. of respondents	Illiterate	Primary school	High school	Higher secondary school	Graduate	Postgraduate
Surgical removal of swollen gum	10 (3.98%)	5 (50%)	1 (10%)	2 (20%)	1 (10%)	1 (10%)	2 (20%)
No treatment needed	35 (13.94%)	8 (22.85%)	5 (14.28%)	9 (25.71%)	4 (11.42%)	1 (2.85%)	2 (5.71%)
Professional scaling	60 (23.90%)	1 (1.66%)	1 (1.66%)	5 (8.33%)	15 (25%)	24 (40%)	28 (46.66%)
Extract the affected tooth	24 (9.56%)	7 (29.16%)	7 (29.16%)	2 (8.33%)	1 (4.16%)	1 (4.16%)	0 (0%)
Medications	72 (28.68%)	8 (11.11%)	5 (6.94%)	17 (23.61%)	15 (20.83%)	13 (18.05%)	15 (20.83%)
Do not know	50 (19.92%)	16 (32%)	9 (18%)	14 (28%)	3 (6%)	2 (4%)	1 (2%)
Chi-squared test = 119.645, df = 25, p-Value = 0.0000							
Q7) What affect can pregnancy gingivitis have on your oral health?	Total no. of respondents	Illiterate	Primary school	High school	Higher secondary school	Graduate	Postgraduate
Overgrowth of gum tissue	27 (10.75%)	2 (7.40%)	1 (3.70%)	3 (11.11%)	5 (18.51%)	6 (22.22%)	7 (25.92%)
No effect	22 (8.76%)	13 (59.09%)	4 (18.18%)	3 (13.63%)	0	1 (4.54%)	1 (4.54%)
Periodontal disease	34 (13.54%)	2 (5.88%)	3 (8.82%)	8 (23.52%)	6 (17.64%)	7 (20.58%)	8 (23.52%)
Tooth decay	40 (15.93%)	11 (27.5%)	7 (17.5%)	12 (30%)	7 (17.5%)	2 (5%)	1 (2.5%)
Tooth sensitivity	67 (26.69%)	4 (5.97%)	3 (4.47%)	11 (16.41%)	13 (19.40%)	14 (20.89%)	21 (31.34%)
Loose tooth	30 (11.95%)	1 (3.33%)	1 (3.33%)	4 (13.33%)	5 (16.66%)	11 (36.66%)	9 (30%)
Do not know	31 (12.35%)	12 (38.70%)	9 (29.03%)	8 (25.80%)	3 (9.67%)	1 (3.22%)	1 (3.22%)
Chi-squared test = 106.470, df = 30, p-Value = 0.000							

continued

continued

Q8). Will pregnancy gingivitis affect the oral health of new born?	Total no. of respondents	Illiterate	Primary school	High school	Higher secondary school	Graduate	Postgraduate
Yes, it will	60 (23.90%)	3 (5%)	2 (3.33%)	12 (20%)	11 (18.33%)	15 (25%)	24 (40%)
No, it won't	72 (28.68%)	18 (25%)	12 (16.66%)	16 (22.22%)	4 (5.55%)	8 (11.11%)	8 (11.11%)
Undecided	119 (47.41%)	24 (20.16%)	14 (11.76%)	21 (17.64%)	24 (20.16%)	19 (15.96%)	16 (13.44%)
Chi-squared test = 39.4812, df = 10, p-Value = 0.00002							
Q9) What can be the adverse pregnancy outcome of pregnancy gingivitis?	Total no. of respondents	Illiterate	Primary school	High school	Higher secondary school	Graduate	Postgraduate
Preterm birth	34 (13.54%)	1 (2.94%)	1 (2.94%)	2 (5.88%)	10 (29.41%)	10 (29.41%)	22 (64.70%)
No effect	55 (21.91%)	15 (27.27%)	8 (14.54%)	13 (23.63%)	7 (12.72%)	7 (12.72%)	5 (9.09%)
Malformation of bones	7 (2.78%)	1 (14.28%)	1 (14.28%)	1 (14.28%)	4 (57.14%)	4 (57.14%)	1 (14.28%)
Cleft Lip/Cleft palate	16 (6.37%)	0 (0%)	1 (6.25%)	1 (6.25%)	4 (25%)	4 (25%)	7 (43.75%)
Do not Know	139 (55.37%)	28 (20.14%)	17 (12.23%)	32 (23.02%)	17 (12.23%)	17 (12.23%)	13 (9.39%)
Chi-squared test = 75.038, df = 20, p-Value = 0.0000							

hyperaemia, edema, and bleeding in periodontal tissues; and these are the risk factors for bacterial infections.⁹ Hence, there arise a need to know the attitude of pregnant women toward their dental care so that oral health promotion strategies can be aimed at in the required direction for early diagnosis and treatment of pregnancy gingivitis and periodontitis, hence leading to prevention of its probable adverse outcomes on new born babies. Thus, the present study was conducted among the child-bearing age group of pregnant women to assess their knowledge, attitude, and perception of pregnancy gingivitis.

When enquired about their awareness of developing gingivitis during pregnancy, majority (42.62%) of the pregnant women said that they had only heard about developing pregnancy gingivitis and were not fully aware about it; other 31.47% of the respondents had not even heard about it, while 25.89% of them were aware about the same. Majority of the women who were fully aware about pregnancy gingivitis were graduates/postgraduates in sync with their educational level, depicting a statistically significant difference between the responses of educated and uneducated respondents. These findings indicate that education has a diverse set of impact on the oral health awareness during pregnancy as is mentioned in previous study conducted by Sunita Bamanikar and Liew KokKee wherein they found vivid difference in good oral health knowledge in educated pregnant women in comparison of those having a little or no education; the reason thereof might be their better social communication with others and hence sharing of views and ideas leading to increase in their level of oral health-related knowledge.¹⁰

In the present study, when the participants were enquired about getting their dental checkup done during their course of pregnancy, majority of them (41.03%) said they did not get their dental checkup done during or before pregnancy. This finding is similar with the previous studies in which it was reported that women decrease or stop visiting the dentists for dental checkups during pregnancy.^{11,12}

This finding is a pointer to misconceptions which the women are having regarding dental care especially during or before pregnancy. It is in contrast to the fact that dental checkup should be undertaken by pregnant women before their pregnancy so that the dentists are able to evaluate their oral health conditions and educate their patients about the expected changes in their oral tissues especially gingiva and also about the preventive measures to be adopted and practiced so as to avoid oral health-related problems. The misconception of harm to the fetus due to dental checkup or dental-treatment leads to aversion of pregnant women to a dentist that ultimately endangers them as well as the oral health of child at later stages.¹³

A good number of respondents (28.68%) agreed that they had swollen and reddened gums; 27.49% of them complained of having bleeding gums, yet other 25.89% said of having bad breath, whereas only 13.9% felt that they had no signs of pregnancy gingivitis. These findings are in concord with previous studies, wherein they found the symptoms like oral malodor, bleeding gums, dental caries, dry mouth, gingival redness persisting among pregnant women; and also stress being one of the causal factors for such poor oral health.^{14,15}

Fig. 1 Frequency distribution of respondents regarding cleaning of oral cavity of children less than 18 months old.

Fig. 2 Frequency distribution of respondents regarding their view about compulsorily having dental education at the start of pregnancy term.

In the present study, 32.66% of respondents did not know about the cause of pregnancy gingivitis, while 31.07% of them reported the hormonal changes to be the cause for the same and 17.92% considered poor oral hygiene as the cause for pregnancy gingivitis. Hormonal changes during pregnancy have been reported in various previously conducted studies as being the main cause of oral tissue changes inducing pregnancy gingivitis, periodontitis, and dental caries. During pregnancy, there is an increase in the metabolism of estrogen by gingiva and also prostaglandins that are synthesized, cause alterations in gingival tissues.¹⁶ The immune system decreases due to change in hormonal (estrogen and progesterone) levels, which has a bearing on the rate of collagen production; and hence body becomes unable to repair the gingiva.¹⁷ Poor oral hygiene was reported, as the cause of pregnancy gingivitis, more by the educated pregnant women than by less/uneducated ones. This finding was analogous to the results showed by previous studies, wherein the authors have mentioned that the health behavior and utilization of health services are directly proportional to the level of education of

pregnant women; the level of education significantly determined the good oral hygiene among pregnant women.¹⁸

In the present study, when the pregnant women were enquired about the measures they believed could prevent pregnancy gingivitis, majority of them (35.05%) were aware of the fact that daily tooth brushing and flossing would be able to prevent the same. Respondents to the tune of 12.74% believed that getting dental checkup done during second trimester would prevent pregnancy gingivitis; and as many as 23.10% of the respondents believed that getting dental checkup done before pregnancy would be beneficial, while as 21.91% of them reported that balanced diet could prevent pregnancy gingivitis. According to the previous literature, all factors such as proper tooth brushing and flossing,¹⁰ regular dental checkups before and during pregnancy, and a balanced diet have their due role in preventing pregnancy gingivitis. In previous studies, it is reported that scaling, polishing, and root planning should be done irrespective of the stage/trimester of pregnancy so as to promote good oral health.^{19,20} It is found that less food intake during pregnancy (especially during first

trimester) was associated with gingival bleeding due to deficient nutrients needed otherwise to protect the gingiva as well as the immune system from getting compromised.²¹

In the present study, around 55.37% of the respondents were unaware about the adverse pregnancy outcome of pregnancy gingivitis and 21.91% believed that it had no such adverse effect, while 13.54% of them reported preterm births as the adverse pregnancy outcome of pregnancy gingivitis. Many previous studies have found the correlation between periodontitis, preterm births, and low weight births. Also maternal periodontitis has been associated with poor oral health of child.^{22,23} Though the association of pregnancy gingivitis/periodontitis with adverse outcomes such as preterm births is still under research, it is better to maintain good oral health by pregnant women to avoid any associated complications as prevention has always been proven to be better than cure.

In the present study, most of the pregnant women (49.40%) reported that using soft moist towel is the proper way of cleaning the oral cavity of children less than 18 months of age. It signifies that most of them were well versed with the way of oral hygiene maintenance among infants, though not all of them were aware about the same. Hence, educative programs for women should include the methods of oral hygiene measures and dental care of infants so as to prevent the development and progression of early childhood caries among infants. Importance of good oral hygiene among infants right from their birth should be emphasized upon so that they develop a healthy primary dentition that would result in the healthy permanent dentition to be carried along throughout life.

Limitations

1. The trimester of pregnancy was not taken into consideration.
2. The oral hygiene practices of the pregnant women were not explored nor were the participants examined for their gingival and periodontal status.
3. The stress level of subjects was not measured.
4. A convenience sample was taken that could limit generalizing of results on larger population.

Conclusion

Within the limits of the study, it was concluded that majority of the pregnant women were unaware about developing gingivitis during their pregnancy period. Most of the women did not know about various measures of prevention and treatment of pregnancy gingivitis and most of the pregnant women did not visit the dentist during pregnancy.

Recommendations

1. Oral health education program should include maternal oral health issues; women should be educated about the importance of good oral health especially during pregnancy and they should be made aware about its adverse effect on the oral health of new born.

2. Pregnant women should be counseled well for undergoing thorough dental examination before and during pregnancy that would prevent pregnancy gingivitis and its outcomes.
3. Preventive and education program should also include the topics regarding balanced, healthy diet needed for the well-being of the expecting mothers as well as the developing fetus.
4. It is recommended that pregnant women should undergo scaling/root planning at any stage of pregnancy without any fear.
5. It is recommended that the dentists should perform surgical procedures such as extractions, restorations, endodontic procedures in second or third trimester, after completion of fetal organogenesis. Moreover, dentists should be aware of the proper seating posture for pregnant women on a dental chair.
6. All health care professionals (dental/medical) should work in collaboration with each other so that they properly opt for relevant referrals for patients especially pregnant women.

Conflict of Interest

None declared.

References

1. Patil S, Thakur R, Madhu. K, Paul ST, Gadicherla P. Oral health coalition: knowledge, attitude, practice behaviours among gynaecologists and dental practitioners. *J Int Oral Health* 2013;5(1):8-15
2. Kandan PM, Menaga V, Kumar RR. Oral health in pregnancy (guidelines to gynaecologists, general physicians & oral health care providers) *J Pak Med Assoc* 2011;61(10):1009-1014
3. Barak S, Oettinger-Barak O, Oettinger M, Machtei EE, Peled M, Ohel G. Common oral manifestations during pregnancy: a review. *Obstet Gynecol Surv* 2003;58(9):624-628
4. Tarsitano BF, Rollings RE. The pregnant dental patient: evaluation and management. *Gen Dent* 1993;41(3):226-234, quiz233-234
5. Lydon-Rochelle MT, Krakowiak P, Huijoe PP, Peters RM. Dental care use and self-reported dental problems in relation to pregnancy. *Am J Public Health* 2004;94(5):765-771
6. al-Kanhal MA, Bani IA. Food habits during pregnancy among Saudi women. *Int J Vitam Nutr Res* 1995;65(3):206-210
7. Michalowicz BS, DiAngelis AJ, Novak MJ, et al. Examining the safety of dental treatment in pregnant women. *J Am Dent Assoc* 2008;139(6):685-695
8. Xiong X, Buekens P, Fraser WD, Beck J, Offenbacher S. Periodontal disease and adverse pregnancy outcomes: a systematic review. *BJOG* 2006;113(2):135-143
9. Taani DQ, Habashneh R, Hammad MM, Batieha A. The periodontal status of pregnant women and its relationship with socio-demographic and clinical variables. *J Oral Rehabil* 2003;30(4):440-445
10. Bamanikar S, Kee LK. Knowledge, attitude and practice of oral and dental healthcare in pregnant women. *Oman Med J* 2013;28(4):288-291
11. Moawad S, Hawsawi A, Al Ahmed SS, Al-Atawi N, Awadien AZ. Knowledge and oral health care practices among Saudi pregnant women. *Life Sci J* 2014;11(5):32-41
12. Al-Swuailam AS, Al-Jamal FS, Helmi MF. Treatment perception and utilization of dental services during pregnancy among

- sampled women in Riyadh, Saudi Arabia. Saudi. J Dent Res 2014;5(2):123–129
- 13 George A, Johnson M, Blinkhorn A, Ellis S, Bhole S, Ajwani S. Promoting oral health during pregnancy: current evidence and implications for Australian midwives. J Clin Nurs 2010;19(23–24):3324–3333
 - 14 Peruzzo DC, Benatti BB, Ambrosano GM, et al. A systematic review of stress and psychological factors as possible risk factors for periodontal disease. J Periodontol 2007;78(8):1491–1504
 - 15 Kumar J, Samelson R. Oral Health Care during Pregnancy and Early Childhood: Practice Guidelines. New York, NY: New York State Department of Health; 2006
 - 16 Lee A, McWilliams M, Janchar T. Care of the pregnant patient in the dental office. Dent Clin North Am 1999;43(3):485–494
 - 17 Hey-Hadavi JH. Women's oral health issues: sex differences and clinical implications. Women's Health Prim Care 2002;5(3): 189–199
 - 18 Yalcin F, Eskinazi E, Soydinc M, et al. The effect of sociocultural status on periodontal conditions in pregnancy. J Periodontol 2002;73(2):178–182
 - 19 Trivedi S, Lal N, Singhal R. Periodontal diseases and pregnancy. J Orol Sci 2015;7:67–68
 - 20 Huda S, Doering H, Tenenbaum HC, Whittle W, Sigal MJ, Glogauer M. Oral neutrophil levels: a screening test for oral inflammatory load in pregnancy in a medical setting. J Periodontol 2015;86(1):72–81
 - 21 Gaffar BO, El Tantawi MM, AlAgf AS, Al-Ansari A. Food intake, oral hygiene and gingival bleeding in pregnancy: does lifestyle make a difference? A cross sectional exploratory study. Int J Health Sci (Qassim) 2016;10(1):13–20
 - 22 Casamassimo PS. Maternal oral health. Dent Clin North Am 2001;45(3):469–478,v–vi
 - 23 Gajendra S, Kumar JV. Oral health and pregnancy: a review. N Y State Dent J 2004;70(1):40–44

Appendix A: Questionnaire

Q1 Are you aware that women can develop pregnancy gingivitis (swollen red gums) during pregnancy?

- (a) Yes, I am fully aware
- (b) No, I am not aware
- (c) I am not fully aware, but heard about it.

Q2 Did you get a dental checkup done before or during the period of your pregnancy?

- (a) No, I did not
- (b) Yes, I did
- (c) I planned to, but did not.
- (d) Yes, due to dental condition

Q3 Do you have any of the following signs of pregnancy gingivitis?

- (a) Swollen and reddened gums
- (b) Bad breath
- (c) Gum bleeding
- (d) Red spot over oral lining
- (e) None

Q4 What according to your knowledge may be the cause of gingivitis (red and swollen gums) during pregnancy?

- (a) Hormonal changes
- (b) Traumatic tooth brushing
- (c) Poor oral hygiene
- (d) Eating fried food
- (e) Malnutrition
- (f) Do not know

Q5 What measures according to your knowledge can prevent pregnancy gingivitis?

- (a) Daily tooth brushing and flossing
- (b) Balanced diet
- (c) Dental checkup before pregnancy
- (d) Scaling every month
- (e) Dental checkup during second trimester
- (f) Do not know

Q6 What according to your knowledge is the treatment, dentists provide for pregnancy gingivitis?

- (a) Surgical removal of swollen gum
- (b) No treatment needed
- (c) Professional scaling
- (d) Extract the affected tooth
- (e) Medication
- (f) Do not know

Q7 What effect can pregnancy gingivitis have on your oral health?

- (a) Overgrowth of gum tissue
- (b) No effect
- (c) Periodontal disease
- (d) Tooth decay
- (e) Tooth sensitivity
- (f) Tooth becoming loose
- (g) Do not know

Q8 Will pregnancy gingivitis have effect on oral health of your new born?

- (a) Yes, it will
- (b) No, it will not
- (c) Undecided

Q9 What effect can gingivitis possibly have on your pregnancy or the child to be born?

- (a) Preterm birth
- (b) No effect
- (c) Malformation of bones
- (d) Cleft lip and palate
- (e) Do not know

Q10 What according to you is the proper way of cleaning the oral cavity of children less than 18 months of age?

- (a) Tooth brush and paste
- (b) Soft moist towel
- (c) Electric tooth brush
- (d) Mouthwash
- (e) Saline

Q11 Do you suggest that pregnant women should have compulsory dental health education at the start of the pregnancy term?

- (a) Yes
- (b) No
- (c) Undecided