Free Flap Skin Paddle: A Painless Donor Site In Vivo

Anupam Golash¹ Raunak Shinde^{1,*} Deepanjan Dey¹

¹Department of Plastic Surgery, Calcutta Medical Research Institute, Kolkata, India

Address for correspondence Raunak Shinde, MS (Gen Surg), MCh (Plastic Surgery), Department of Plastic Surgery, HB 5, Muni Nagar, Ujjain 456010, Madhya Pradesh, India (e-mail: drraunak@hotmail.com).

Indian J Plast Surg 2019;52:256–257

Dear Sir,

Free flap reconstruction is an essential tool in the coverage of composite large defects for varied etiology and at varied anatomical locations. Often, in case of composite defects where the critical area of the wound is covered by the free flap, some raw area may remain at the recipient site that needs to be covered secondarily. We present an innovative way to address the issue by using the free flap skin paddle as the donor site a skin graft. Most of the times, in our clinical practice, the remnant raw area was often large enough to require a small skin graft for healing. In such cases, the skin paddle of the free flap can be used as the donor site, avoiding morbidity at a new anatomical location.

Once the flap is deemed reasonably stable (around 3–4 weeks) and the pending raw area is amenable for grafting, the skin paddle should be stabilized, gently stretched, and, with the help of skin grafting handle/dermatome, graft can be harvested from the skin paddle and applied to the raw area.

Following is a case example illustrating the use of this technique.

A middle-aged female presented with posttraumatic scalp avulsion with the skull bone exposed. The wound covered with free latissimus dorsi musculocutaneous flap with skin graft over the muscle. A residual area due to graft loss over the muscle was addressed with skin graft from the skin paddle of the free flap. The recipient raw area and the graft donor site healed well (► Figs. 1–4).

Fig. 1 Preoperative-posttraumatic scalp avulsion with exposed skull bone.

Fig. 2 Postreconstruction with free latissimus dorsi myocutaneous flap with the remnant raw area.

Fig. 3 Skin paddle of latissimus dorsi flap as the donor site for split-thickness skin graft.

Fig. 4 Healed donor site and raw area at 2 weeks.

Conflict of Interest

None.

References

- 1 Dabernig J, Watson S. Flap skin as a donor site for split skin grafts. *Ann Plast Surg* 2005;55(4):439–440
- 2 Li AKM, Ghanem T, Use of skin grafts in free flap reconstruction. *InSkin Grafts*. London, UK: IntechOpen Limited; 2013