unfavorable outcome, while 7 died. At a cutoff of 3, STEPSS had 93.3% sensitivity, 80.8% specificity, and negative predictive value of 99% for unfavorable outcome. The negative predictive values for death and treatment failure (refractory or super-refractory status epilepticus) were 99% and 98%, respectively. At a cutoff of 2, STEPSS had 100% sensitivity, 60.6% specificity for unfavorable outcome, and 100% negative predictive value for unfavorable outcome, death, and treatment failure. The predictive accuracy was comparable to that of STESS in adults.

Conclusion: The STEPSS, a simple bedside clinical score was found to be useful to predict the outcome and treatment response in children with status epilepticus.

A0042: Role of Pyridoxal Phosphate in Modulation of Endogenous Kynurenic Acid Synthesis Associated with Hippocampal Sclerosis

Soumil Dey,¹ Aparna Dixit,¹ Manjari Tripathi,¹ P. Sarat Chandra,¹ Jyotirmoy Banerjee¹

¹Center of Excellence for Epilepsy, A Collaborative Project between All India Institute of Medical Sciences, New Delhi, National Brain Research Center, Manesar, India

Objective: Hippocampal sclerosis (HS) is the most common form of drug-resistant epilepsy where temporal lobe structures are responsible for unprovoked seizures. The hallmark of HS is enhanced glutamatergic excitatory neurotransmission. Pyridoxal 5-phosphate (pLP) is an active form of vitamin B6 and plays an important role as a cofactor of various enzymes including kynurenine aminotransferase II (KAT II), which catalyzes kynurenic acid (KYNA) synthesis from kynurenine within cortical astrocytes. KYNA, a tryptophan metabolite, is the only natural inhibitor of glutamate receptors. The present study was designed to test the hypothesis that in HS, altered PLP concentration is responsible for reduced endogenous kynurenic acid synthesis in hippocampus.

Methods: Hippocampus from HS patients ($n = 33$) were used for this study. Tissues resected from tumor margin during brain tumor surgery of seizure-free patients as nonepileptic control ($n = 14$) were used. To determine total KYNA and PLP concentration, tissues were kept in perchloric acid at -80°C . Tissues were homogenized, centrifuged, and supernatants were collected and estimated using HPLC with fluorescence detection (pLP excitation 300 nm, emission 400 nm; KYNA excitation 344 nm, and emission 404 nm).

Results: KYNA concentration was significantly less in HS hippocampus (0.0244 ± 0.0219 ng/mg of protein) compared with nonepileptic controls (0.2117 ± 0.1332 ng/mg of protein). PLP concentration was also significantly less in HS patients (1.77 ± 3.96 mg/mg of protein) compared with controls (10.91 ± 11.34 ng/mg of protein).

Conclusion: PLP concentration is reduced which, we suspect, may consequences in decreased KYNA due to dysfunctional machinery to synthesize kynurenic acid from kynurenine in HS patients.

Acknowledgment This work was supported by the Centre of Excellence for Epilepsy, a collaborative project between All India Institute of Medical Sciences, New Delhi,

and National Brain Research Centre, Manesar; and funded by Department of Biotechnology, Ministry of Science and Technology, Govt. of India (Grant: BT/01/COE/09/08).

A0043: A Home-Based, Primary-Care Model for Epilepsy Care in Low and Middle Income Countries: Basis and Design

Suman Sharma,¹ Anuraag Chowdhury,¹ Sarit Sharma,¹ Rajinder Bansal,¹ Gagandeep Singh¹

¹Department of Neurology, Dayanand Medical College, Ludhiana, Punjab, India

Objective: A cluster-randomized trial of home-based care using primary care resources for people with epilepsy has been set up to optimize epilepsy care in resource-limited communities in low- and middle- income countries. The primary aim is to determine whether treatment adherence to antiepileptic drugs is better with home-based care or with routine clinic-based care. Secondary aims are to compare the effects of the two care pathways on seizure control and quality of life.

Methods: The home-based intervention comprises epilepsy medication provision, adherence reinforcement, and epilepsy self-management and stigma management guidance provided by an auxiliary nurse-midwife equivalent. The experimental group will be compared with a routine clinic-based care group using a cluster-randomized design in which the unit of analysis is a cluster of 10 people with epilepsy residing in an area cared for by a single accredited government grass-root health worker. The primary outcome is treatment adherence as measured by monthly tablet counts and two self-completed questionnaires. The secondary outcomes include monthly seizure-frequency, time to first seizure (in days) after enrolment, proportion of subjects experiencing seizure freedom for the duration of the study and quality of life measured by the "Personal Impact of Epilepsy Scale," all assessed by an independent study nurse.

Results: The screening phase and neurological evaluations and randomizations have been recently completed and follow-up is currently underway.

Conclusion: The Results of the trial are likely to have substantial bearing on the development of governmental policies and strategies to provide coverage and care for epilepsy in resource-limited countries.

A0044: Clinical and Demographic Profile of Epilepsy Patients in Rural Rajasthan

Surbhi Chaturvedi,¹ R.K Sureka,¹ Amit Agarwal¹

¹Mahatma Gandhi Medical College and Hospital, Jaipur, Rajasthan, India

Objective: The present study attempted to explore the clinical and demographic profile of epileptic patients, and pharmacological management in these patients.

Methods: The prospective study was done in center situated at Ratannagar, a Community Health Centre in district Churu under the auspicious of Epilepsy Care and Research foundation, an NGO involved with epilepsy work. It included 6,993 epileptic patients who visited the center from majority

of states from all over India from 1994 to 2018. Seizures were classified according to the International League against Epilepsy Classification (1981). Detailed clinical history and neurological examination was done in all patients. Routine blood tests, computerized tomography of head, and electroencephalography (EEG) were done only in some patients due to nonavailability, limited resources, and poor economic strata of the patients. All patients were given free medicines and were followed-up monthly for assessing the compliance, the response, and side effects of antiepileptic drugs.

Results: Male: female ratio was 2.51. The 36.45% patients were of age group 21 to 30 years and 20.9% were of age group from 11 to 20 years. Most of the patients (86%) were from rural areas. GTCS was the most common seizure type accounting for 67% followed by complex partial seizures (21%). In 68% of the patients no cause could be found for the seizures and they were labeled idiopathic. The commonest causes for symptomatic epilepsy were hypoxic ischemic insult and CNS infections. Majority of the patients were fully controlled on drugs. Fifty percent patients were controlled on monotherapy, 30% were controlled on two drugs, and 18% patients required more than two drugs. The most common drug used was phenytoin followed by phenobarbitone and levetiracetam, sodium valproate, and carbamazepine. Noncompliance of the drug was found to be the most common cause of the recurrent episodes of seizures followed by sleep deprivation and fever. The incidence of mental retardation was 12% and behavioral disorder was 22%. Loss of memory was reported in 48% of the patients, which was relatively very high.

Conclusion: Preventable causes of epilepsy share a significant portion in the etiology of the disease. Maximum patients can be managed at rural center without sophisticated investigations.

A0045: Bilateral Rasmussen's Encephalitis: A Rare and Difficult Entity—Case Report

Swapan Gupta,¹ Neera Chaudhry,¹ Vinod Puri¹

¹G.B. Pant Institute of Postgraduate Medical Education and Research (GIPMER), Delhi, India

Introduction: Rasmussen encephalitis is a rare chronic disorder characterized by unilateral inflammation of the cerebral cortex, drug resistant epilepsy, and progressive neurological deterioration. Functional hemispherotomy is the choice of management in drug-resistant cases. The existence of bilateral disease is debated.

Case Report: A 12-year-old girl, born of a nonconsanguineous caesarean section with normal milestones had seizures since 9 years of age. She started with left focal motor (tonic-clonic) seizures with lower limb onset, which evolved to *epilepsia partialis continua* (EPC) over time, resistant to multiple antiepileptics. Her MRI revealed right temporo-parieto-occipital gliosis and atrophy and EEG showed right hemispherical slowing with superimposed epileptiform discharges. Left hemisphere was completely normal on EEG. She was diagnosed as a case of Rasmussen's encephalitis after ruling out other etiologies and given immunotherapy. After initial failure with steroids, IVIG and pheresis, she

later responded favorably with rituximab but after 6 months, developed right-sided focal motor seizures with lower limb onset which over a short period of time progressed to EPC. EEG revealed slowing in bilateral hemispheres with left hemispherical interictal epileptiform discharges and seizure onset from left-sided (central and midline) channels. MRI revealed T2/FLAIR hyperintense signal in left frontal parasagittal region corresponding to the leg region. She was again given rituximab with modest benefit.

Conclusion: Rasmussen's encephalitis can be rarely bilateral. Management of such patients is difficult since surgical option is not feasible owing to bilaterality.

A0046: Early Onset Seizures in Vacuolating Megalencephalic Leukoencephalopathy with Subcortical Cysts (MLC-Vander Knaap Disease) in a Brahmin Family Puja Rani,¹ Swapan Gupta,¹ Geeta Anjum Khwaja¹

¹Department of Neurology, G.B. Pant Institute of Postgraduate Medical Education and Research (GIPMER), Delhi, India

Background: Epileptic seizures are common and occur early in most gray matter diseases but are rare in white matter disease. MLC, a white matter disease, has been mostly reported from the Agarwal community in India. It usually presents as macrocephaly with developmental delay, cerebellar and pyramidal dysfunction. We report a rare case of MLC from a Brahmin family, with early onset seizure as a rare manifestation.

Case: An 11-year-old boy from Uttarakhand, a Northern state of India, presented with recurrent seizures since age of age. The seizure frequency was variable and semiology was focal motor, evolving to bilateral tonic-clonic seizures. He was born out of nonconsanguineous marriage and belonged to the Brahmin community. Birth history was normal but there was a history of delayed motor and mental milestones. His elder sibling also had recurrent seizures with developmental delay and both had been labeled as cases of cerebral palsy with refractory epilepsy. EEG showed focal as well as generalized spike, polyspike, and sharp wave discharges with eye closure sensitivity. MRI brain was suggestive of diffuse white matter changes with subcortical cysts. Genetic analysis proved the diagnosis of MLC in our case.

Conclusion: MLC which has primarily been reported from the Agarwal community in India can also occur in other communities, and seizures can be an early and initial manifestation of this white matter disorder. This condition is probably under-reported because of its rare occurrence and misdiagnosis.

A0047: Effect of Pentylentetrazole-Induced Seizures on Serum Brain-Derived Neurotrophic Factor (BDNF) Levels and Cognition in Rats

Vibha Patel,¹ Jatinder Katyal,¹ Haroon Rashid,¹ Yogendra K. Gupta¹

¹Neuropharmacology Laboratory, Department of Pharmacology, All India Institute of Medical Sciences, New Delhi, India

Objective: Pentylentetrazole (pTZ)-induced seizures affect the serum brain-derived neurotrophic factor (BDNF) levels and also cognition in rats. In this study, effect of acute