Guest Editorial

Cerebral Artery Hypoplasia among Black Kenyans

Bukkambudhi V. Murlimanju¹

¹Department of Anatomy, Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Mangalore, Karnataka, India

J Neurosci Rural Pract 2019;10:392

Mortality due to stroke has declined but the global burden of stroke is still in increase.¹ The vascular anatomy and pathology have to be clearly understood for the effective management of the stroke. The circulus arteriosus (Circle of Willis) is an arterial anastomosis which maintains the cerebral blood flow. The strokes are of two varieties, ischemic and hemorrhagic. Among all the ischemic strokes, the anterior cerebral artery territory constitutes only approximately 0.6 to 3%.² The cerebral blood flow of different anatomical variations can be of help for the understanding of collateral circulation.³ In this study by Ogengo et al,⁴ the authors made an attempt to observe the cerebral artery hypoplasia in the East African sample population. The “Charcot artery of cerebral hemorrhage,” which is a branch of the middle cerebral artery, is the most common artery to bleed. Here, in this study from the Kenyan population, Ogengo et al⁴ observed that the middle cerebral artery is least commonly affected with the hypoplasia. This suggests that the middle cerebral artery is prone for hemorrhagic stroke than ischemic. In this study, 73.4% of the cadaveric brain specimens showed the arterial hypoplasia. Perhaps this is the reason for most of the people who suffer from headache, migraine, and transient cerebral ischemia. Hypertension, cigarette smoking, and diabetes are the risk factors for the stroke.⁵ Cerebrovascular accidents can be prevented by taking aspirin prophylaxis and lifestyle modification, like routine exercises. However, few things are genetically determined that persist and progress into the adult life.

In this Kenyan African study,⁴ the authors observed that the posterior communicating artery, which connects the carotid and vertebral arterial system, is more prone for hypoplasia. This factor may hinder the effective anastomosis between

the major arterial inputs of brain. The authors have also compared their observation bilaterally for the paired arteries. The frequency was also compared with other population groups of various nations, globally. I believe that this type of population-based studies is of paramount importance. These studies will help the clinician to have effective population-based strategies such as control of risk factors. The literature will aid in the reduction of stroke-related morbidity and mortality.⁶

Funding

None.

Conflict of Interest

None declared.

References

- 1 Hankey GJ. Stroke. Lancet 2017;389(10069):641–654
- 2 Lakhotia M, Pahadiya HR, Prajapati GR, Choudhary A, Gandhi R, Jangid H. A case of anterior cerebral artery A1 segment hypoplasia syndrome presenting with right lower limb monoplegia, abulia, and urinary incontinence. J Neurosci Rural Pract 2016;7(1):189–191
- 3 Zhu G, Yuan Q, Yang J, Yeo J. Experimental study of hemodynamics in the Circle of Willis. Biomed Eng Online 2015;14(Suppl 1):S10
- 4 Ogengo JA, Cheruiyot I, Amuti T, et al. Cerebral artery hypoplasia in a select Kenyan adult population. J Neurosci Rural Pract 2019;10(3):423–429
- 5 Dhiman D, Mahajan SK, Sharma S, Raina R. The evolving pattern and outcome of stroke at moderate altitude. J Neurosci Rural Pract 2018;9(1):68–72
- 6 Verma R. Stroke: a neglected epidemic in India. J Neurosci Rural Pract 2018;9(4):453–454

Address for correspondence

Bukkambudhi V. Murlimanju, MD,
Department of Anatomy, Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Mangalore 575001, Karnataka, India
(e-mail: flutesnowmm@gmail.com).

DOI <https://doi.org/10.1055/s-0039-1695695>
ISSN 0976-3147.

©2019 Association for Helping Neurosurgical Sick People

License terms