Endoscopic Endonasal Transodontoid Approach for Degenerative Pseudotumor of the Craniocervical Junction

Andrew S. Venteicher^{1,*} Ezequiel Goldschmidt^{2,*} Michael M. McDowell² Eric W. Wang³
 Carl H. Snyderman³ Paul A. Gardner²

¹Department of Neurosurgery, University of Minnesota, Minneapolis, Minnesota, United States

²Department of Neurological Surgery, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, United States

³Department of Otolaryngology, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, United States

Address for correspondence Paul A. Gardner, MD, Department of Neurological Surgery, 200 Lothrop Street, Suite B400, University of Pittsburgh Medical Center, Pittsburgh, PA 15213, United States (e-mail: gardpa@upmc.edu).

J Neurol Surg B 2019;80(suppl S4):S368–S369.

Abstract

Keywords

- transodontoid
- endoscopic endonasal approach
- craniocervical junction
- retroodontoid degenerative pseudotumor
- periodontoid pannus

Ventral masses of the craniocervical junction threaten the medulla and upper spinal cord leading to lower cranial nerve dysfunction and myelopathy. Traditional transoral and newer endonasal approaches can access ventral pathology in this region, though both remain challenging due to the competing goals of achieving sufficient decompression, yet mitigating risk to nearby critical neurovascular structures. Transoral approaches have traditionally been avoided for degenerative pseudotumor which generally slowly regresses following posterior fixation. Here, we present two cases of patients with significant retroodontoid degenerative pseudotumor and pannus causing dysphagia and myelopathy. The first patient was found with a large broad pannus requiring a wide decompression extending from lower clivus to inferior aspect of the C1 arch and odontoid process. The second patient had a more focal mass effect due to pannus and synovial cyst at the level of the inferior half of C1 and midodontoid. Both patients underwent an endoscopic endonasal transodontoid approach for immediate decompression followed by a posterior C1–C2 fusion. In these surgical videos, we highlight anatomic considerations in this critical area, demonstrate nuances of technique, and outline strategies to avoid complications and maximize exposure. The link to the video can be found at: <https://youtu.be/19l-GQYGIr4>.

www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

* Co-first authors.

received
 March 30, 2019
 accepted after revision
 August 25, 2019
 published online
 November 4, 2019

DOI <https://doi.org/10.1055/s-0039-1700892>.
 ISSN 2193-6331.

© 2019 Georg Thieme Verlag KG
 Stuttgart · New York

License terms

Fig. 1 Patient 1 with broad retroodontoid pseudotumor causing cervicomedullary stenosis. (A) Preoperative MRI T2. (B) Postoperative CT. (C) Endoscopic view after pannus was resected. CT, computed tomography; MRI, magnetic resonance imaging.

Fig. 2 Patient 2 with retroodontoid pseudotumor and synovial cyst causing cervicomedullary stenosis. (A) Preoperative MRI T2. (B) Postoperative CT. (C) Endoscopic view removing synovial cyst wall. CT, computed tomography; MRI, magnetic resonance imaging.

Publication Comments

The authors present two cases of endonasal endoscopic resection of the odontoid for pannus and a synovial cyst. Use of a myofascial flap is demonstrated.

Jeffery Sorenson, MD
University of Tennessee Health Science Center
Memphis, TN

Conflict of Interest

Dr. Gardner reports other from SPIWay, LLC, outside the submitted work; Dr. Snyderman reports other from SPIWay, LLC, outside the submitted work.

Acknowledgments

The authors thank the Rhoton Collection for access to anatomic photographs used in this video. We also thank Paul Stanick for assistance in producing this video.