Current Trends in Practice of Residents in the Saudi Board of Endodontics Program

Mothanna K. AlRahabi¹, Ayman M. AlKady¹

¹Department of Restorative Dental Science, College of Dentistry, Taibah University, Al Madinah Al Munawwarah, Kingdom of Saudi Arabia

Address for correspondence Mothanna AlRahabi, MSc, PhD, Department of Restorative Dental Science, College of Dentistry, Taibah University, PO Box 2898, Al Madinah Al Munawwarah 43353, Kingdom of Saudi Arabia (e-mail: dr.rahabi@gmail.com; mrahabi@taibahu.edu.sa).

Eur J Dent 2019;13:619–624

Abstract

Objectives The purpose of the current study was to collect information about some techniques and armamentarium currently used by the Saudi Board of Endodontics residents in relation to the technical steps of root canal therapy.

Materials and Methods A web-based survey was sent to the Saudi Board of Endodontics residents, based in the western area of Saudi Arabia. The survey assessed controversial concepts, and collected information regarding new instruments and materials, used in the technical steps of nonsurgical root canal treatment. A one-sample chi-square test, with a 95% level of significance, was applied to determine whether there were significant differences between respondents' answers.

Results A total of 45 out of 50 residents (90%) responded to the questionnaire. The majority of Saudi Board residents of Endodontics, who participated in this survey, used the mean of working length (ML) measured by periapical X-Ray and apex locator in the presence of a radiographic lesion (63.3%), and maintained apical patency in all cases (80.0%). ProTaper Universal (40%) and ProTaper Next (41.2%) were the most common NiTi rotary systems used for root canal instrumentation. Thirty percent of respondents used adjunct device with irrigation, and 80% advocated smear layer removal. Zinc oxide–eugenol-based sealers were the most common used sealers (70%), and most residents (86.6%) did not advocate sealer extrusion.

Conclusions This study revealed that residents of the Saudi Board of Endodontics program adopt new endodontic technologies. There is, however, a need for more investigations regarding this objective, including responses from all residents in Saudi Arabia.

Keywords

- Saudi Board of Endodontics
- endodontic therapy
- radiography
- root canal
- surveys
- technology

Introduction

Recent years have witnessed a revolution in trends and technologies within the field of endodontics.^{1,2} Electronic apex locators, smart electronic motors, nickel–titanium rotary and reciprocating files, irrigation devices and activators, ultrasonic endodontic tips, operating microscope, thermoplastic obturation devices, and cone beam-computed tomography, are all advances that have enhanced and improved the practice of root canal therapy,³ thereby affecting the daily practice of endodontics.⁴ A questionnaire survey is a common tool to

collect information regarding attitudes toward new advances and technologies in the endodontics world.^{3,5–9} The following surveys have been performed with respect to different parts of endodontics practice: irrigation solutions and methods,^{6,10} antibiotic use by endodontists,¹¹ cone beam-computed tomography in endodontic practice,¹² endodontic procedures related to mineral trioxide aggregate usage,¹³ confidence in performing endodontic treatment,¹⁴ cleaning and shaping of root canal systems with nickel–titanium instruments,¹⁵ magnification,⁷ and knowledge and attitude in the management of dental trauma.¹⁶ A decreasing dentists-to-population ratio,

and the growing tendency for patient referral from general practitioners to specialty dentists, have both made it necessary to increase the number of specialized dentists.¹⁷ The Saudi Commission for Health Specialties (SCFHS) established the Saudi Board of Endodontics in 2007. This four-year program has been designed to graduate endodontists with satisfactory knowledge, skills, and clinical experience, along with sufficient background in basic biological science. These objectives are achieved by clinical training, lectures, book reviews, current and classic literature, seminars, quizzes, as well as case presentations. Until now, there have been no studies regarding the trends of Saudi Board of Endodontics' residents during their training and practice of daily endodontic treatment. The purpose of this study was to collect information about certain techniques and armamentarium currently used by them to perform the technical steps of root canal therapy.

Materials and Methods

This study was approved by the research ethics committee of the College of Dentistry-Taibah University Protocol no. TUCDREC/20171220/AlRahabi in agreement with the guidelines of the Helsinki Declaration revised in 1975. A web-based survey, created using Google forms, was emailed to the residents of Saudi Board of Endodontics in Saudi Arabia. The survey included the aim and the importance of this research and seven multiple choices; ►Table 1 shows the included questions. The link was sent to 50 residents. E-mail reminders were sent to those who had not responded within 3 weeks after the first message. The survey was designed to address controversial concepts, and collect information regarding certain instruments and materials, used recently in relation to the technical steps of nonsurgical root canal treatment: modification of working length (WL) in the presence of radiographic lesion; maintenance of apical patency of the root canal during cleaning and shaping; NiTi system used for root canal instrumentation; use of an adjunct devices for root canal irrigation during cleaning and shaping; attitude regarding smear layer removal; type of sealer used for root canal obturation; and attitude regarding sealer extrusion from the apex; ►Table 1 shows the questionnaire items. A one-sample chi-square test, at 95% significance level, was applied to determine whether there were significant differences among residents' answers.

Results

A total of 45 out of 50 residents (90%) responded to the questionnaire. The frequency of residents—who depended on the mean of WL, as measured by periapical X-Ray and apex locator in the presence of a radiographic lesion (63.3%)—was significantly higher than those who measured the WL by apex locator only or by reducing the length by 1 to 2 mm from radiographic apex. Maintaining apical patency in all cases was significantly higher (80%) than

Table 1 Questionnaire items

No.	Question	Answers
1	Do you advocate changing working length based on the presence of a radiographic lesion?	1. Yes, I advocate reducing WL 1–2 mm from radiographic apex 2. We use measurements of apex locator 3. We use measurements of apex locator and paper point bleeding point 4. We use the mean of WL measured by X-Ray and apex locator
2	Do you maintain apical patency?	1. In all cases 2. In necrotic cases 3. In retreatment cases 4. In necrotic and retreatment cases
3	What NiTi system do you use for root canal instrumentation?	1. Not specific 2. ProTaper Universal 3. ProTaper Next 4. Reciproc 5. Vortex blue 6. Others specify
4	Do you use any adjunct device to activate irrigation?	1. No activation 2. Endo activator 3. Endo vac 4. Others specify
5	Do you advocate remove smear layer?	1. Yes 2. No 3. There is no difference
6	What is the sealer type do you use?	1. Zinc oxide–eugenol sealers 2. Calcium hydroxide sealers 3. GIC sealers 4. Epoxy or methacrylate resins sealers 5. Bioceram sealers 6. Others specify
7	Do you advocate sealer extrusion from the apex?	1. I do not advocate sealer extrusion 2. I advocate sealer extrusion 3. I advocate sealer extrusion in necrotic cases 4. I advocate sealer extrusion in retreatment cases 5. I advocate sealer extrusion in all cases

Abbreviation: WL, working length.

those who only maintained it in cases of necrosis. ProTaper Next (43.3%) and ProTaper Universal (40%) NiTi rotary systems were significantly used more than other types of NiTi systems. The majority of respondents (70%) performed irrigation of root canal system without any type activation. Most of the respondents (80%) were routinely removing the smear layer during endodontic treatment. Zinc oxide–eugenol-based sealers were the most commonly used sealers (66.7%) and significantly higher than other types of sealers. The majority of respondents (86.6%) were against the sealer extrusion from apical foramen. The responses of participants were summarized in ►Table 2 using frequencies and percentages for different items on the

Table 2 Frequencies and percentages for residents' answers to questionnaire items

No.	Studied Item	Answers	Number of respondents	Percentage
1	Determination of working length in the presence of a radiographic lesion	Reduce WL 1–2 mm from radiographic apex	1	3.3%
		Using the measurements of apex locator	15	33.3%
		Use the mean of WL, as measured by X-Ray and apex locator	29	63.3%
2	Maintain apical patency	In all cases	36	80.0%
		In necrotic and retreatment cases	9	20.0%
3	NiTi system used for root canal instrumentation	No specific preference	1	3.3%
		ProTaper Universal	18	40.0%
		ProTaper Next	19	41.2%
		Reciproc	6	12.2%
		Vortex blue	1	3.3%
4	Use of adjunct device for irrigation	No activation	31	70.0%
		Endo activator	12	26.7%
		Endo vac	2	3.3%
5	Removal of smear layer	Advocate smear layer removal	36	80.0%
		Do not advocate smear layer removal	1	3.3%
		No difference in outcome	8	16.7%
6	Sealer type	Zinc oxide–eugenol sealers	30	66.7%
		Calcium hydroxide sealers	5	11.1%
		Epoxy or methacrylate resin sealers	5	11.1%
		Bioceramic sealers	5	11.1%
7	Advocate sealer extrusion from the apex	Do not advocate sealer extrusion	39	86.6%
		Advocate sealer extrusion in necrotic cases	1	3.3%
		Advocate sealer extrusion in all cases	5	11.1%

Abbreviation: WL, working length.

questionnaire. The attitudes of residents toward investigated techniques and armamentarium during their practice are shown in ► **Figs. 1 and 2.**

Discussion

The present study was designed to collect information regarding the trends and attitudes of the residents of Saudi Board of Endodontics toward some **controversial** concepts, and new instruments and materials, used during nonsurgical root canal treatment. The majority of participants used the mean of WL, as measured by both periapical radiograph and apex locator; only 33.3% depended on the measurement of apex locator. The recommended method to measure WL involves electronic devices, followed by radiographic confirmation.¹⁸ Modern apex locators has proven to be a reliable device for WL measurement¹⁹. Maintaining apical patency in all cases

was significantly higher than those who only maintained it in cases of necrosis According to the American Association of Endodontists, apical patency is a technique where the apical portion of the canal is maintained free of debris by recapitulation with a small file through the apical foramen²⁰. Since 1997, 50% of dental schools in the United States have taught the patency concept²¹. One study revealed that maintaining apical patency did not introduce microorganisms into the periapical tissues²² and improved irrigation efficiency in the apical third.²³ Further, maintenance of apical patency did not increase postoperative pain of root canal treatment in necrotic cases, where all these cases in this study had preoperative radiolucency.²⁴ NiTi rotary instrumentation is considered a basic component of the recent endodontic practice. Several rotary nickel–titanium (Ni–Ti) file systems have been introduced for the preparation of root canals. Ni–Ti instruments provide many advantages compared with conventional files.

Fig. 1 Attitudes of Saudi Board of Endodontics residents during their training part 1.

Fig. 2 Attitudes of Saudi Board of Endodontics residents during their training part 2.

Increased flexibility and shortened working time are the major advantages of Ni–Ti files.²⁵ In this study, ProTaper Next and ProTaper Universal NiTi rotary systems were significantly used more than other types of NiTi systems. However, the selection of NiTi system maybe be affected by their availability. ProTaper systems are produced by Dentsply-Tulsa Dental [Oklahoma, USA], which has an active marketing strategy that might explain the widespread preference for this system.²⁶ Both ProTaper Universal and ProTaper Next can preserve the original curvature of the canal.²⁷ Reciproc, which is a single-file system was used by 12.2% of participants in this survey, has been specifically designed for use in reciprocating motion, with high resistance to cyclic fatigue,²⁸ Moreover, the use of only one NiTi instrument is more cost-effective.²⁹ However, the selection of NiTi system maybe be affected by the

availability or desire to experiment with a new NiTi system. Vortex Blue NiTi system was used with a small percentage of participants, and it can also be attributed to availability or marketing. Vortex Blue has a high-cyclic fatigue resistance.³⁰ A recent survey revealed that 82% of respondents stated they used a multi-file rotary system when performing root canal treatment (RCT), and 18% reported using both multi-file and single-file systems. the most widely used belonged to the ProTaper multi-file brands, accounting for 78% of responses. The most used single-file reciprocation brands were WaveOne WO and WaveOne Gold WOG, used by 26% of respondents.³¹ This study showed that the majority of residents did not use adjunct devices to activate irrigants. However, this maybe be affected by the availability. In one report to evaluate current trends in irrigation among American Association

of Endodontists members, half of the respondents were using an adjunct, such as ultrasonic activation, to aid in their irrigation technique.⁶ EndoActivator system showed significant effectiveness in removal of collagen from the canal surface, relative to syringe irrigation alone.³² The use of sonic activation with EndoActivator did not significantly improve sealer penetration compared with conventional irrigation.³³ There was a significant percentage of residents who advocated smear layer removal. A web-based Survey emailed to the American Association of Endodontists revealed that endodontists routinely remove the smear layer during endodontic treatment.⁶ One study showed that smear layer presence might decrease the efficiency of sodium hypochlorite irrigant,³⁴ while another study showed that smear layer removal improves sealer penetration into dentinal tubules.³⁵ In this study, the most common sealers used by residents were zinc oxide–eugenol-based sealers. This selection maybe be affected by the availability and marketing. Zinc oxide–eugenol root canal sealer, showed minimal microleakage compared with calcium hydroxide and resin-based sealers.³⁶ Zinc oxide–eugenol, and bioceramic and resin-based root canal sealers exhibited antibacterial effects against *E. faecalis* in the dentinal tubules. Bioceramic and resin-based root canal sealers exhibited superior antibacterial effects compared with zinc oxide–eugenol root canal sealers,^{37,38} as their antibacterial efficacy continued after setting.³⁸ Cytotoxicity and genotoxicity of bioceramic-based sealers were less than resin-based sealers AH Plus.³⁹ The majority of residents did not advocate sealer extrusion. Extrusion of obturation material may result in an undesired outcome, such as inflammation and severe neurotoxic damage.⁴⁰ Information regarding the effect of apical extrusion of sealer on root canal treatment outcome is scarce.⁴¹ To evaluate the radiographic healing of a periapical lesion of permanent teeth, after extrusion of an AH Plus sealer, an investigation was conducted, which revealed that extruded AH Plus does not prevent periapical healing, but can contribute to delayed healing in children.⁴² However, a new study revealed that new ceramic-based root canal sealer may be considered minimally cytotoxic, if accidentally extruded into the periapical tissues.⁴³

Conclusion

In conclusion, the current study reveals that residents in the Saudi Board of Endodontics program adopt new technologies in Endodontics, such as the use of an apex locator and NiTi rotary instrumentation; a high percentage of residents also advocate in favor of keeping canal apical patency. Most residents used zinc oxide–eugenol sealers and did not advocate sealer extrusion into the apical foramen. Further, most residents did not use adjunct devices in irrigation, but did advocate smear layer removal. In this study, there exist certain limitations, including the fact that not all residents participated in the survey, limited responses to the questionnaire, inability to verify the accuracy of answers, and the possible biases of respondents. Thus, we are in need of more investigation.

Authors' Contributions

The authors contributed equally and all authors are in agreement with this manuscript.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of Interest

None declared.

Acknowledgments

We would like to thank the Saudi Board of Endodontics residents who participated in this study. We would also like to thank Editage (www.editage.com) for English language editing.

References

- 1 Zou H, Li Y, Lian X, Yan Y, Dai X, Wang G. Frequency and influencing factors of rubber dam usage in tianjin: a questionnaire survey. *Int J Dent* 2016;2016:7383212. doi: 10.1155/2016/7383212
- 2 Lababidi EA. Discuss the impact technological advances in equipment and materials have made on the delivery and outcome of endodontic treatment. *Aust Endod J* 2013;39(3):92–97
- 3 Savani GM, Sabbah W, Sedgley CM, Whitten B. Current trends in endodontic treatment by general dental practitioners: report of a United States national survey. *J Endod* 2014;40(5):618–624
- 4 Lee M, Winkler J, Hartwell G, Stewart J, Caine R. Current trends in endodontic practice: emergency treatments and technological armamentarium. *J Endod* 2009;35(1):35–39
- 5 Clarkson RM, Podlich HM, Savage NW, Moule AJ. A survey of sodium hypochlorite use by general dental practitioners and endodontists in Australia. *Aust Dent J* 2003;48(1):20–26
- 6 Dutner J, Mines P, Anderson A. Irrigation trends among American Association of Endodontists members: a web-based survey. *J Endod* 2012;38(1):37–40
- 7 Kersten DD, Mines P, Sweet M. Use of the microscope in endodontics: results of a questionnaire. *J Endod* 2008;34(7):804–807
- 8 Locke M, Thomas MB, Dummer PM. A survey of adoption of endodontic nickel-titanium rotary instrumentation part 1: general dental practitioners in Wales. *Br Dent J* 2013;214(3):E6
- 9 Lynch CD, McConnell RJ. Attitudes and use of rubber dam by Irish general dental practitioners. *Int Endod J* 2007;40(6):427–432
- 10 Willershausen I, Wolf TG, Schmidtman I, et al. Survey of root canal irrigating solutions used in dental practices within Germany. *Int Endod J* 2015;48(7):654–660
- 11 Germack M, Sedgley CM, Sabbah W, Whitten B. Antibiotic use in 2016 by members of the American Association of Endodontists: report of a national survey. *J Endod* 2017;43(10):1615–1622
- 12 Setzer FC, Hinckley N, Kohli MR, Karabucak B. A Survey of cone-beam computed tomographic use among endodontic practitioners in the United States. *J Endod* 2017;43(5):699–704
- 13 Ha WN, Duckmanton P, Kahler B, Walsh LJ. A survey of various endodontic procedures related to mineral trioxide aggregate usage by members of the Australian Society of Endodontology. *Aust Endod J* 2016;42(3):132–138

- 14 Alrahabi M. The confidence of undergraduate dental students in Saudi Arabia in performing endodontic treatment. *Eur J Dent* 2017;11(1):17–21
- 15 Bird DC, Chambers D, Peters OA. Usage parameters of nickel-titanium rotary instruments: a survey of endodontists in the United States. *J Endod* 2009;35(9):1193–1197
- 16 Al-Shamiri HM, Alaizari NA, Al-Maweri SA, Tarakji B. Knowledge and attitude of dental trauma among dental students in Saudi Arabia. *Eur J Dent* 2015;9(4):518–522
- 17 Glickman GN, Gluskin AH, Johnson WT, Lin J. The crisis in endodontic education: current perspectives and strategies for change. *J Endod* 2005;31(4):255–261
- 18 European Society of Endodontology. Quality guidelines for endodontic treatment: consensus report of the European Society of Endodontology. *Int Endod J* 2006;39(12):921–930
- 19 Aggarwal G, Bogra P, Gupta S, Jindal A, Jain N. Determination of apical constriction and apical foramen using electronic apex locator in vivo: Comparison between vital and nonvital teeth. *Saudi Endod J* 2018;8:99–105
- 20 American Association of Endodontists, Glossary of Endodontic Terms. 9th ed. Chicago: American Association of Endodontists; 2015
- 21 Cailleteau JG, Mullaney TP. Prevalence of teaching apical patency and various instrumentation and obturation techniques in United States dental schools. *J Endod* 1997;23(6):394–396
- 22 Izu KH, Thomas SJ, Zhang P, Izu AE, Michalek S. Effectiveness of sodium hypochlorite in preventing inoculation of periapical tissues with contaminated patency files. *J Endod* 2004;30(2):92–94
- 23 Vera J, Arias A, Romero M. Dynamic movement of intracanal gas bubbles during cleaning and shaping procedures: the effect of maintaining apical patency on their presence in the middle and cervical thirds of human root canals-an in vivo study. *J Endod* 2012;38(2):200–203
- 24 Arora M, Sangwan P, Tewari S, Duhan J. Effect of maintaining apical patency on endodontic pain in posterior teeth with pulp necrosis and apical periodontitis: a randomized controlled trial. *Int Endod J* 2016;49(4):317–324
- 25 Ustun Y, Aslan T, Sagsen B, Kesim B. The effects of different nickel-titanium instruments on dentinal microcrack formations during root canal preparation. *Eur J Dent* 2015;9(1):41–46
- 26 AlRahabi M. Attitudes of general practice dentists in private dental clinics in Almadinah Almunawarah toward novel endodontic technologies. *G Ital Endod* 2016;30:10–13
- 27 Alrahabi M, Alkady A. Comparison of the shaping ability of various nickel-titanium file systems in simulated curved canals. *Saudi Endod J* 2017;7:97–101
- 28 Özyürek T. Cyclic fatigue resistance of Reciproc, WaveOne, and WaveOne Gold nickel-titanium instruments. *J Endod* 2016;42(10):1536–1539
- 29 Berutti E, Chiandussi G, Paolino DS, et al. Canal shaping with WaveOne Primary reciprocating files and ProTaper system: a comparative study. *J Endod* 2012;38(4):505–509
- 30 Plotino G, Grande NM, Cotti E, Testarelli L, Gambarini G. Blue treatment enhances cyclic fatigue resistance of vortex nickel-titanium rotary files. *J Endod* 2014;40(9):1451–1453
- 31 Blacher JD, Safavi KE, Aseltine RH, Kaufman BM. Defining endodontic residents' clinical experiences: a national survey. *JDE* 2019;83(5):504–509
- 32 Bryce G, MacBeth N, Gulabivala K, Ng YL. The efficacy of supplementary sonic irrigation using the EndoActivator system determined by removal of a collagen film from an ex vivo model. *Int Endod J* 2018;51(4):489–497
- 33 Bolles JA, He J, Svoboda KK, Schneiderman E, Glickman GN. Comparison of vibringe, endoactivator, and needle irrigation on sealer penetration in extracted human teeth. *J Endod* 2013;39(5):708–711
- 34 Morago A, Ordinola-Zapata R, Ferrer-Luque CM, Baca P, Ruiz-Linares M, Arias-Moliz MT. Influence of smear layer on the antimicrobial activity of a sodium hypochlorite/etidronic acid irrigating solution in infected dentin. *J Endod* 2016;42(11):1647–1650
- 35 Kuçi A, Alaçam T, Yavaş O, Ergul-Ulger Z, Kayaoglu G. Sealer penetration into dentinal tubules in the presence or absence of smear layer: a confocal laser scanning microscopic study. *J Endod* 2014;40(10):1627–1631
- 36 Shetty V, Hegde P, Chauhan RS, Chaurasia VR, Sharma AM, Taranath M. A spectro photometric comparative evaluation of apical sealing ability of three different sealers; calcium hydroxide based, resin based and zinc oxide eugenol based sealers. *J Int Oral Health* 2015;7(2):25–27
- 37 Singh G, Gupta I, Elshamy FM, Boreak N, Homeida HE. In vitro comparison of antibacterial properties of bioceramic-based sealer, resin-based sealer and zinc oxide eugenol based sealer and two mineral trioxide aggregates. *Eur J Dent* 2016;10(3):366–369
- 38 Wang Z, Shen Y, Haapasalo M. Dentin extends the antibacterial effect of endodontic sealers against *Enterococcus faecalis* biofilms. *J Endod* 2014;40(4):505–508
- 39 Candeiro GTM, Moura-Netto C, D'Almeida-Couto RS, et al. Cytotoxicity, genotoxicity and antibacterial effectiveness of a bioceramic endodontic sealer. *Int Endod J* 2016;49(9):858–864
- 40 Tuğ Kılış B, Er K, Taşdemir T, et al. Neurotoxicity of various root canal sealers on rat sciatic nerve: an electrophysiologic and histopathologic study. *Clin Oral Investig* 2015;19(8):2091–2100
- 41 Ricucci D, Rôças IN, Alves FR, Loghin S, Siqueira JF Jr. Apically extruded sealers: fate and influence on treatment outcome. *J Endod* 2016;42(2):243–249
- 42 Sari S, Durutürk L. Radiographic evaluation of periapical healing of permanent teeth with periapical lesions after extrusion of AH Plus sealer. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007;104(3):e54–e59
- 43 Chakar S, Changotade S, Osta N, Khalil I. Cytotoxic evaluation of a new ceramic-based root canal sealer on human fibroblasts. *Eur J Dent* 2017;11(2):141–148