When Mount Fuji Can Erupt after Seven Days: A Case Report of Delayed Posttraumatic Tension Pneumocephalus with Literature Review

V. T. T. Rajan¹ G. Krrithvi Dharini III² V. S. Sathish Anand¹ H. S. Nandish¹

¹Department of Neurosurgery, Saveetha University, Saveetha Medical College and Hospital, Chennai, Tamil Nadu, India

²Saveetha University, Saveetha Medical College and Hospital, Chennai, Tamil Nadu, India

Address for correspondence G. Krrithvi Dharini III, MBBS, Saveetha University, Saveetha Medical College and Hospital, Thandalam, Chennai-600124, Tamil Nadu, India (e-mail: krrithvidharini@gmail.com).

J Neurosci Rural Pract 2020;11:187–190

Abstract

Tension pneumocephalus (TPC) is a neurosurgical emergency that occurs when there is an expansion of trapped intracranial gas causing raised intracranial pressure. Rarely, posttraumatic TPC can occur even after 72 hours although the initial scans are normal. There are less than 20 cases of delayed TPC in the reported literature. Here, we report a case of delayed TPC that occurred 7 days after the initial injury and presented as sudden neurological deterioration. It was promptly diagnosed with a computed tomography brain and appropriate surgical intervention was performed and the outcome was good. We also did a literature review of reported cases of delayed TPC and looked out for factors that may predict its occurrence. The occurrence of an episode of cerebrospinal fluid rhinorrhea, followed by worsening of headache and sensorium in a patient with anterior cranial fossa fracture should alert a neurosurgeon to the possibility of delayed TPC.

Keywords

- traumatic tension pneumocephalus
- delayed tension pneumocephalus

Introduction

Pneumocephalus (PC) is the presence of air or gas trapped within cranial cavity. It complicates approximately 3.9 to 9.7% cases of head injury.^{1,2} Most of it remaining asymptomatic, it is uncommon for PC to morph into tension pneumocephalus (TPC). TPC is a neurosurgical emergency that occurs as a result of increased intracranial pressure due to presence of intracranial air. TPC can be acute (<72 hours) or delayed (>72 hours), with delayed being even rarer.³ Although head injury is commonly attributed to the development of TPC, postoperative status, following chronic subdural hematoma accounts for 2.5 to 16% of cases.^{4,5} Other etiological factors include infections and postventriculoperitoneal shunt, or theco-peritoneal shunt.^{3,6} We describe a case of head injury following road traffic accident, where the patient developed TPC after 7 days of head injury. This case emphasized the importance of serial radiological evaluation and neurological assessment even in mild-to-moderate head injuries.

Case Report

A 45-year-old male patient, with alleged history of road traffic accident under the influence of alcohol, was presented to the emergency department. Patient had a history of epistaxis and loss of consciousness for 10 minutes after which he recovered and had persistent mild headache. On examination, his Glasgow Coma Scale (GCS) was 15/15, decreased vision in the right eye with relative afferent pupillary defect of the right pupil. He was tested negative for other cranial nerves, sensory, or motor deficits. At the time of presentation, patient did not have cerebrospinal fluid (CSF) rhinorrhea. Computed tomography (CT) brain and facial bones (►Fig. 1) revealed multiple facial bone fractures involving inner table of frontal sinus, right zygomatic, and lateral wall of optic foramen. There was no evidence of brain parenchymal injury or PC. A postinjury 24-hour follow-up scan showed a small speck of PC in the frontal region. Magnetic resonance imaging brain with orbit revealed signal changes in the right optic nerve. Visual evoked potential suggested

Fig. 1 Computed tomography brain done on the day of admission.

for right optic neuropathy, for which the patient was started on methylprednisolone.

On postadmission day 6, he developed an episode of CSF rhinorrhea. On postadmission day 7, patient had worsening of headache with a drop in GCS of two points (13/15). Repeat CT brain (►Fig. 2) revealed bifrontal TPC with classical Mount Fuji sign. Hence, the patient was subjected to emergency burr-hole and tapping and release of TPC. During the postoperative period, patient's headache was relieved, GCS improved to 15/15, and he did not have further CSF rhinorrhea. Postsurgery scan showed resolution of TPC and the mass effect on the frontal lobes and ventricles was relieved. Patient had no further episodes of CSF rhinorrhea at 6 months of follow-up and his vision in the right eye improved significantly after a course of steroids. Currently, the patient is on regular follow-up with no significant complaints.

Discussion

Pneumocephalus was first described in 1741 by Lecat.⁷ In 1962, Ectors, Kessler, and Stern first reported a case of TPC.⁸ The two theories regarding the etiology of TPC are Dandy theory of ball valve mechanism and Horowitz theory.^{9,10}

Dandy theory states that unidirectional flow of air through a defect and trapping of air within the cavity causes

Fig. 2 Computed tomography brain done on seventh postadmission day.

TPC. Horowitz explains that TPC is due to negative pressure within the cavity as a result of excessive loss of CSF (inverted soda bottle effect). In our case, these two theories might be the most probable pathophysiology.

Etiology of PC as reported in literature shows trauma, postsurgery, neoplasms, and infections. In a large prospective study conducted in level one trauma center by Paul tran et al, the incidence of PC was 0.77% (68/8747). Trauma contributed 81% (55/68), postsurgical 13% (9/68), tumor 1.5% (1/68), and presumed IV injection or idiopathic 4% (3/68).⁷ Hence, trauma is found to be the most common etiology of PC then follows postsurgical sequel.

The incidence of TPC is very rare, limited to case reports, and small case series. Delayed TPC is characterized by PC occurring 72 hours after injury. Some cases occur even later as described by Sprague¹¹ and Poulgrain et al.¹²

The common manifestations noticed are headache, vomiting, CSF rhinorrhea, decreased vision, lethargy, and stupor progressing to coma leading to death. Diagnosis in most case was definitive with plain CT and by the characteristic Mount Fuji sign. This sign is characteristic of TPC. It appears as bilateral

frontal subdural air on CT with widened interhemispheric fissure, representing the compression and separation of frontal lobes by subdural air. TPC can also occur without classical Mount Fuji sign and in any compartment of the brain.

Review of literature for previously reported cases of delayed TPC in the past 20 years has been documented in ► **Table 1**. Search was done using keywords TPC, posttraumatic and delayed in PubMed and Google Scholar.

In this case report, fracture involving the inner table of frontal sinuses, one of the most common associations with the development of PC apart from ethmoid sinus fracture led to unilateral flow of air based on the Dandy theory, producing PC. CSF rhinorrhea further created a negative pressure resulting in PC based on the Horowitz theory.

Tension pneumocephalus always warrants surgical intervention, as in this case, urgent decompression with burr-hole/needle aspiration. If TPC is diagnosed timely with appropriate intervention, the outcome is generally very good. Challenge of delayed TPC lies in the timely diagnosis, which only can be attained by regular clinical monitoring and obtaining a plain CT brain when required.

Table 1 Review of literature for previously reported cases of delayed TPC in the past 20 years

Author	Year	Patient profile	Clinical presentation	Management	Outcome
Ruge et al	1985	28/Male 7/Male	Delayed TPC 11 days following shunt surgery Delayed TPC 1 week following shunt surgery	Surgery Surgery	Good Good
Kon et al	2003	46/Male	Delayed TPC after 7 years of craniotomy	Surgery	Good
Sankhala et al	2004	19/Male	Delayed TPC after 6 months of shunt surgery	Surgery	Good
Hong et al	2005	64/Male 38/Female	Delayed TPC after ventriculoperitoneal shunt Delayed TPC after 12 years of trauma	Surgery Surgery	Good Good
Kıymaz et al	2005	63/Male	Delayed TPC 11 days after RTA	Surgery	Good
Chandran et al	2007	18/Male	Delayed spontaneous TPC	Surgery	Good
Leong et al	2008	42/Male	Traumatic delayed TPC	Conservative	Good
Lee et al	2009	45/Male	Delayed TPC caused by spinal tapping in a patient with basal skull fracture and pneumothorax	Surgery	Good
Rathore et al	2011	30/Male	Delayed TPC 2 weeks after fall from vehicle	Surgery	Good
Solomiichuk et al	2013	75/Male	Delayed TPC 5 days after open penetrating wound	Surgery	Good
Kankane et al ²	2016	30/Male	Delayed TPC after 1 month of trauma	Surgery	Good
Wang et al ¹⁰	2016	25/Male	Delayed TPC 16 weeks after bifrontal craniotomy for self-inflicted gunshot wound	Surgery	Good
Chen et al	2018	27/Female	Delayed TPC 9 years after craniotomy	Surgery	Good

Abbreviations: RTA, road traffic accident; TPC, tension pneumocephalus.

Conclusion

Regular assessment with standardized scales such as GCS in patient with neurotrauma is essential. CT is diagnostic of TPC. A sudden worsening of headache following CSF rhinorrhea in a posttraumatic patient even with previous normal CT should raise a suspicion of TPC.

Funding

None.

Conflict of Interest

None declared.

References

- 1 Al-Aieb A, Peralta R, Ellabib M, El-Menyar A, Al-Thani H. Traumatic tension pneumocephalus: two case reports. *Int J Surg Case Rep* 2017;31:145–149
- 2 Kankane VK, Jaiswal G, Gupta TK. Posttraumatic delayed tension pneumocephalus: rare case with review of literature. *Asian J Neurosurg* 2016;11(4):343–347
- 3 Pillai P, Sharma R, MacKenzie L, et al. Traumatic tension pneumocephalus - two cases and comprehensive review of literature. *Int J Crit Illn Inj Sci* 2017;7(1):58–64
- 4 Shaikh N, Masood I, Hanssens Y, Louon A, Hafiz A. Tension pneumocephalus as complication of burr-hole drainage of chronic subdural hematoma: a case report. *Surg Neurol Int* 2010;1:27
- 5 Rathore AS, Satyarthee GD, Mahapatra AK. Post-traumatic tension pneumocephalus: series of four patients and review of literature. *Turk Neurosurg* 2016;26(2):302–305
- 6 Tran P, Reed EJ, Hahn F, Lambrecht JE, McClay JC, Omojola MF. Incidence, radiographical features, and proposed mechanism for pneumocephalus from intravenous injection of air. *West J Emerg Med* 2010;11(2):180–185
- 7 Kessler LA, Stern WZ. The ventriculopleural shunt procedure for hydrocephalus. Case report of an unusual complication. *J Pediatr* 1962;60:418–420
- 8 Dandy WE. Pneumocephalus (intracranial pneumatocele or arocele). *Arch Surg* 1969;12:949–982
- 9 Horowitz M. Intracranial pneumatocele: an unusual complication following mastoid surgery. *J Laryngol Otol* 1964;78:128–134
- 10 Wang A, Solli E, Carberry N, Hillard V, Tandon A. Delayed tension pneumocephalus following gunshot wound to the head: a case report and review of literature. *Case Rep Surg* 2016;2016:7534571