The Impact of a Stabilization Exercise on Neck Pain: A Systematic Review and Meta-analysis

Binbin Wu¹ Hongyan Yuan¹ Deyu Geng¹ Liang Zhang¹ Cheng Zhang¹

¹Department of Orthopedics, Qingdao Hospital of Traditional Chinese Medicine (Qingdao Hiser Hospital), Shandong, China

J Neurol Surg A 2020;81:342–347.

Address for correspondence Dr. Cheng Zhang, Department of Orthopedics, Qingdao Hospital of Traditional Chinese Medicine (Qingdao Hiser Hospital), 4 Renmin Road, Qingdao, Shandong, 325000, China (e-mail: zangcheng100@163.com).

Abstract

Introduction The efficacy of a stabilization exercise for the relief of neck pain remains controversial. We conducted a systematic review and meta-analysis to explore the effectiveness of a stabilization exercise on neck pain.

Methods We searched Embase, Web of Science, EBSCO Information Services, and the Cochrane Library databases through May 2019 for randomized controlled trials (RCTs) assessing the impact of a stabilization exercise on neck pain. This meta-analysis was performed using the random effects model.

Results Six RCTs are included in the meta-analysis. Compared with the control group of patients with neck pain, a stabilization exercise can significantly reduce pain scores at 4 to 6 weeks (mean difference [MD]: -2.41 ; 95% confidence interval [CI], -4.46 to -0.35 ; $p = 0.02$), Neck Disability Index [NDI] at 10 to 12 weeks (MD: -6.75 ; 95% CI, -11.71 to -1.79 ; $p = 0.008$), and depression scale at 4 to 6 weeks (MD: -4.65 ; 95% CI, -7.00 to -2.31 ; $p = 0.02$), but it has no obvious impact on pain scores at 10 to 12 weeks (MD: -1.07 ; 95% CI, -3.42 to 1.28 ; $p = 0.37$) or at 6 months (MD: -1.02 ; 95% CI, -3.43 to 1.39 ; $p = 0.41$).

Conclusions A stabilization exercise can provide some benefits to control neck pain.

Keywords

- ▶ stabilization exercise
- ▶ neck pain
- ▶ randomized controlled trials

Introduction

Neck pain is reported to rank as the second most prevalent musculoskeletal disorder.^{1–3} It is estimated that ~ 67% of people may experience neck pain during their lifetime.^{4,5} Neck pain results in a reduced quality of daily life and has a considerable economic impact on health care systems. Its diagnosis and treatment are still challenging because the anatomical source and cause of neck pain remains elusive in most cases.⁶

Neck pain may be associated with a reduction in the strength and endurance capacity of the cervical muscles.^{7,8} For instance, the strength of deep and anterior cervical flexors in the cervical spine is reduced in patients with neck pain.⁹ Exercise can possibly alleviate neck pain because of its ability to gain muscle strength, flexibility, and endurance,

as well as restore injured tissues.¹⁰ Stabilization exercises serve as an important approach to alleviate back and pelvic pain,^{11–13} and they have the ability to achieve a stable injury-free state for the cervical spine.^{14,15}

Several studies reported on the treatment efficacy of stabilization exercises on neck pain, but the results are conflicting.^{4,16–18} This meta-analysis of randomized controlled trials (RCTs) aimed to investigate the treatment efficacy and function of stabilization exercises versus general exercises for patients with neck pain.

Materials and Methods

This systematic review and meta-analysis were performed based on the guidance of the Preferred Reporting Items for Systematic Reviews and Meta-analysis statement and

received

March 12, 2019

accepted after revision

March 12, 2019

published online

March 6, 2020

© 2020 Georg Thieme Verlag KG
Stuttgart · New York

DOI <https://doi.org/10.1055/s-0039-3400953>.
ISSN 2193-6315.

Cochrane Handbook for Systematic Reviews of Interventions.^{19,20} No ethical approval and patient consent were required because all analyses were based on previously published studies.

Literature Search and Selection Criteria

We systematically searched several databases including Embase, Web of Science, EBSCO Information Services, and the Cochrane Library from inception to May 2019 with the key words *stabilization exercise* and *neck pain*. The reference lists of retrieved studies and relevant reviews were also hand-searched, and the search process as described was performed repeatedly to include additional eligible studies.

The inclusion criteria are presented as follows: (1) study design is RCT, (2) patients are diagnosed with neck pain, and (3) intervention treatments are stabilization exercises versus general exercise. Patients with cervical radiculopathy were excluded.

Data Extraction and Outcome Measures

Some baseline information was extracted from the original studies including first author, number of patients, age, female, body mass index (BMI), and detailed methods in the two groups. Data were extracted independently by two investigators, and discrepancies were resolved by consensus. We contacted the corresponding author to obtain data when necessary.

The primary outcomes are pain scores at 4 to 6 weeks and 10 to 12 weeks. Secondary outcomes are pain scores at 6 months, Neck Disability Index (NDI) at 10 to 12 weeks, and depression scale at 4 to 6 weeks.

Quality Assessment in Individual Studies

The methodological quality of each RCT was assessed by the Jadad Scale that consists of three evaluation elements: randomization (0–2 points), blinding (0–2 points), and drop-outs and withdrawals (0–1 points).²¹ One point is allocated to each element if they have been conducted and mentioned appropriately in the original article. The Jadad Scale scores vary from 0 to 5 points. An article with a Jadad score ≤ 2 is considered low quality; it is considered high quality with a Jadad score ≥ 3 .²²

Statistical Analysis

We assessed mean differences (MDs) with 95% confidence intervals (CIs) for continuous outcomes (pain scores at 4–6 weeks, 10–12 weeks, and 6 months; NDI at 10–12 weeks; and depression scale at 4–6 weeks). Heterogeneity was evaluated using the I^2 statistic with $I^2 > 50\%$ indicating significant heterogeneity.²³ The random effects model was used for all meta-analyses. We searched for potential sources of heterogeneity for significant heterogeneity. Sensitivity analysis was performed to detect the influence of a single study on the overall estimate via omitting one study in turn or performing the subgroup analysis. Owing to the limited number (< 10) of the included studies, publication bias was not assessed. Results were considered statistically significant for $p < 0.05$. All statistical analyses were performed using RevMan, v.5.3 (The Cochrane Collaboration, Software Update, Oxford, United Kingdom).

Results

Literature Search, Study Characteristics, and Quality Assessment

A detailed flowchart of the search and selection results is shown in ►Fig. 1. A total of 469 potentially relevant articles were identified initially. Six RCTs that ultimately met our inclusion criteria were included in the meta-analysis.^{4,16–18,24,25}

►Table 1 presents the main characteristics of the six included RCTs. The six studies were published between 2003 and 2019, and the total sample size was 395. Five included RCTs involved chronic neck pain,^{4,16,18,24,25} and the remaining RCT involved neck pain caused by posttraumatic stress disorder.¹⁷

Among the six RCTs, four included RCTs that reported pain scores at 4 to 6 weeks,^{16–18,25} two included RCTs that reported pain scores at 10 to 12 weeks,^{4,25} two included RCTs that reported pain scores at 6 months,^{18,25} two included RCTs that reported the NDI at 10 to 12 weeks,^{4,25} and two included RCTs that reported the depression scale at 4 to 6 weeks.^{17,25} Jadad scores of the six included studies varied from 3 to 5 points, and all six studies were considered high quality, according to the quality assessment.

Primary Outcomes: Pain Scores at 4 to 6 Weeks and 10 to 12 Weeks

The random effects model was used for the analysis of pain scores at 4 to 6 weeks and 10 to 12 weeks. Compared with the control group of patients with neck pain, stabilization exercises were associated with significantly reduced pain scores at 4 to 6 weeks (MD: -2.41 ; 95% CI, -4.46 to -0.35 ; $p = 0.02$) with significant heterogeneity among the studies (I^2 : 94%; heterogeneity $p < 0.00001$; ►Fig. 2) but showed no notable impact on pain scores at 10 to 12 weeks (MD: -1.07 ; 95% CI, -3.42 to 1.28 ; $p = 0.37$) with significant heterogeneity among the studies (I^2 : 91%; heterogeneity $p = 0.0009$; ►Fig. 3).

Sensitivity Analysis

Significant heterogeneity was observed among the included studies for the primary outcomes, and significant heterogeneity remained when performing sensitivity analysis by omitting one study in turn.

Secondary Outcomes

Stabilization exercises were found to have no substantial influence on pain scores at 6 months compared with the control group for neck pain (MD: -1.02 ; 95% CI, -3.43 to 1.39 ; $p = 0.41$; ►Fig. 4), but did result in a reduction in NDI at 10 to 12 weeks (MD: -6.75 ; 95% CI, -11.71 to -1.79 ; $p = 0.008$; ►Fig. 5) and depression scale at 4 to 6 weeks (MD: -4.65 ; 95% CI, -7.00 to -2.31 ; $p = 0.02$; ►Fig. 6).

Discussion

Altered motor control is confirmed in chronic neck pain.^{6,26,27} Different aspects are associated with altered motor control and include co-contraction of agonistic muscles, more activity for superficial flexors and extensors,

Fig. 1 Flow diagram of study search and selection process.

delayed onset time for neck muscles, and weakness of deep muscles.^{6,28–30} The muscles are found to provide ~ 80% of spinal stabilization.³¹ Deep and superficial muscles have different roles for cervical stabilization. Deep muscles are segmental stabilizers and provide stabilization between segments that is the basis for participation of the superficial muscles.⁶

Patients with neck pain have weak deep cervical flexors that may result in increased activity level of the superficial flexor muscles.⁶ These patients have reduced deep muscle activity and increased superficial muscle activity during cognitive tasks and functional activities. Increased activity of superficial flexors during isometric contraction is also reported to cause neck pain.²⁸ High co-contraction of a weak sternocleidomastoid and the anterior scalene can lead to neck pain and disability.³² A specific exercise program should be designed to target the deep flexors first and then the superficial ones.

Cervical stabilization exercises can improve neck pain and cervical muscle performance in patients with cervicogenic headache.¹² When compared with isometric and stretching

exercises, stabilization exercises may be more effective in improving disability and pain control for patients with neck pain.²⁵ Our meta-analysis concluded that stabilization exercises can substantially reduce pain scores at 4 to 6 weeks, NDI at 10 to 12 weeks, and depression scale at 4 to 6 weeks for neck pain patients, but they show no obvious influence on pain scores at 10 to 12 weeks or 6 months. In one included RCT involving the influence of exercise programs on neck pain, the results found significantly increased deep flexor muscle endurance for stabilization exercise and also increased endurance for a group of routine exercises.⁴

Significant heterogeneity is observed when omitting one study in turn for the sensitivity analysis. Several factors may account for this heterogeneity, and they include different causes (e.g., chronic neck pain and posttraumatic stress disorder) of neck pain, various combinations with stabilization exercises (e.g., isometric neck strengthening and physical therapy), and duration of administration (ranging from 6 weeks to 12 months).

This study has several possible limitations. First, only six RCTs were included in this meta-analysis, and five of them

Table 1 Characteristics of included studies

No.	Study	Stabilization exercise group						Control group						Follow-up time	Jada score
		N	Age, y	Female, n	BMI, kg/m ²	Duration, mo	Methods	N	Age, y	Female, n	BMI, kg/m ²	Duration, mo	Methods		
1	Shiravi et al ¹⁶	45	27.6 ± 2.06	–	22.2 ± 1.16	–	Scapular stabilization exercises for neck movement impairment	45	25.11 ± 1.99	–	22.3 ± 1.43	–	Active exercise performed by the patient	6 wk	4
2	Ghaderi et al ⁴	20	35.97 ± 2.5	–	–	–	Stabilization exercise and routine electrotherapy for nonspecific chronic neck pain	20	36.346 ± 3.06	–	–	–	Routine exercise and electrotherapy	10 wk	3
3	Park and Kim ¹⁷	15	57.5 ± 6.7	2	23.3 ± 3.5	24.6 ± 12.1	Cervical exercises for 30 min, 3 times/week for 6 wk and physical therapy for posttraumatic stress disorder	16	62.86 ± 7.9	4	23.76 ± 3.3	20.2 ± 10.1	Physical therapy	6 wk	4
4	Griffiths et al ¹⁸	37	51.1 ± 14.00	20	–	30 (median)	Specific neck stabilization exercises with general neck advice and exercise program for chronic neck disorders	37	51.5 ± 13.6	26	–	24 (median)	General neck advice and exercise program	6 mo	5
5	Dusunceli et al ²⁵	20	50.2 ± 4.8	14	–	45.0 ± 46.8	Physical therapy agents and neck stabilization exercises for nonspecific neck pain	20	53.4 ± 6.8	12	–	43.2 ± 40.6	Physical therapy agents	12 mo	4
6	Ylinen et al ⁶	60	45 ± 6	–	25 ± 3	96 ± 72	High-intensity isometric neck strengthening and stabilization exercises for chronic neck pain	60	46 ± 5	–	26 ± 4	96 ± 60	General exercises	12 mo	4

Abbreviation: BMI, body mass index.

Fig. 2 Forest plot for meta-analysis of pain scores at 4 to 6 weeks. CI, confidence interval; IV, inverse variance; SD, standard deviation.

Fig. 3 Forest plot for meta-analysis of pain scores at 10 to 12 weeks. CI, confidence interval; IV, inverse variance; SD, standard deviation.

Fig. 4 Forest plot for meta-analysis of pain scores at 6 months. CI, confidence interval; IV, inverse variance; SD, standard deviation.

Fig. 5 Forest plot for meta-analysis of Neck Disability Index (NDI) at 10 to 12 weeks. CI, confidence interval; IV, inverse variance; SD, standard deviation.

Fig. 6 Forest plot for meta-analysis of depression scale at 4 to 6 weeks. CI, confidence interval; IV, inverse variance; SD, standard deviation.

had a small sample size ($n < 100$). Overestimation of the treatment effect is more likely in smaller trials compared with larger samples. Next, significant heterogeneity was observed in this meta-analysis and may have been caused by different methods and duration of stabilization exercises, as well as sex. Finally, it was not feasible to perform a meta-analysis of some important outcomes such as endurance or the SF-36 Short Form. In conclusion, stabilization exercises can provide some benefit for pain relief and cervical function for patients with neck pain.

Conflicts of Interest

None declared.

References

- 1 Côté P, Cassidy JD, Carroll L. The Saskatchewan Health and Back Pain Survey. The prevalence of neck pain and related disability in Saskatchewan adults. *Spine* 1998;23(15):1689-1698
- 2 Genebra CVDS, Maciel NM, Bento TPF, Simeão SFAP, Vitta A. Prevalence and factors associated with neck pain: a population-based study. *Braz J Phys Ther* 2017;21(04):274-280

- 3 Ye S, Jing Q, Wei C, Lu J. Risk factors of non-specific neck pain and low back pain in computer-using office workers in China: a cross-sectional study. *BMJ Open* 2017;7(04):e014914
- 4 Ghaderi F, Jafarabadi MA, Javanshir K. The clinical and EMG assessment of the effects of stabilization exercise on nonspecific chronic neck pain: a randomized controlled trial. *J Back Musculoskeletal Rehabil* 2017;30(02):211–219
- 5 Fougere N, Fleiter B. Temporomandibular disorder and comorbid neck pain: facts and hypotheses regarding pain-induced and rehabilitation-induced motor activity changes. *Can J Physiol Pharmacol* 2018;96(11):1051–1059
- 6 Falla D. Unravelling the complexity of muscle impairment in chronic neck pain. *Man Ther* 2004;9(03):125–133
- 7 Chiu TT, Sing KL. Evaluation of cervical range of motion and isometric neck muscle strength: reliability and validity. *Clin Rehabil* 2002;16(08):851–858
- 8 Ylinen J, Salo P, Nykänen M, Kautiainen H, Häkkinen A. Decreased isometric neck strength in women with chronic neck pain and the repeatability of neck strength measurements. *Arch Phys Med Rehabil* 2004;85(08):1303–1308
- 9 Gogia PP, Sabbahi MA. Electromyographic analysis of neck muscle fatigue in patients with osteoarthritis of the cervical spine. *Spine* 1994;19(05):502–506
- 10 Wolsko PM, Eisenberg DM, Davis RB, Kessler R, Phillips RS. Patterns and perceptions of care for treatment of back and neck pain: results of a national survey. *Spine* 2003;28(03):292–297; discussion 298
- 11 Kose G, Hepguler S, Atamaz F, Oder G. A comparison of four disability scales for Turkish patients with neck pain. *J Rehabil Med* 2007;39(05):358–362
- 12 Jull G, Trott P, Potter H, et al. A randomized controlled trial of exercise and manipulative therapy for cervicogenic headache. *Spine* 2002;27(17):1835–1843; discussion 1843
- 13 Jordan A, Mehlsen J, Ostergaard K. A comparison of physical characteristics between patients seeking treatment for neck pain and age-matched healthy people. *J Manipulative Physiol Ther* 1997;20(07):468–475
- 14 Monticone M, Barbarino A, Testi C, Arzano S, Moschi A, Negrini S. Symptomatic efficacy of stabilizing treatment versus laser therapy for sub-acute low back pain with positive tests for sacroiliac dysfunction: a randomised clinical controlled trial with 1 year follow-up. *Eura Medicophys* 2004;40(04):263–268
- 15 Arokoski JP, Valta T, Airaksinen O, Kankaanpää M. Back and abdominal muscle function during stabilization exercises. *Arch Phys Med Rehabil* 2001;82(08):1089–1098
- 16 Shiravi S, Letafatkar A, Bertozzi L, Pillastrini P, Khaleghi Tazji M. Efficacy of abdominal control feedback and scapula stabilization exercises in participants with forward head, round shoulder postures and neck movement impairment. *Sports Health* 2019; 11(03):272–279
- 17 Park SD, Kim SY. Clinical feasibility of cervical exercise to improve neck pain, body function, and psychosocial factors in patients with post-traumatic stress disorder: a randomized controlled trial. *J Phys Ther Sci* 2015;27(05):1369–1372
- 18 Griffiths C, Dziedzic K, Waterfield J, Sim J. Effectiveness of specific neck stabilization exercises or a general neck exercise program for chronic neck disorders: a randomized controlled trial. *J Rheumatol* 2009;36(02):390–397
- 19 Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009;339:b2535
- 20 Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions* [version 5.1.0]. Chichester, UK: John Wiley; 2011
- 21 Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17(01):1–12
- 22 Kjaergard LL, Villumsen J, Gluud C. Reported methodologic quality and discrepancies between large and small randomized trials in meta-analyses. *Ann Intern Med* 2001;135(11):982–989
- 23 Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539–1558
- 24 Ylinen J, Takala EP, Nykänen M, et al. Active neck muscle training in the treatment of chronic neck pain in women: a randomized controlled trial. *JAMA* 2003;289(19):2509–2516
- 25 Dusunceli Y, Ozturk C, Atamaz F, Hepguler S, Durmaz B. Efficacy of neck stabilization exercises for neck pain: a randomized controlled study. *J Rehabil Med* 2009;41(08):626–631
- 26 Woodhouse A, Vasseljen O. Altered motor control patterns in whiplash and chronic neck pain. *BMC Musculoskelet Disord* 2008; 9:90
- 27 De Pauw R, Coppieters I, Caeyenberghs K, et al. Associations between brain morphology and motor performance in chronic neck pain: a whole-brain surface-based morphometry approach. *Hum Brain Mapp* 2019
- 28 Falla D, Jull G, Hodges PW. Feedforward activity of the cervical flexor muscles during voluntary arm movements is delayed in chronic neck pain. *Exp Brain Res* 2004;157(01):43–48
- 29 Falla D, Lindstrøm R, Rechter L, Farina D. Effect of pain on the modulation in discharge rate of sternocleidomastoid motor units with force direction. *Clin Neurophysiol* 2010;121(05):744–753
- 30 Falla D, Rainoldi A, Merletti R, Jull G. Spatio-temporal evaluation of neck muscle activation during postural perturbations in healthy subjects. *J Electromyogr Kinesiol* 2004;14(04):463–474
- 31 Hides JA, Richardson CA, Jull GA. Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine* 1996;21(23):2763–2769
- 32 Lindstrøm R, Schomacher J, Farina D, Rechter L, Falla D. Association between neck muscle coactivation, pain, and strength in women with neck pain. *Man Ther* 2011;16(01):80–86