Aortic Coarctation

Umberto G. Rossi, MD, EBIR^{1,2} Anna Maria Ierardi, MD² Gianpaolo Carrafiello, MD, Prof³
Maurizio Cariati, MD²

¹Department of Diagnostic Imaging, Interventional Radiology Unit, Ente Ospedaliero Galliera Hospital, Genova, Italy

²Department of Diagnostic and Therapeutic Advanced Technology, Diagnostic and Interventional Radiology Unit, Azienda Socio Sanitaria Territoriale Santi Paolo and Carlo Hospital, Milano, Italy

³Department of Services and Preventive Medicine, Radiology Unit, Ca' Granda Foundation, Maggiore Policlinic Hospital, University of Milan, Milano, Italy

Address for correspondence Umberto G. Rossi, MD, EBIR, Department of Diagnostic Imaging, Interventional Radiology Unit, Ente Ospedaliero Galliera Hospital, Mura delle Cappuccine, 14-16128 Genova, Italy
(e-mail: urossi76@hotmail.com).

AORTA 2020;8:46–47.

Abstract

Keywords

- aorta
- coarctation
- vascular pathology
- computed tomography
- imaging
- aging

We report a case of a 45-year-old male suffering from arterial hypertension who was found to have an aortic coarctation with marked hypertrophic compensatory collateral arterial circulation. Although coarctation is relatively rare, this must be included in the differential diagnosis in patients with arterial hypertension with a positive gradient between upper and lower limbs.

A 45-year-old male was evaluated for arterial hypertension. A pressure gradient (18 mm Hg) between upper and lower limbs was noted. Among the various diagnostic examinations, the patient underwent multidetector computed tomography (MD-CT) angiography. This showed an aortic coarctation at the level of isthmus proximal third of thoracic aorta with significant hypertrophic compensatory collateral arterial circulation (► **Fig. 1**). At the time of diagnosis, endovascular or open surgery treatment were offered to the patient, but he refused all intervention. Therefore, the patient was started on life-long antihypertensive medical treatment with clinical and imaging follow-up.

Aortic coarctation in adults is generally recognized via systemic arterial hypertension associated with a pressure

gradient between the upper and lower extremities.¹ Precise vascular imaging (MD-CT or magnetic resonance imaging) is mandatory for a complete evaluation of the thoracic aorta, its branches, and possible collateral vessels.^{2,3} Criteria for invasive treatment in adult patients include translesional pressure gradient (>20 mm Hg) and/or evidence of significant collateral vessels.^{1,4–6} Choice between open surgery versus percutaneous endovascular treatment should be determined by a multidisciplinary team specialists (surgeons, interventional radiologists, and cardiologists).^{1,7} Finally, all patients affected by aortic coarctation require a life-long treatment of arterial pressure and close follow-up (clinical and imaging).

received
September 14, 2018
accepted after revision
December 18, 2019

DOI <https://doi.org/10.1055/s-0040-1701522>.
ISSN 2325-4637.

Copyright © 2020 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 760-0888.

License terms

Fig. 1 Multidetector computed tomography angiography sagittal reconstruction demonstrating the presence of aortic coarctation at the level of isthmus proximal third of thoracic aorta (arrowhead) with hypertrophic compensatory collateral arterial circulation from the intercostal and mammary arteries (arrows).

Funding

This research was funded solely through institutional sources.

Conflict of Interest

The authors do not report any conflict of interest related to this article.

Acknowledgment

None.

References

- 1 Warnes CA, Williams RG, Bashore TM, et al. ACC/AHA 2008 guidelines for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (writing committee to develop guidelines on the management of adults with congenital heart disease). *Circulation* 2008;118(23):e714–e833
- 2 Ferro C, Rossi UG, Seitun S, Scarano F, Passerone G, Williams DM. Aortic branch artery pseudoaneurysms associated with intramural hematoma: when and how to do endovascular embolization. *Cardiovasc Intervent Radiol* 2013;36(02):422–432
- 3 Rossi UG, Seitun S, Cariati M. Aortic dissection: the flood tide sign. *J Cardiovasc Comput Tomogr* 2013;7(05):330–332
- 4 Ruys TP, Bekkers JA, Duvekot JJ, Roos-Hesselink JW. A pregnant patient with native aortic coarctation and aneurysm. *Aorta (Stamford)* 2014;2(03):110–112
- 5 Inaraja Pérez GC, Spitzer Cano E, Angulo Hervías E, López Ramón M, Salazar González JJ, Merchante García P. Images in vascular medicine. Asymptomatic aortic coarctation diagnosed because of a vast calcified collateral circulation. *Vasc Med* 2014;19(01):75–76
- 6 Beckmann E, Jassar AS. Coarctation repair-redo challenges in the adults: what to do? *J Vis Surg* 2018;4:76–88
- 7 Rossi UG, Santuari D, Dallatana R, Cariati M. Use of iliac branch device for endovascular treatment for abdominal aorta aneurysm with small diameter neck. *Aorta (Stamford)* 2017;5(06):181–183