Original Article

A SYSTEMATIC EVALUATION OF RAPID DOT-EIA, BLOOD CULTURE AND WIDAL TEST IN THE DIAGNOSIS OF TYPHOID FEVER

Sanjeev H¹, Sweettha Nayak², Pai Asha K.B.³, Rai Rekha⁴, Vimal Karnaker⁵ & Ganesh H.R.²

^{1,3}Assistant Professors, ²Tutors, ⁴Associate Professor, ⁵Professor & Head,

Department of Microbiology, K.S. Hegde Medical Academy, Nitte University, Mangalore - 575 018. India.

Correspondence:

Sanjeev H.

Assistant Professor

Department of Microbiology, K.S. Hegde Medical Academy, Nitte University, Mangalore - 575 018, Karnataka, India

Phone: +91 824 2204490/92. Mobile : +91 99722 12280 E-mail : drsanjeevh@gmail.com

Abstract :

Background : Typhoid fever, caused by *Salmonella enterica* serotype Typhi, is endemic in the Indian sub-continent including Bangladesh, South-east and Far-east Asia, Africa and South Central America. The disease can occur in all age group with highest incidence among children. Blood culture is regarded as the gold standard for diagnosis and carry 70-75% diagnostic yield in the first week of illness. However, this requires laboratory equipment and technical training that are beyond the means of most primary health care facilities in the developing world. Typhidot is a rapid dot-enzyme immune assay (EIA), which detects IgG and IgM antibodies to a specific 50 kD outer membrane protein (OMP) antigen of *Salmonella enterica* serotype Typhi. Typhidot becomes positive as early as in the first week of fever. The results can be visually interpreted and is available within one hour.

Materials and method : Fifty blood samples, collected aseptically from patients clinically diagnosed of Typhoid fever, were evaluated by blood culture, Widal test and Typhidot.

Results : Of the 50 patients, 33 (66%) were positive by blood culture. Widal test was positive in 33(66%) patients which included 26 in blood culture positive patients and 7 in blood culture negative patients. Typhidot was positive in 37 (74%) patients. Thus, in comparison to the gold standard test i.e. blood culture, Typhidot and Widal test had sensitivity and specificity of 100% & 76% and 78.78% & 58.82% respectively.

Conclusion : Typhidot is found to have high sensitivity and good specificity and could be applied as a good alternate in resource poor nation. Further, it is simple to perform, reliable when compared to Widal test, and rapid, with results being available in one hour when compared to 48 hours for blood culture and 18 hours for Widal test.

Key words : Typhidot, Blood culture, Widal, *Salmonella enterica*

Introduction :

Typhoid fever, caused by *Salmonella enterica* serotype Typhi, is a major cause of morbidity and mortality world-wide, causing an estimated 16.6 million new infections and 600,000 deaths each year.¹ It is endemic in the Indian sub-continent including Bangladesh, South-east and Far-east Asia, Africa and South Central America. The disease can

occur in all age group with highest incidence among children.² The annual incidence of typhoid fever has been reported as more than 13 million cases in Asia.³

Isolation of *Salmonella* from blood, urine or stool is the most reliable means of confirming an infection. Blood culture is regarded as the gold standard for diagnosis and carry 70-75% diagnostic yield in the first week of illness.⁴ However, this requires laboratory equipment and technical training that are beyond the means of most primary health care facilities in the developing world.¹ In addition, easy availability and widespread use of antibiotics in the community makes it frequently difficult to isolate the organism from blood culture and alternate methods such as bone marrow cultures may be required, which is invasive and difficult to carry out.²

Thus one has to rely on serological diagnosis, which is the mainstay of diagnosis of typhoid fever in most laboratories

Access this article online

Quick Response Code

in developing world.⁵ Unfortunately, neither the Widal test, which remains in widespread use in the developing world, nor any of the serodiagnostic tests that have since been developed has proven sufficiently sensitive, specific and practical to be of value in areas where this disease is endemic.¹

Typhidot is a rapid dot-enzyme immune assay (EIA), which detects IgG and IgM antibodies to a specific 50 kD outer membrane protein (OMP) antigen of *Salmonella enterica* serotype Typhi. Typhidot becomes positive as early as in the first week of fever, the results can be visually interpreted and is available within one hour.⁶

This study was undertaken to systematically evaluate the utility of Typhidot in diagnosis of typhoid fever in terms of sensitivity and specificity.

Materials and methods:

The study was conducted in the Department of Microbiology, K.S. Hegde Medical College and Hospital, located in Mangalore. Blood was collected from 50 patients who were clinically diagnosed with typhoid fever.

Ten ml of blood was inoculated into blood culture media (BHI broth) and incubated at 37°C. Subcultures were done on every alternate day till the 7th day. The growth of *Salmonella* was identified as per standard protocol and confirmed by agglutination with *Salmonella* polyvalent "O", "O"9 and H:d' antisera.⁷ The Widal test was performed by tube agglutination method and it was considered positive when a titre of equal to or more than 1:160 was observed.⁸ Typhidot is a rapid, qualitative test which detects specific IgG and IgM antibodies to a specific 50-kD OMP antigen on *Salmonella typhi enterica* serotype Typhi strains. The test was done as per manufacturer's kit instructions (Typhidot; Malaysian Bidiagnostic Research SDN BHD, Kuala Lumpur, Malaysia). Reaction tray was divided into two columns and marked "M" and "G" and strips were placed respectively. 250µl of the sample diluents was then added to reaction wells. 2.5µl of control/test serum was added to each well and incubated for 20 minutes at room temperature on a rocker platform.

The strip was washed thrice and 250µl Anti human IgM and Anti human IgG conjugates were added to respective wells and incubated for 15 minutes. The strip washed as before, and 250µl of colour development solution was added and incubated for another 15 min. The reaction was stopped, the strip was washed in distilled water and the result was read. When both the dots on the test strip were as dark as or darker than their corresponding dots on the positive control strip, they were reported as positive. The absence of visible spot indicated a negative test result. If the spots were fainter than control, that sample was also considered negative, as per directions given in the kit. In case of discrepant appearance of duplicate spots the test was repeated and only if both spots were darker than control, the sample was considered positive.

Results:

Fifty blood samples were evaluated for typhoid fever by blood culture, Widal test and Typhidot. Of the 50 patients, 33 (66%) were positive by blood culture. *Salmonella enterica* serotype Typhi was isolated in 30 patients where as *Salmonella enterica* serotype Paratyphi A was isolated in 3 patients. Widal test was positive in 33(66%) patients which included 26 in blood culture positive patients and 7 in blood culture negative patients. Typhidot was positive in 37 (74%) patients. Of the 37 Typhidot positive cases 30 were positive for both IgG and IgM antibodies, 4 for IgG antibodies only and 3 for IgM antibodies only. Of the 37 cases positive by Typhidot, 33 were also positive by blood culture and 4 were negative by blood culture. These 4 patients who were negative on blood culture but positive by Typhidot were also positive by Widal test. Thus, in comparison to the gold standard test i.e blood culture, Typhidot and Widal test had sensitivity and specificity of 100% & 76% and 78.78% & 58.82% respectively.(Table 1, Table 2). A comparison of Widal and Typhidot is described in Table 3.

Discussion:

Typhoid fever is a systemic illness with a significant morbidity and mortality in developing countries.⁹ Emergence of multidrug resistant strains of *Salmonella*

enterica serotype Typhi has only added to the burden of the disease. Any delay in diagnosis and institution of appropriate therapy only increases the risk of outcome.²

Blood culture has remained the gold standard for diagnosis of typhoid fever. Of the 50 patients clinical diagnosis of typhoid fever evaluated, 66% (33/50) were positive by blood culture. Of the 33 positive blood cultures, 30 were *Salmonella enterica* serotype Typhi and 3 were *Salmonella enterica* serotype Paratyphi A. This rate of isolation for blood culture correlates well documented data, which reports sensitivity of 65-68%.⁹ The high rate of isolation in the present study could be attributed to a) awareness among people leading to early presentation to secondary or tertiary health care facility, b) the present study involved adult population thus adequate quantities of blood could be taken for diagnostic test and c) improvements in blood culture methods especially use of automated systems. While, in contrast to the present study, many reports have quoted sensitivity ranging from 8.9-43%.^{5, 10-14} The relative low rate of isolation from blood culture had been attributed to delay in diagnosis, widespread and irrational use of antibiotics and low volume of blood obtained for cultures especially among children.

Widal test has been used for over a century in developing countries for diagnosing typhoid fever but it has been reported to have low sensitivity, specificity and positive predictive value.⁹ In the present study, Widal test was positive in 66% (33/50) of the patients. Widal test was positive in 26 of the 33 blood culture positive patients and 7 of blood culture negative patients. Thus the test had sensitivity of 78.78% and specificity of 58.82%. This corroborates well with data from endemic areas.^{5, 9, 15} The finding of the present study indicate a low specificity for Widal test. This is very much in consonance with published data especially from endemic areas where there may be high levels of specific and cross reacting antibodies. Though higher sensitivity and specificity for Widal has been reported, its use in endemic areas should not be encouraged. Ideally a fourfold rise in antibody titre in a paired serum is more diagnostic.¹⁶

Typhidot is a new, inexpensive and reliable serodiagnostic test available commercially and have been reported to have good sensitivity and specificity.⁹ In the present study, the sensitivity and specificity of Typhidot was 100% and 76.5% respectively. The sensitivity is very much in agreement with other studies, while higher specificities have been reported¹⁷. A specificity of 76.5% is comparable to studies from Indian sub-continent who have reported specificity of 77-87%.^{9, 18} Typhidot was positive in all blood culture positive patients and in 4 blood culture negative patients. Interestingly, these 4 patients were also positive by Widal test. This finding could be attributed to a) endemicity of typhoid fever in the region indicating pre-existing antibody from previous exposure or may be due to sub clinical infection, and this is supported by the fact that these patients were positive for IgG antibodies only, and b) that the patient may have had non-typhoidal fever. All the 50 patients in the study group had strong clinical suspicion of Enteric fever, were treated for the same and recovered completely without requiring any changes in the prescribed antibiotics. Thus, it could be argued that, though Typhidot gave false positive result in 4 cases, these patients may actually have had *Salmonella* infection, either in the past or sub clinical, which the blood culture failed to detect.

Conclusion :

It can be concluded that though blood culture is still the gold standard, Typhidot is found to have high sensitivity and good specificity and could be applied as a good alternate in resource poor nation. Further, it is simple to perform, reliable when compared to Widal test and rapid, with results being available in one hour when compared to 48 hours for blood culture and 18 hours for Widal test.

Table 1-Comparison of blood culture with Widal test

Widal	Blood culture			
		Positive	Negative	Total
	Positive	26	07	33
	Negative	07	10	17
	Total	33	17	50

Sensitivity-78.78%

Specificity-58.82%

Table 2- Comparison of blood culture with Typhidot

Typhidot	Blood culture			
		Positive	Negative	Total
	Positive	33	04	37
	Negative	00	13	13
	Total	33	17	50

Sensitivity-100%

Specificity-76.58%

Table 3- Comparison of Widal test and Typhidot.

Widal	Typhidot			
		Positive	Negative	Total
	Positive	30	03	33
	Negative	07	10	17
	Total	37	13	50

References :

- Olsen SJ, Pruckler J, Bibb W, Thi My Thanh N, MyTrinh T, ThiMinh N et al. Evaluation of Rapid Diagnostic Tests for Typhoid fever. *J Clin Microbial* 2004; 42:1885-1889.
- Begum Z, Hossain Md A, Shamsuzzaman AKM, Ahsan Md M, Musa AKM, Mahmud Md C et al. Evalutaion of Typhidot(IgM) for Early Diagnosis of Typhoid fever. *Bangladesh J Med Microbiol* 2009; 3:10-13.
- Ivanoff B, Levine MM, Lambert PH. Vaccination against typhoid fever, present status. *Bull WHO* 1994;72(6).957-71.
- Krishna S, Desai S, Anjana VK, Paranthaaman RG. Typhidot(IgM) as a reliable and rapid diagnostic test for typhoid fever. *Ann Trop Med Pub Health* 2011; 4:42-44.
- Narayanappa D, Sripathi R, Kumar JK, Rajani HS. Comparative study of Dot Enzyme Immunoassay (Typhidot-M) and Widal test in the diagnosis of Typhoid fever. *Indian Pediatrics* 2009; 47: 331-333
- Choo KE, Openheimer SJ, Ismail AB, Ong KH. Rapid serodiagnosis of typhoid fever by dot enzyme immunoassay in an endemic area. *Clin Infect Dis* 1994;19: 172-176.
- Koneman EW, Allen SD, Janda WM, Schreckenberger PC, Winn WC. *Colour Atlas and Textbook of Diagnostic Microbiology*. 5th ed. New York: Lippincott; 1997
- Old DC. *Salmonella infection*. In: Collee JG, Fraster AG, Marmion BP, Simons a, eds. *Mackie and McCartney practical medical microbiology*, 14th ed. New York: Churchill Livingstone; 1996:385-402.
- Sherwal BI, Dhamija RK, Randhawa VS, Jais M, Kaintura A, Kumar M. A comparative study of Typhidot and Widal test in patients of Typhoid fever. *JIACM* 2004; 5:244-246.
- Abucejo PE, Capeding MR, Lupsian SP, Arcay J, Sombrero LT, Ruutu P, Herva E. Blood culture confirmed typhoid fever in a provincial hospital in Philippines. *Southeast Asian J Trop Med Public Health* 2001; 32:531-536.
- Wain J, Diep ST, BeBay PV, Walsh AL, HaVinh Duong NM et al. Specimen and culture media for the laboratory diagnosis of typhoid fever. *J Infect Developing Countries* 2008; 2:469-474.
- Akoh JA. Relative sensitivity of blood and bone marrow culture in typhoid fever. *Trop Doct* 1991; 21:174-176.
- Jesudason MV, Sivakuma S. Propective evaluation of a rapid diagnostic test Typhidot for typhoid fever. *Indian J Med Res* 2006; 123:513-516.
- Saha MR, Duta P, Palit A, Dutt D, Bhattacharya MK, Mitra U et al. A note on incidence of typhoid fever in diverse age groups in Kolkota, India. *Jpn J Inf Dis* 2003; 56:121-122.
- Koharo HK. A comparative study of the Typhidot (Dot-EIA) and Widal test in blood culture positive cases of typhoid fever. *Tropical Doctor* 2011; 4:136-138.
- Anagha K, Bhalerao D, Shariar R, Kulkarni S. The easy and early diagnosis of typhoid frver. *J Clin Diag Res* 2012; 6:198-199.
- Choo KE, Davis TM, Ismail A, Ibrahim TA, Ghazali WN. Rapid and reliable serological diagnosis of enteric fever: comparative sensitivity and specificity of Typhidot and Typhidot- M tests in febrile Malaysian children. *Acta Tropica* 1999; 72:175-183.
- Butta ZA, Mansurali N. Rapid serological diagnosis of pediatric typhoid fever in an endemic area: a prospective comparative evaluation of two dot-enzyme immunoassays and widal test. *Am J Trop Med Hyg* 1999; 61:645-647.