SCREENING OF SELECTED PLANT ESSENTIAL OILS FOR THEIR ANTIFUNGAL ACTIVITY AGAINST CANDIDA SPECIES ISOLATED FROM DENTURE STOMATITIS PATIENTS

Vinaya Bhat¹, S.M. Sharma², Veena Shetty³, C.S. Shastry⁴, Vaman Rao⁵,
Shilpa M. Shenoy⁶, Santanu Saha⁷ & Sriram Balaji⁸

¹PhD Research Scholar, ⁸PG Student, Department of Prosthodontics, ²HOD, Professor & PhD Guide, Department of Oral & Maxillofacial Surgery, A B Shetty Memorial Institute of Dental Sciences, ³Co-Guide, ⁴PhD Research Scholar, Department of Microbiology, K. S. Hegde Medical Academy, ⁵Principal & Co-Guide, NGSM Institute of Pharmaceutical Sciences, ⁷Senior Lecturer, Department of Pharmacognosy, NGSM Institute of Pharmaceutical Sciences, Nitte University, Mangalore, ⁶Co-Guide, Department of Biosciences, NMAM Institute of Technology, Nitte, Mangalore, INDIA.

Correspondence

Vinaya Bhat

PhD Research Scholar, Department of Prosthodontics, A B Shetty Memorial Institute of Dental Sciences, Nitte University Deralakatte, Mangalore - 575 018, Karnataka, India.

Mobile: +91 94819 21180 E-mail : drvinayabhat@gmail.com

Abstract

Background: Many efforts have been made to discover new antimicrobial compounds from various sources such as micro-organisms, animals and plants. Use of herbal medicines has a long history in Asian population in treating different ailments. Many plants have been found to have properties like antibacterial, antifungal and antioxidant activities.

In a complete denture wearer, due to either local or systemic factors, denture stomatitis has been found to have high prevalence. In such patients, higher incidence of *Candida* infection and increased levels of *Candida* spp. also have been identified. Existing antifungal agents like Nystatin & Fluconazole have been found to have toxic effects on long term application and chances of development of drug resistance is high. Medicinal plants have been used to inhibit growth and development of fungal infection from *Candida* isolated from sites other than oral cavity.

Objectives: To evaluate the antifungal activity of some of the medicinal essential oils against oral isolates of *Candida* obtained from denture wearing patients identified through a survey.

Materials and Methods: The oral isolates of *Candida* species were collected after conducting a survey of complete denture wearers, wearing the denture for at least more than a year. The speciation of the organism was done after identifying it from the specimens. It was confirmed with the help of germ tube test, other biochemical tests and chlamydospore formation tests. A lab isolate also was included in the study. The essential oils of different herbs were extracted using hydro distillation methods. The antifungal activity of these oils was tested and compared with antifungal activity of Nystatin and fluconazole using modified Kirby-Bauer method. The inhibition zones of different oils were measured and compared among themselves against all the clinical isolates and the lab isolate.

Results: About 55 complete denture wearing patients have been surveyed and among them 27 have been found to be positive for *Candida*-associated denture stomatitis. Among the *Candida* species, *Candida albicans* was more prevalent followed by *C. tropicalis* and *C. glabrata*.

Among the test essential oils, *Origanum vulgare* (stored & fresh) and clove oil gave positive results against all isolates. The fresh samples gave better results than the stored and all three oils showed more antifungal activity than Nystatin & fluconazole.

Interpretation: The *Candida*-associated denture stomatitis is prevalent in patients who are wearing denture prosthesis for more than a year.

Candida albicans is the most prevalent causative species followed by *C. tropicalis* and *C. glabrata*.

Origanum vulgare and clove oil are potent antifungal agents against oral species of *Candida*. They can be either used separately or their synergistic activity could be explored against denture stomatitis.

Key words: Denture stomatitis, herbal essential oils, clove oil, *Origanum vulgare*, *Candida* spp., nystatin, fluconazole

Access this article online

Quick Response Code

Introduction

High prevalence of denture stomatitis has been observed in patients who are wearing complete denture prosthesis. About 60 to 70 % of prevalence has been reported by various authors in the past ¹⁻⁶. In geriatric patients, when there is immune suppression due to any reason, there is a conversion of the normal oral commensal *Candida* into an infection causing pathogen ⁷⁻¹⁵. The pathogenic fungi *Candida* grow in large numbers forming biofilms on the surface of the prosthesis and on the oral mucosa. These biofilms have been found to be highly resistant to existing antifungal agents like Nystatin and Fluconazole. Moreover, these drugs when administered on a long term basis have been known to cause toxicity in human beings.

Efforts are being made to discover new antifungal agents from sources like micro organisms, animals and plants, which are either less toxic or not toxic at all. Use of plant extracts has a long history in Indian scenario for treatment of different ailments. Plants have been found to have rich source of antibacterial, antioxidant and antifungal activities. Several studies using plant essential oils have been carried out against *Candida* isolated from areas other than oral cavity ¹⁶⁻²⁰. No studies are reported in the literature regarding their effect against oral species of *Candida* in a denture wearer. Hence this study was undertaken with the aim of evaluating the effect of eight different plant essential oils on oral isolates of *Candida* retrieved from a denture wearing patients. Conventional antifungal agents like Nystatin and Fluconazole were also added as controls. A lab isolate was taken for comparative purposes and Agar Well diffusion method was implemented to study the susceptibility tests.

Materials and Methodology:

Ethical clearance for the present study was obtained from the ethical committee of NITTE University, Mangalore, Karnataka, India. The clinical isolates of *Candida* required for the study were collected through a survey conducted in the department of Prosthodontics. Fifty five patients wearing complete denture prosthesis for more than a year were selected based on the selection criteria.

Selection criteria:

Inclusion Criteria:

1. Patients wearing a denture >1 year
2. Patients not on any antifungal agent
3. Patients willing to participate in the study

Exclusion criteria:

1. Patients unwilling to participate in the study
2. Patients wearing a denture for <1 year
3. Patients already on antifungal therapy

A structured questionnaire was prepared to obtain the necessary information from the patients. Once selected, they were explained about the study and a written consent was obtained. About 27 patients were found to be positive for denture stomatitis observed through clinical signs and symptoms.

Collection of *Candida* samples: The isolates were collected from saliva samples of the selected patients. The patient was made to sit comfortably on the dental chair. The maxillary denture was removed and using a sterile cotton bud sample was collected from the palatal aspect of the tissue surface of the denture. (Fig.01). It was immediately placed in a test tube containing Phosphate Buffer Solution and was transported to the laboratory where it was cultured on the Sabouraud's Dextrose Agar (SDA) to separate the *Candida*.

Isolation of different species:

Once the *Candida* was identified, it was once again incubated in SDA medium at 37°C for 48 hours. *Candida* was confirmed using Germ Tube Test, Biochemical assays and CHROMagar tests (Fig. 02). Based on color of the colonies in CHROMagar, species in the samples were identified as *C. albicans* (pale green, 13/27), *C. tropicalis* (09/27), and *C. glabrata* (05/27). Once isolated, they were separated and subcultured for further use.

A lab isolate ATCC90028 of *C. albicans* was added in the study for comparative purposes.

Collection of plant material:

The dry leaves of *Origanum vulgare* were purchased from JKH Impex Pvt. Ltd. Mumbai. Leaves of *Pogostemon*

patchouli, *Oscimum sanctum* and Neem were collected locally and the leaves were identified taxonomically at the University of Agricultural Sciences, Bengaluru. Commercially available *Origanum vulgare*, clove and caprylate were purchased from departmental store.

Preparation of herbal essential oils:

The fresh leaves of *Pogostemon patchouli*, *Oscimum*

sanctum and Neem were dried at room temperature and were powdered. Essential oils from all plants were extracted using hydrodistillation procedure and the oils were stored at -4 °C till use. Nystatin and fluconazole were added as controls.

Antifungal susceptibility testing (Well diffusion method):

A modified Kirby-Baur method was used. Here, the

Table 1 : Antifungal activity of the essential oils of the collected plants

Plant name	Botanical name	Zone of inhibition (mm/20µl)			
		<i>Candida albicans</i>	<i>Candida glabrata</i>	<i>Candida tropicalis</i>	<i>Candida albicans</i> (Lab isolate ATCC90028)
Origanum (stored extract)	<i>Origanum vulgare</i>	30±5	35±2	30±2	30±4
Origanum (fresh extract)	<i>Origanum vulgare</i>	30±3	36±2	32±3	30±2
Origanum (commercial)	<i>Origanum vulgare</i>	----	----	----	----
Patche Kaduru (Kannada)	<i>Pogostemon Patchouli</i>	----	----	----	----
Tulasi	<i>Oscimum Sanctum</i>	----	----	----	----
Clove	<i>Syzygium aromaticum</i>	33±4	30±2	30±2	33±1
Neem	<i>Azadirachta indica</i> L	----	----	----	----
Coconut	<i>Cocosnificera Sanctum</i>	----	----	----	----

Table II : Descriptive analysis of data

Comparison of three essential oils in each group

		N	Mean	std. Deviation	df	Mean Square	F	Sig.
candida albicans	origanum vulgare (o p) stored at 4	10	29.9	3.281	2	9.733	1.547	0.23
	origanum vulgare fresh extract	10	30.1	2.025				
	clove oil	10	31.7	2.003				
	Total	30	30.57	2.555				
candida glabrata	origanum vulgare (o p) stored at 4	10	35	1.491	2	103.333	35.769	<0.00
	origanum vulgare fresh extract	10	36	2.055				
	clove oil	10	30	1.491				
	Total	30	33.67	3.133				
candida tropicalis	origanum vulgare (o p) stored at 4	10	30	1.491	2	14.033	4.802	0.01
	origanum vulgare fresh extract	10	32	2.16				
	clove oil	10	29.9	1.37				
	Total	30	30.63	1.921				
candida albicans lab isolate (control) (ATCC90028)	origanum vulgare (o p) stored at 4	10	29.9	2.644	2	29.233	8.919	0.00
	origanum vulgare fresh extract	10	30.2	1.476				
	clove oil	10	33	0.816				
	Total	30	31.03	20251				

There is significant difference in the candida glabrata, tropicalis and lab isolate

Candida species were suspended in SDA broth and it was made equivalent to 0.5 McFarland units. Six millimeter wells were created on the SDA medium and 20 microlitre each of the antifungal agents was delivered into each well. It was incubated again for further 48 hours at 37 ° C with agitation. After 48 hours, zone of inhibition for each essential oil and for each species was measured using a flexible scale (Fig.03). The results were tabulated (Table I) and were subjected to statistical analysis.

Results:

Among eight essential oils (*Origanum vulgare* in three forms, i.e., fresh extract, stored extract & commercial, *Pogostemon patchouli*, *Oscimum sanctum*, *Neem*, clove and caprylate) three showed positive effects against *Candida* species. *Origanum vulgare* in two forms, i.e. Fresh extract, and stored extract and Clove oil showed significant results ($p < 0.001$) on all species of *Candida*. All showed more antifungal effect than Nystatin (22mm zone of inhibition) and fluconazole (19mm zone of inhibition). Among fresh and preserved forms of *Origanum vulgare*, fresh samples showed better results (30 ± 3 , 36 ± 2 , 32 ± 3 , 30 ± 2) against all species. There was no significant difference ($P < 0.231$) in action of three oils on *Candida albicans* where as there was significant difference in *Candida glabrata*, *C. tropicalis* and lab isolate ($P < 0.001$, 0.016, 0.001). This indicates that all three oils have similar activity against *Candida albicans* and the action is different for other three species. *C. glabrata* and *C. tropicalis* are more susceptible to *Origanum vulgare* rather than clove oil, where as lab isolate is more susceptible to clove oil. Hence any of the three oils can be used effectively against oral *Candida albicans* and *Origanum vulgare* is also effective on other oral isolates. Clove oil is more effective on lab isolate of *Candida albicans* ATCC90028. *Origanum vulgare* also showed significant difference as compared to Fluconazole & Nystatin ($P < 0.001$) (Table II) & (Table III).

Discussion :

Denture stomatitis is a very common type of infection of the oral mucosa found in 60 to 65% of the denture wearing population^{21, 22}. Various reasons have been cited by the

authors for this prevalence. Denture hygiene, trauma and allergy to the denture base materials are the commonest causes. In addition, when the immune status of the patient is lowered due to systemic diseases or due to immunosuppressive therapy, the otherwise normal oral commensal *Candida* turns pathogenic and causes denture stomatitis. Histologic sections of the oral mucosa have been shown to exhibit *Candida* blastospores, hyphae and pseudohyphae.

In diabetes mellitus, the patients have been found to be more susceptible to oral candidiasis than patients without diabetes²³⁻²⁴. If the patient is a denture wearer, it has been found to favor the growth of *Candida* to cause candidiasis²⁴⁻²⁵. In a study by Sanita et al⁶, they said that *Candida* species were more prevalent in patients with denture stomatitis in both, diabetic and non diabetic conditions. Diabetes mellitus has been a risk factor for the development of DS. In a study by Emami et al⁵, they found that the patient who is wearing a removable denture whether partial or complete is at a risk of developing DS. Presence of dental prostheses has also been found to be associated with HIV/AIDS¹⁵. They are of the opinion that CD4 count plays an important role in the development of oral candidiasis.

Candida albicans has been found to be the most prevalent species in *Candida*-related denture stomatitis. Under normal circumstances, it is a common oral commensal not causing any pathosis. However, during favourable conditions it transforms itself into a virulent pathogen using various mechanisms.

Even though *Candida albicans* is the frequently associated species in denture stomatitis, *C. glabrata* is the emerging species next to it²⁶⁻²⁹. However, Sanita et al⁶, in their study concluded that *C. albicans* and *C. tropicalis* were more prevalent in a denture wearer with denture stomatitis. In severe candidiasis, *C. tropicalis* was more prevalent. Mixed infections caused by *C. albicans* and *C. glabrata* have been demonstrated in vitro oral epithelium³⁰. Reconstructed Human Oral epithelium was used to demonstrate the ability of *C. albicans* to enhance the invasiveness of *C. glabrata* in causing the infection.

To arrive at a conclusion regarding the dominance of the species in the local population the authors conducted a survey to include denture wearers from where the samples were collected for species characterization of *Candida*. In this survey, we found that after *C. albicans*, it was *C. tropicalis* and then *C. glabrata* which were the next most prevalent species of *Candida*.

Existing antifungal agents like Nystatin and Fluconazole have been found to be toxic on long term use³¹⁻³⁵. Also, development of resistance by the organisms has been recorded.[®] Hence, there is a need to develop a new antifungal agent which may be less or not toxic at all. Plants have a long positive history for being used as natural remedies for various ailments^{36-38, 18}. Several herbal agents have been tried against *Candida* of various regions in the human body¹⁸⁻²⁰. However, no studies have been reported regarding the investigation of antifungal effects of any herbs on the *Candida* strains obtained from the oral candidiasis of a denture wearing patient. This study was undertaken to screen some of the herbal products that are commonly available against *Candida* spp. of oral candidiasis.

A modified Kirby-Baur method was implemented to study the antifungal susceptibility potential of the eight herbs. Zone of inhibition for each was measured and tabulated. Three oils (two types of *Origanum vulgare* and clove oil) showed strong positive antifungal effects on all strains (Clinical and lab) of *Candida*, where as, the other oils did not show any measurable activity at all. Zone of inhibition for stored *Origanum vulgare* was 30mm±5 for *Candida albicans*, 30mm±2 for *Candida tropicalis*, 35mm±2 for *Candida glabrata* and 30mm±4 for *Candida* lab isolate. For fresh sample of *Origanum vulgare* was, 30mm±3 for *C. albicans*, 32mm±3 for *C. tropicalis*, 36mm±2 for *C. glabrata* and 30mm±2 for the lab isolate. Clove oil exhibited 33mm±4mm for *C. albicans*, 30mm±2 mm for *C. tropicalis*, 30mm±2 mm for *C. glabrata* and 33mm±1 mm for lab isolate. Nystatin has been found to show an antifungal effect of up to 22mm¹⁸ and fluconazole upto 19mm³⁹. In this study, the antifungal effects of *Origanum vulgare* and

clove oil were found to be substantially high as compared to Nystatin and Fluconazole, indicating that they can have better results in eradicating candidiasis. And being of plant origin, they may be less toxic. Further studies on the toxic effects of these oils on animals are indicated.

Origanum vulgare is a plant belonging to the family Lamiaceae which contains the phenolic compound Carvacrol as a primary compound⁴⁰. In a few earlier studies, the essential oil of *Origanum vulgare* has been studied against *Candida* spp. isolated from sites such as, vaginal mucous membrane of dogs, cutaneous tegument of a dog, cutaneous tegument of a capuchin monkey²⁰, culture collection *Candida albicans* ATCC10231³⁸, ATCC 14053¹⁸, female mice¹⁹ and from human stool specimen¹⁸. In all the above studies *Origanum vulgare* has been found to be possessing anti-candidal activity. In the present study, the *Candida* spp. was obtained from the oral cavity and this plant was found to be very effective against different species of *Candida* and so also a lab species of *Candida albicans*. Both, the stored and the fresh extracts showed good antifungal effect as compared to fluconazole and nystatin, indicating that proper storage does not decrease the potency of the oil.

Syzygium aromaticum (clove) had been used against lab isolate of *Candida albicans* ATCC 10231 by Hammer et al.³⁸. In their study of antimicrobial activity of essential oils and plant extracts carried on several organisms they found that clove had an antifungal effect against *Candida albicans*. In our study also, the essential oil of clove showed a strong antifungal activity against several species of *Candida* obtained from the oral cavity thus confirming its possession of positive antifungal effects.

Conclusion :

This study has scientifically proved the antifungal efficacy of three essential oils obtained from fresh & stored samples of *Origanum vulgare* and Clove oil. Further, it was found that the effects are much more than the existing antifungals like Nystatin & Fluconazole. The usage of *Origanum vulgare* or clove oil or the combination of the two essential oils has the potential of being used in treating

oral candidiasis. The present in-vitro study has confirmed the potentiality of the two essential oils to be used as antifungal agents against *Candida* spp. However, further studies to determine their MIC & MFC and also the phase II

studies on animals have to be carried out. Also, their pharmacological effects like antibacterial, anti carcinogenic and anti-inflammatory activities can be investigated.

References

1. Kebe K, Morii K, Matsuda Kk, Hata K, Nokubi T. Journal of Oral Rehabilitation 2006;33:36-42.
2. Liu X, Hua H. Oral Manifestation of Chronic Mucocutaneous Candidiasis: Seven Case Reports. J Oral Pathol Med 2007;36:528-32.
3. Pereira-cenci T, Del bel cury AA, Crielard W, Ten cate JM. Development of Candida-associated Denture Stomatitis: New Insights. J Appl Oral Sci. 2008;16:86-945.
4. Williams D, Lewis M. Pathogenesis and treatment of oral candidosis. Journal of Oral Microbiology 2011;3.
5. Emami E, Taraf H, Grandmont P, Gauthier G, Koninck L, Lamarche C, de Souza R. The association of Denture Stomatitis and Partial Removable dental Prostheses: A Systematic Review. Int J Prosthodont 2012;25:113-9.
6. Sanita PV, Pavarina AC, Eunice TG, Silva MM, Mima EGO, Ribeiro DG, Vergani CE. Candida spp. prevalence in well controlled type 2 diabetic patients with denture stomatitis. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2011;111:726-33
7. Dangi YS, Soni ML, Namdeo KP. Oral Candidiasis: A Review. International Journal of Pharmacy and Pharmaceutical Sciences; 2010; 2(4).
8. Dongari-Bagtzoglou A et al. Oral Candida Infection and Colonization in Solid Organ Transplant Recipients. Oral Microbiol Immunol 2009;24:249-54.
9. Olczak-Kowalczyk D et al. Oral Candidiasis in Immunosuppressed Children and Young Adults after Liver or Kidney Transplantation. Pediatric Dent 2010;32:189-94.
10. Reichart PA, Samaranayake LP, Bendick Ch, Schmidt-Westhausen M, Jayalathike MS. Prevalence of Oral Candida Species in Leprosy Patients from Cambodia and Thailand. J Oral Pathol Med 2007;36:342-6.
11. Samaranayake, LP et al. Fungal Infections associated with HIV Infection. Oral Diseases. 2002;2:151-60.
12. Sashikumar R, Kannan R. Salivary Glucose Levels and Oral Candidal Carriage in Type II Diabetics. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2010;109:706-711.
13. Scully C, El-Kabir M, Samaranayake LP. Candida and Oral Candidosis: A Review. Critical Reviews in Oral Biology and Medicine, 1994;5:125-157.
14. Thompson GR et al. Oropharyngeal candidiasis in the era of antiretroviral therapy. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2010;109:488-95.
15. Witzel AL, Pires MFC, Carli ML, Rabelo GD, Nunes TB. Candida albicans isolation from Buccal Mucosa of Patients with HIV wearing removable dental prostheses. Int J Prosthodont 2012;25:127-131.
16. Javidnia K, Miri R, Assadollahi M, Gholami M, Ghaderi M. Screening of selected plants growing in Iran for antimicrobial activity. Iranian Journal of Science & Technology, Transaction A. 2009;33:329-333
17. Rosato A, Vitali C, Piarulli M, Mazzotta M, Argentieri MP, Rosanna M. In vitro efficacy of the combination of Nystatin with the essential oils of *Origanum vulgare* and *Pelargonium graveolens* against some *Candida* species. International Journal of Phytotherapy & amp; Phytopharmacology. 2009;Elsevier.
18. Stiles JC, Sparks W, Ronzio RA. The Inhibition of *Candida albicans* by *Oregano*. J of Applied Nutrition 1995;47:4:1-7.
19. Manohar V, Ingram C, Gray J, Talpur NA, Echard BW, Bagchi D et al. Antifungal Activities of *Origanum* oil against *Candida albicans*. Molecular and Cellular Biochemistry. 2001;228:111-117.
20. Cleff MB, Meinerz AR, Xavier M, Schuch LF, Meireles MCA, Rodrigues MRA et al. In Vitro activity of *Origanum vulgare* essential oil against *Candida* species. Braz. J. Microbiol. 2010;41(1).
21. Webb BC, Thomas CJ, Willcox MD, Harty DW, Knox KW. Candida-associated denture stomatitis. Aetiology and management: a review. Part 2. Oral diseases caused by *Candida* species. Aust Dent J. 1998;43:160-6.
22. Salerno C, Pascale M, Contaldo M, Esposito V, Busciolano, Milillo L, Guida A, Petrucci M, Serpico R. Candida-associated denture stomatitis. Oral Med and Pathol. 2010.
23. Belazi M, Velegraki A, Fleva A, Gidarakou I, Papanau L, Baka D, et al. Candidal overgrowth in diabetic patients: Potential predisposing factors. Mycoses 2005;48:192-6.
24. Khosravi AR, Yarahmadi S, Baiat M, Shokri H, Pourkabir M. Factors affecting the prevalence of yeasts in the oral cavity of patients with diabetes mellitus. J Mycol Med 2008;18:83-8.
25. Goncalves RH, Miranda ET, Zaia JE, Giannini MJ. Species diversity of yeast in oral colonization of insulin-treated diabetes mellitus patients. Mycopathologia 2006;162:83-9,
26. Li L, Redding S, Dongari-Bagtzoglou AD. *Candida glabrata*, an emerging oral opportunistic pathogen. J Dent Res 2007;86:204-15.
27. Redding SW, Kirkpatrick WR, Dib O, Fothergill AW, Rinaldi MG, Patterson T. The epidemiology of non-albicans *Candida* in oropharyngeal candidiasis in HIV patients. Spec Care Dentist 2000;20:178-81.
28. Redding SW, Kirkpatrick WR, Coco BJ, et al. *Candida glabrata* oropharyngeal candidiasis in patients receiving radiation treatment for head and neck cancer. J Clin Microbiol 2002;40:1879-81.
29. Coco BJ, Bagg J, Cross LJ, Jose A, Cross J, Ramage G. Mixed *Candida albicans* and *Candida glabrata* populations associated with pathogenesis of denture stomatitis. Oral Microbiol Immunol 2008;23:377-83
30. Silva S, Henriques M, Hayes A, Oliveira R, Azeredo J, Williams DW. *Candida glabrata* and *Candida albicans* co-infection of an in vitro oral epithelium. J Oral Pathol Med. 2011;40:421-27
31. Charlier C et al. Fluconazole for the management of invasive candidiasis: where do we stand after 15 years? J of Antimicrobial Chemotherapy, 2006;57:384-410.
32. Jafari-nodoushan AA, Kazemi A, Mirzaii F, Dehghani M. Fluconazole Susceptibility Profile of *Candida* isolates Recovered from Patients Specimens Admitted to yard Central Laboratory. Iranian J of Pharmaceutical Research 2008;7:69-75.
33. Martin MV. The Use of Fluconazole and Itraconazole in the Treatment of *Candida albicans* Infections: a Review. J of Antimicrobial Chemotherapy 1999;44:429-37.
34. Sheehan DJ, Hitchcock CA, Sibley CM. Current and Emerging Azole Antifungal Agents. Clin Microbiol Rev 1999;12:40-79.
35. Wu T, Wright K, Hurst SF, Morrison CJ. Enhanced extracellular Production of Aspartyl Proteinase, a virulence factor, by *C. albicans* isolates following growth in subinhibitory concentrations of fluconazole. Antimicrobial Agents and Chemotherapy. 2000;44:1200-1208.
36. Malabadi RB, Kumar SV. Assessment of Antifungal Activity of Some Medicinal Plants. Intl. J of Pharmacology. 2007;3:499-504.
37. Meena AK et al. Review on antifungal activities of Ayurvedic Medicinal Plants. Drug Invention Today. 2009;2:146-48.
38. Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. Journal of Applied Microbiology 1999;86:985-990.
39. Pfaller MA, Barry AL. Evaluation of a Novel Colorimetric Broth Micro dilution Method for Antifungal Susceptibility Testing of Yeast Isolates. J of Clinical Microbiology, Aug. 1994;32:1992-96.
40. Taxon: *Origanum vulgare* L. GRIN Taxonomy for Plants. United States Dept. of Agriculture.