R. L. REYES, M. SATO, T. IWAI, K. SUZUKI, S. MAEDA, M. SAWAMURA* (HOKKAIDO UNIVERSITY, SAPPORO, JAPAN)

Asymmetric Remote C–H Borylation of Aliphatic Amides and Esters with a Modular Iridium Catalyst

Science **2020**, 369, 970–974, DOI: 10.1126/science.abc8320.

Asymmetric γ -Borylation of Aliphatic Amides and Esters

Selected examples:

Derivatizations:

Significance: The authors report on an asymmetric borylation of aliphatic amides and esters in γ -position. The efficient catalytic system delivers chiral pinacolboronates in high yield and enantiomeric excess.

Comment: The complex catalytic system is comprised of an iridium precatalyst, a chiral ligand and a receptor ligand. The receptor ligand has a pyridine moiety intended for binding to the catalyst and a urea moiety for substrate binding. The absence of either of these components results in no reaction. 2,6-Lutidine has a positive impact on both yield and enantiomeric excess. Preliminary quantum chemical calculations show that the amide substrate is bound to the catalytic cavity not only through hydrogen bonding with receptor ligand but also through other noncovalent interactions.

SYNFACTS Contributors: Paul Knochel, Dimitrije Djukanovic
 Synfacts 2020, 16(11), 1301 Published online: 20.10.2020
 DOI: 10.1055/s-0040-1706512; Reg-No.: P12620SF

© 2020, Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Metals in Synthesis

Key words

C(sp³)–H activation

borylation

asymmetric catalysis

Synfact
of the
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.