A. M. BERMAN, J. S. JOHNSON* (UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, USA)

Copper-Catalyzed Electrophilic Amination of Diorganozinc Reagents

J. Am. Chem. Soc. **2004**, 126, 5680–5681.

Electrophilic Amination of Diorganozinc Reagents Using *N*-Hydroxylamine Benzoates

Selected examples:

Significance: In 2004, Berman and Johnson developed the first electrophilic amination of *N*-hydroxylamine benzoates using diorganozinc reagents in the presence of $[\text{Cu}(\text{OTf})_2 \cdot \text{C}_6\text{H}_6]$. This umpolung-type reaction allowed access to a broad range of highly functionalized secondary and tertiary amines under mild conditions.

Comment: The outstanding performance of *N*-hydroxylamine benzoates as electrophilic nitrogen sources has inspired various following publications, opening up new opportunities for amination reactions. Based on experimental studies, the authors excluded an oxidative addition/reductive elimination pathway and proposed a $\text{S}_{\text{N}}2$ -type pathway. For further insights into electrophilic aminations based on this pioneering work, see: *Synthesis* **2011**, 24, 3954–3964.

SYNFACTS Contributors: Paul Knochel, Simon Graßl
Synfacts 2020, 16(06), 0687 Published online: 15.05.2020
DOI: 10.1055/s-0040-1706966; Reg-No.: P04820SF

© 2020, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Metals in Synthesis

Key words

electrophilic
amination

organozinc reagents

copper catalysis

Synfact
Classic

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.