Case Report

A rare case of fulminant Toxic Shock Syndrome Associated with clostridium sordellii septic abortion

Ahmed S. Elagwany¹ & Nehal Anas²

^{1,2}Lecturer, Department of Obstetrics and Gynecology, Alexandria University, Egypt.

*Corresponding Author : Ahmed S. El-agwany, Lecturer, Department of Obstetrics and Gynecology, El-shatby maternity hospital, Alexandria University, Alexandria, Egypt. Ph.: +20 12282 54247 E-mail: Ahmedsamyagwany@gmail.com

Received : 30-09-2015

Review Completed : 30-04-2016

Accepted : 02-05-2016

Keywords : Rare, Fulminant, Toxic Shock Syndrome, Clostridium Sordellii, Septic Abortion

Abstract :

Endometritis and toxic shock syndrome associated with *Clostridium sordellii* have previously been reported after childbirth and, in one case, after medical abortion. Clinical findings included tachycardia, hypotension, edema, hemoconcentration, profound leukocytosis, and absence of fever. We describe a young woman who developed *Clostridium sordellii* toxic shock syndrome after having had an abortion surgically evacuated five days before admission to our hospital. Although the patient was aggressively treated, death occurred <3 days after admission. It is hoped that very early recognition of this disease will decrease the mortality associated with this rarely reported ailment.

Access this article online

Quick Response Code

Introduction

Clostridium sordellii is a gram-positive anaerobic bacillus that has been reported as a cause of infection in the female genital tract and fatal toxic shock syndrome. Of 10 cases identified in the literature, 8 occurred after delivery of live-born infants, 1 occurred after a medical abortion, and 1 was not associated with pregnancy. (1-4)

We report one additional deaths due to *C. sordellii* toxic shock syndrome that occurred among previously healthy women after abortion.

Case report

In march 2012, a 35-year-old, G5P2+3, previously healthy woman was admitted to emergency unit of our department complaining of severe abdominal pain, vomiting, diarrhea, fever and foul-smelling vaginal discharge. Five days before admission, the patient had a dilatation and curettage in a district hospital for a missed abortion of ten weeks' gestation that was induced by

misoprostol. All her children were delivered by normal vaginal delivery.

On examination, the patient was disoriented, pale and toxic. Vital signs were unstable with ABP 90/40 mmHg, heart rate 120 beats/min and temperature 39°C. Her abdomen was distended, and vaginal examination revealed an enlarged tender uterus with offensive vaginal discharge. Laboratory investigations revealed a high white-cell count of 34,800 cells per microliter, hemoglobin level of 7gm/dl, and platelet count of 100,000 cells per microliter platelets and a hematocrit of 40 percent. Liver function and renal function tests were normal.

Abdominal ultrasound demonstrated an enlarged AVF uterus with a heterogeneous intrauterine mass about 4cm in diameter highly suspicious of remnants of conception.

So, the decision was made to correct the general condition of the patient then evacuate the uterine cavity under

umbrella of antibiotics. The patient was treated in the intensive care unit with high doses of penicillin, clindamycin, and ciprofloxacin, as well as with intravenously administered fluid and blood transfusion. Approximately 24 hours after the patient was admitted to the intensive care unit, uterine dilatation and blunt curettage was done. (Figure 1) Tissues were biopsied for histological diagnoses and culture and sensitivity for antibiotics. Numerous, large gram-positive bacilli that were suggestive of *Clostridium* species were seen on a Gram stain of the specimen.

The patient's general condition deteriorate over the next 24 hours where drowsiness and severe vaginal bleeding occurred. Vital signs revealed heart rate 140 /min, ABP 70/30 mmHg. Laboratory investigations revealed HB 6 gm./dl, a hematocrit of 45 percent, white-cell count of 50,000 cells per microliter, platelet cell count of 60,000 cells per microliter and INR 3 with continuous blood, fluid and antibiotic therapy. SGOT 500 U/L, SGPT 400 U/L, BUN 50 mg/dl, serum creatinine 2 and albumin 1.5 gm/dl.

An abdominal hysterectomy with bilateral salpingoophorectomy was performed immediately. (Figure 2) Soon after surgery, intractable shock with unrecorded ABP, weak pulse of 150 beat / min, temp 35°C, anuria, coma and disseminated intravascular coagulation developed. Laboratory investigations revealed Hb 7 gm/dl, white cell count of 50,000 cells per microliter, platelet cell count of 40,000 cells per microliter, INR 7, SGOT 4000 U/L, SGPT 5000 U/L, BUN 60 mg/dl, serum creatinine 3 mg/dl and serum albumin 0.8 gm/dl. Death eventually followed in less than three days after admission.

Microscopy of the hysterectomy specimen revealed massive coagulation necrosis of the uterine wall, the fallopian tubes, and the ovaries, with leukocytoclastic inflammation of the adjacent viable tissue. Bacterial cultures of samples obtained from the uterus yielded *Clostridium sordellii*, Coagulable necrosis of the decidua and the superficial myometrium associated with hemorrhage and acute inflammation of the adjacent viable tissue. Evidence of *C. sordellii* infection was established

with the use of PCR assays performed on fixed uterine tissue.

Figure 1 : showing the infected remnants of infection

Figure 2 : showing hysterectomy specimen

Discussion

C. sordellii, a pathogen that is infrequently found in humans, has a broad spectrum of clinical presentations. Among them, a fulminant toxic shock syndrome characterized by marked leukemoid reaction (WBC >80,000) and hemoconcentration (Hct >50%) and caused by a massive capillary leak has been reported almost exclusively in association with infections of the uterus or the perineum after either infected episiotomy (in 2 patients) or postpartum endometritis (in 4 patients). In 1 patient, the syndrome manifested as spontaneous endometritis (1-4).

Highly active hemorrhagic and lethal toxins of *C. sordellii* play a central role in the pathogenesis of the syndrome by targeting the Ras guanosine triphosphate binding proteins and glycosylating them (5,6).

The rate of vaginal colonization with *Clostridium* species in the period after abortion occurs has been reported to be as high as 29%, whereas these bacteria have been isolated in the vaginal secretions of 5% 10% of nonpregnant women

(7). Opening of the cervix during labor or abortion, which permits the passage of vaginal pathogens, appears to be the critical event that leads to infection of the endometrium.

To improve diagnosis and therapy, clinicians should be aware of the distinctive features of this potentially fatal entity, including tachycardia, hypotension, edema, hemoconcentration, profound leukocytosis, and absence of fever. Evidence of *C. sordellii* infection was established with the use of anti-clostridium species immunohistochemical assay and both organism-specific and broad-range PCR assays performed on fixed uterine tissue.

There are limited data regarding the optimal therapy for *C. sordellii*. As with other severe histotoxic clostridial infections, aggressive surgical wound débridement, removal of infected organs (e.g., by means of hysterectomy), and antibacterial agents with good anaerobic activity are logical first steps to decrease the bacterial load and minimize further production of toxins. In vitro susceptibility testing on 24 *C. sordellii* strains showed low minimal inhibitory concentrations for penicillin, ampicillin, erythromycin, rifampin, tetracycline, cefoxitin, clindamycin, and metronidazole; antibiotics that interfere with bacterial protein synthesis (such as clindamycin) may have additional benefit. However, débridement, surgery, and antibacterial therapy will not mitigate the effects of preformed toxin. (8,9)

Treatment with antitoxins could, in principle, be clinically useful. Antitoxin therapy directed against the toxins of clostridial species causing gas gangrene was used

intensively from 1918 to the end of World War II, but it was later proven to be ineffective. *C. sordellii* antitoxins are commercially available for in vitro neutralization of *Clostridium difficile* toxin B in cytotoxin cell assays, and they probably would be present in human immunoglobulins, albeit not in a high concentration. Although potentially useful, they have never been tested for clinical efficacy (10). Experience with anticlostridial toxins is limited; however, other than for use in the treatment of tetanus and botulism, anticlostridial toxins probably could be useful only for prevention of the syndrome in subjects exposed to *C. sordellii* or, at best, could be useful as treatment if administered very early in the infectious process. Obviously, animal studies are needed before a compassionate protocol could be approved for such treatment.

This syndrome is very rare and, to date, has claimed the lives of all individuals who have been affected by it. We believe that the association of a capillary leak with hemoconcentration and a marked leukemoid reaction in a patient during the postpartum or postabortion period is very characteristic and should hasten early recognition of this disease before development of irreversible shock. Given the present state of knowledge regarding this disease, we can only hope that very early recognition of this disease, along with an aggressive surgical approach and appropriate antimicrobial therapy and resuscitation techniques, will decrease the mortality associated with an ailment that occurs mostly among young, otherwise healthy women.

References

1. McGregor JA, Soper DE, Lovell G. Maternal deaths associated with *Clostridium sordellii* infection. *Am J Obstet Gynecol* 1989; 161:987-95?
2. Rorbye C, Petersen IS, Nilas L. Postpartum *Clostridium sordellii* infection associated with fatal toxic shock syndrome. *Acta Obstet Gynecol Scand* 2000; 79:1134-5.
3. Bitti A, Mastrantonio P, Spigaglia P. A fatal postpartum *Clostridium sordellii* associated toxic shock syndrome. *J Clin Pathol* 1997; 50:259-60.
4. Hogan SF, Ireland K. Fatal acute spontaneous endometritis resulting from *Clostridium sordellii*. *Am J Clin Pathol* 1989; 91:104-6?
5. Qa'dan M, Spyres LM, Ballard JD. PH-enhanced cytopathic effect of *Clostridium sordellii* lethal toxin. *Infect Immun* 2001; 69:5487-93.
6. Abdulla A, Yee L. The clinical spectrum of *Clostridium sordellii* bacteraemia: two case reports and a review of the literature. *J Clin Pathol* 2000; 53:709-12.
7. Bartlett JG, Onderdonk AB, Drude E. Quantitative bacteriology of the vaginal flora. *J Infect Dis* 1977; 136:271-7.
8. Grimwood K, Evans GA, Govender ST, Woods DE. *Clostridium sordellii* infection and toxin neutralization. *Pediatr Infect Dis J* 1990; 9:582-5.
9. Adamkiewicz TV, Goodman D, Burke B, Lysterly DM, Goswitz J, Ferrieri P. Neonatal *Clostridium sordellii* toxic omphalitis. *Pediatr Infect Dis J* 1993; 12:253-7.
10. Lorber B. Gas gangrene and other *Clostridium*-associated diseases. In: Mandell GL, Bennett JE, Dolin R. Principles and practice of infectious diseases. Philadelphia: Churchill Livingstone 1999:2549-61