The Central Role of Acute Phase Proteins in Rheumatoid Arthritis: Involvement in Disease Autoimmunity, Inflammatory Responses, and the Heightened Risk of Cardiovascular Disease

Johannes A. Bezuidenhout, MSc¹ Etheresia Pretorius, PhD¹

¹ Department of Physiological Sciences, Stellenbosch University, Stellenbosch, South Africa

Semin Thromb Hemost 2020;46:465–483.

Address for correspondence: Etheresia Pretorius, PhD, Department of Physiological Sciences, Faculty of Science, Stellenbosch University, Private Bag X1 Matieland, Stellenbosch 7602, South Africa (e-mail: resiap@sun.ac.za).

Abstract

Rheumatoid arthritis (RA) is an autoimmune disease of complex etiopathogenic origin and traditionally characterized by chronic synovitis and articular erosions. Furthermore, there is strong evidence that infectious agents, including those that become dormant within the host, play a major role in much of the etiology of RA and its hallmark of inflammation. A combination of genetic predisposition, environmental exposure, and presence of infectious agents may therefore lead to a loss of immune tolerance to citrullinated proteins, which present as self-antigens to the human immune system. This results in generation of highly RA-specific autoantibodies, known as anti-citrullinated protein antibodies (ACPAs). Protein citrullination occurs via posttranslational deamination of arginine residues by peptidylarginine deiminase enzymes, which have confirmed sources of both endogenous and infectious origins. A recognized plasma protein target of citrullination and RA autoantibody generation is fibrin and its soluble precursor fibrinogen, both key components of hemostasis and acute phase reaction. Increased titers of ACPAs that accompany rapid progression to clinical RA disease have been shown to drive a variety of proinflammatory processes, and therefore results in aberrant fibrin clot formation and increased cardiovascular risk. However, the full extent to which hemostasis is affected in RA remains controversial, owing to the differential impact that citrullinated fibrin(ogen) and concurrent systemic inflammation may have on resulting hemostatic outcome. This review highlights key events in initiation of autoimmune-driven inflammatory events, including the role of bacterial infectious agents, which subsequently result in clinical RA disease and associated secondary cardiovascular disease risk, with specific focus on plasma proteins that are heavily involved throughout the immunopathological progression process.

Keywords

- rheumatoid arthritis
- hemostasis
- coagulation
- citrullination
- periodontitis

Rheumatoid arthritis (RA) is the most common autoimmune disease, displaying a worldwide prevalence of 0.5 to 1%.^{1,2} RA typically has a middle-aged onset that disproportionately affects females (threefold higher prevalence than males).³ Symptomatic RA is characterized by chronic systemic inflammation, resulting in bilateral and symmetrical polyarticular

arthritis, persistent erosive synovitis, and development of autoantibodies.^{1,4} The clinical manifestations of chronic inflammation in RA are most commonly observed in the small joints of hands and feet.⁵ Common disease symptoms include pain, swelling of the joint, stiffness, and progressive destruction of joint structure.⁶ Irreversible damage to joint structure

and subsequent loss of mobility can occur within 3 years of diagnosis in 20 to 30% of RA patients when appropriate treatment measures are not undertaken.⁷

RA is considered to be a prototypic polygenic disorder, as disease development is thought to occur as a result of the interaction between predisposing genetic risk factors and environmental exposures.⁸ The strongest genetic link associated with RA is the single nucleotide polymorphisms in the human leukocyte antigen (HLA)-DR β 1 gene, which in some study cohorts have shown to be present in approximately 64 to 70% of RA patients and 55% of their first-degree relatives.^{9,10} HLA molecules belong to the major histocompatibility complex (MHC) and is strongly linked to anti-citrullinated protein antibodies (ACPA)⁺-RA.⁵ HLA-DRB1 genes exert major influence on the cluster of differentiation 4 (CD4)⁺ $\alpha\beta$ T cell repertoire, through their “shared epitope” (SE) alleles that present as self-peptides to CD4⁺ T-cells in the thymus.^{5,11}

The most compelling current explanation for the development and perpetuation of clinical RA disease is the infection of certain mucosal sites (oral, intestinal, and pulmonary) where microbes, under the influence of environmental factors such as smoke and drugs along with genetic predisposition, trigger mucosal inflammation.¹² Subsequent migration of mucosal macrophages and lymphocytes to articular spaces results in synovial inflammation and joint destruction.¹³ Research by Rashid and Ebringer has strongly linked the bacterium *Proteus mirabilis* to the development of RA,¹⁴ stemming from an increased incidence rate of urinary tract infections (UTIs) observed in affected individuals.¹⁵ *P. mirabilis* is a major causative agent of UTIs and is detected to a greater extent in RA patient urine, along with increased levels of antibodies directed toward the bacterium.¹⁶ The closely matched amino acid sequences of “E-S-R-R-A-L” contained in *Proteus spp.* hemolysins and “E-Q/K-R-R-A-A” in the “SE” of RA-linked HLA-DR molecules result in the binding of cross-reactive anti-*P. mirabilis* antibodies to various “SE” containing immune cells, causing a plethora of proinflammatory responses related to clinical RA disease.¹⁷ The nature of the *Proteus spp.* relationship with RA possibly explains the disproportionate occurrence of disease in middle-aged women, owing to inherent difference in the genitourinary tract resulting in higher frequencies of UTIs in females.¹⁸

Lipopolysaccharides (LPS) are major endotoxin components of gram-negative bacterial cell walls, and act as strong pathogen-associated molecular patterns (PAMPs) to stimulate various innate immune responses.¹⁹ High levels of LPS are present within the gut lumen as a result of numerous residing commensal bacteria, and LPS is normally unable to penetrate across healthy intestinal epithelium.²⁰ However, LPS-mediated inflammation and breach of intestinal permeability occurs when homeostasis of the normal gut microbiome is disturbed.²¹ LPS is therefore able to enter blood circulation and contribute to various systemic inflammatory processes. The disruption of the normal gut microbiota, known as gut dysbiosis, is highly prominent in RA patients and could therefore contribute significantly to the chronic inflammatory state.^{22,23} LPS ligates with toll-like receptor (TLR-4) leading to the production of various proinflammatory cytokines (tumor necrosis factor α [TNF α], interleukin-6

[IL-6], interleukin-8 [IL-8], interleukin-12 [IL-12]) through the nuclear factor kappa B (NF κ B) signal transduction pathway.²⁴ TLR-4 is expressed on various immune cells involved in RA inflammatory signaling.

The high levels of LPS associated with gut dysbiosis, which is common in RA patients, might also influence the hypercoagulable state seen in a high proportion of affected individuals. As discussed previously, LPS stimulate the release of proinflammatory cytokines via TLR-4 dependent pathways. This process has been shown to facilitate the increased expression of tissue factor (TF) by immune cells, thereby amplifying coagulation via the TF pathway.^{25,26} It has also been demonstrated that LPS, due to its highly lipophilic nature, is able to bind to fibrinogen and cause more acute fibrin polymerization.²⁷

The general overview of the key immunopathogenic events in the clinical RA disease development is illustrated in **Fig. 1**. We consider RA pathogenesis within a systems biology approach, providing evidence of bacterial infectious agents generating the autoimmune inflammatory response, while emphasizing the importance of hematological biomarkers of acute systemic inflammation (hypercoagulation and β -amyloid formation) in disease diagnosis and treatment.²⁸

Cardiovascular Complications in Rheumatoid Arthritis

Together with the immunopathogenic profile in RA, there is evidence for increased cardiovascular disease (CVD) risk in these patients.^{29–31} Conventional risk factors (age, sex, obesity, hypertension, and hypercholesterolemia) do not fully account for premature development of CVD in RA, thus indicating a prominent role for chronic systemic inflammation in alterations of hemostatic and thrombotic function.³² Highly sensitive molecular markers of hemostatic function such as prothrombin factor 1 + 2 (PF₁₊₂), thrombin–antithrombin complex (TAT), D-dimer, thrombin-activatable fibrinolysis inhibitor (TAFI), TF, and fibrinogen are significantly elevated in RA patients compared with healthy individuals.^{33–35}

Increased prevalence of an activated platelet profile has been observed in RA patients.³⁶ Platelets have a dual role in affecting both inflammatory and hemostatic responses in RA. Activated platelets express P-selectin which binds soluble CD40 ligand (sCD40L), triggering the expression of inflammatory adhesion receptors, proinflammatory cytokines, and matrix metalloproteinases (MMPs).³⁷ Increased levels of platelet activity markers (sCD40L, P-selectin) are associated with RA disease activity.³⁶ In addition to stimulation of platelets by TNF α , ACPAs are also capable of directly influencing platelet activity through Fc γ -RIIa-dependent binding.³⁶ Some proteins released from platelet granules are associated with formation of more compact fibrin clot structures (e.g., increasing concentrations of platelet factor 4 and factor XIII which have fibrin cross-linking properties).^{38,39}

TF is a transmembrane glycoprotein that functions as a cellular receptor and an important initiator of coagulation.⁴⁰ TF expression is upregulated in the RA synovium and present on key proinflammatory mediators such as fibroblast-like synoviocytes, B-cells, and macrophages.⁴¹ Proinflammatory

Fig. 1 The outline of key events in the immunopathogenesis of rheumatoid arthritis (RA). (Original figure; created with Biorender.com).

cytokines (TNF α , IL-1) facilitate thrombus formation by stimulating the procoagulant activity of endothelial cells, leading to increased TF expression.⁴²

Fibrin fiber generation and structure are abnormally altered in RA as a result of the underlying inflammatory processes.⁴³ Aberrant fibrin deposition in RA plasma as seen by scanning electron microscopy (SEM) is illustrated in ►Fig. 2. The increased levels of fibrin deposition contribute the structural integrity of pannus in RA synovia.⁴⁴ Pannus refers to the abnormal proliferation of synovial tissue lining, composed of stimulated fibroblasts, immune cells, and enriched vasculature.⁴⁵ The presence of pannus formation in the synovium contributes to perpetual stimulation of local inflammation and erosion of extra-articular cartilage and bone.⁴⁵

Levels of hemostasis factors responsible for thrombus generation (fibrinogen, von Willebrand factor, plasminogen-activator inhibitor I) are increased in RA patients.^{31,35} Characteristics of fibrin clot structure in RA include thick, matted layers of fibrin fibers with increased diameter.⁴³ Unusual projections in RA patient fibrin fibers are also present owing to interactions with various proinflammatory mediators.³⁴ These ultrastructural alterations result in fibrin clots that are significantly less stable, permeable, and lysable compared with clots formed under physiological conditions.³⁴

Increased levels of fibrinolysis biomarkers (i.e., D-dimer, PF₁₊₂, TAT) have been associated with RA disease activity and inflammation.^{33,35} However, inflammation also induced TAFI, which favors fibrin persistence and prevents the protein breakdown.³⁵ Previously described ultrastructural alterations also contribute to decreased breakdown of fibrin clots.^{34,43} Citrullinated-fibrinogen immune complexes impair the activity of plasmin, leading to decreased levels of fibrinolysis and sustained synovial damage through proinflammatory effec-

tors.⁴⁶ These findings further support the notion of an impaired hemostatic balance of fibrin clot formation and breakdown in RA patients. CVD risk, inflammation, hypercoagulation, and immunopathology may be closely interlinked in RA pathology. The main immunological hallmarks and diagnostic biomarkers of RA are known to be rheumatoid factor (RF) and ACPAs. The development of autoimmunity in RA may be driven by a bacterial presence that may play important roles in the development of hypercoagulation in RA.

Diagnostic Markers in Rheumatoid Arthritis with Specific Focus on Anti-Citrullinated Protein Antibodies

The presence of two notable autoantibodies in RA patient sera, namely RF and ACPA, is a hallmark characteristic of the disease. RFs were the first identified autoantibodies to be linked with RA and target the Fc portion of immunoglobulin G (IgG).⁴⁷ RF has very low specificity for RA, as it is detected in other rheumatic disorders and in healthy individuals,^{48,49} whereas there is strong evidence for the role of ACPAs in RA immunopathogenesis.⁵⁰ However, the presence of both IgA-RF and ACPA in the preclinical phase is predictive of more rapid disease onset, with increased severity and decreased possibility of remission.⁵¹ The concomitant effect of RF and ACPAs in RA patients stimulates in vivo proinflammatory cytokine production, which increases the risk of significant bone mineral density loss.⁵² Recently, the cross-reactivity of RF with citrullinated fibrinogen, which is distinct from its primary IgG1-Fc target, has been reported.⁵³ This occurs as a result of sequence homology between the Fc and citrullinated fibrinogen epitopes, both of which contain arginine residues.⁵³ These findings could therefore mechanistically explain the

Fig. 2 A series of scanning electron microscopy images illustrating the difference in fibrin clot structure of a healthy control (A) and RA patients (B–D). (Unpublished work by authors).

occurrence of more severe disease severity in RA patients with positive sera for both RFs and ACPAs.

Citrullination is a physiologically occurring process during apoptosis, inflammation, or epithelial keratinization.⁵⁴ The amino-acid arginine is posttranslationally modified by peptidylarginine deaminase (PAD) enzymes to form citrulline, a nonstandard amino acid.⁵⁴ Modification of proteins in this manner results in an increase of molecular mass (1 kDa compared with regular arginine residues)⁵⁵ and a reduction of the net positive charge, causing the loss of potential ionic bonds and interference of hydrogen bonds.⁵⁶ These conformational changes ultimately result in unfolding of protein structure.⁵⁶ Various protein targets for citrullination exist, including human fibrin(ogen),⁵⁷ histones,⁵⁸ vimentin,⁵⁹ α -enolase,⁶⁰ and types I- and II collagen.^{61,62} Citrullination enhances peptide affinity toward MHC molecules, leading to activation of CD4⁺ T-cells.⁶³ ACPAs are then produced in response to the presence of citrullinated peptides or proteins.^{54,64} ACPAs are highly specific to RA,⁶⁵ present prior to clinical disease onset,⁶⁶ and are indicative of more severe disease activity and rapid structural damage.⁶⁷

Generation of Citrullinated Autoantigens

PADs are intracellular, Ca^{2+} -dependent deaminase enzymes that are usually inactive under normal physiological conditions.⁶⁸ PAD enzymes become activated during large-scale apoptotic or inflammatory events as a result of plasma or

organelle membrane disintegration, causing extracellular Ca^{2+} influx or PAD enzyme liberation from cellular compartments.^{69–71} The abnormal humoral response to the presence of citrullinated proteins results from insufficient phagocytic clearance of apoptotic remnants under severe inflammatory conditions.^{70–72} Five mammalian isoforms of PAD are known (PAD1, PAD2, PAD3, PAD4–5, PAD6), of which PAD2 and PAD4 are primarily responsible for the generation of autoantibodies in RA.^{4,73} PAD2 and PAD4 are expressed by several peripheral blood leukocytes present in the RA synovium, in addition to performing regulatory functions related to gene expression, central nervous system development, and tumorigenesis.^{4,71,73,74} PAD2 and PAD4 exhibit different subcellular localization and antigen target specificities in RA and are associated with the presence of inflammatory markers.^{75,76} The most prominent protein targets for PAD-driven citrullination are human fibrin(ogen),⁵⁷ histones,⁵⁸ vimentin,⁵⁹ α -enolase,⁶⁰ and types I- and II collagen.^{61,62}

The primary environmental risk factors associated with RA development are infection⁷⁶ and smoking.⁷⁷ There has also been strong emerging evidence of the role of infectious agents in the initiation and progression of rheumatic diseases. The prevalence of periodontitis in RA patients (35% in all patients and 37% in ACPA⁺-RA patients) is significantly increased compared with individuals not diagnosed with RA.⁷⁸ Periodontitis has been associated with various RA disease markers, such as disease activity scores (DAS-28), C-reactive protein

(CRP), erythrocyte sedimentation rate, and anticyclic citrullinated peptide (CCP) antibodies.^{78–80} Okada et al also demonstrated that clinical disease presentation is improved with periodontal treatment in RA patients.⁸¹ Emerging evidence suggests that RA may develop at extra-articular mucosal sites (lungs and oral cavities), where exposure of HLA-DR β 1 risk genes to environmental sources of inflammation (tobacco smoke and bacterial or viral infection) is most likely to occur.⁷⁶ This results in the generation of an aggressive autoimmune response and subsequent assault of self-tissues.⁷⁶

Anti-Citrullinated Autoantibodies

Synovial fluid-derived B cells are responsible for generating ACPAs in RA.⁸² Varying degrees of ACPA cross-reactivity have been demonstrated, ranging from monoreactive antibodies to reactivity with multiple citrullinated antigens.^{83,84} Li et al indicated that 66.7% of plasmablast-derived ACPAs from RA patients demonstrated cross-reactivity with different citrullinated epitopes, while 33% of the patients had monoreactive ACPA subsets that recognize either citrullinated fibrinogen, citrullinated enolase, or citrullinated vimentin.⁸⁵ The mode in which ACPAs target citrulline-containing peptide antigens is dependent on appropriate molecular motifs, specifically sequences in which citrulline residues are flanked by other small and neutral amino acids.⁸⁶

In the preclinical stage of RA development, increased ACPA titers are accompanied by increased levels of epitope spreading to recognize a broader range of citrullinated antigens.^{87,88} However, no further expansion of epitope recognition is observed after disease onset.⁸⁹ The presence and level of antigen recognition by ACPAs in the RA synovium determine the magnitude of the immune response.⁹⁰ The magnitude of the patient ACPA isotype profile is also strongly predictive of long-term radiographic progression.⁹¹

The presence of ACPAs represent a specified subgroup that constitutes 70 to 80% of all RA patients.^{74,75} These observations indicate a strong role of ACPAs in development of clinical RA, ACPAs are therefore listed as an important criterion for RA disease classification in American College of Rheumatology/European League against Rheumatism (ACR-EULAR) guidelines.⁹² The discovery of the first citrullinated antigens (flag-grin) and ACPAs (antikeratin factor and antiperinuclear factor) in RA^{93,94} led to the development of enzyme-linked immunosorbent assay (ELISA)-based methods for detecting ACPAs.⁹⁵ Known as anti-CCP tests, these detection methods are considered the gold standard for detection of ACPA presence in RA patients. First generation anti-CCP tests displayed 68% sensitivity and 97 to 98% specificity for RA diagnosis,⁹⁵ with improved results in second generation anti-CCP2 tests (80% sensitivity and 98% specificity).⁹⁶ However, anti-CCP detection captures overlapping ACPA cross-reactivity and therefore cannot determine single ACPA-subtype specificity.⁸³

Fibrinogen in Rheumatoid Arthritis

Fibrinogen is a large (340 kDa) glycoprotein that is synthesized in the liver and circulates in blood at concentrations between 2 and 4 g/L.⁹⁷ The triglobular structure consists of paired α -, β -, and γ -polypeptide chains linked by disulfide bonds via their N-

terminal regions.^{97–99} Fibrinogen is the soluble precursor of insoluble fibrin, the terminal component of the common coagulation pathway with thrombin responsible for the proteolysis of fibrinogen.^{98,100} Thrombin, a trypsin-like serine protease, cleaves fibrinogen between Arg-16 and Gly-17 of the A α chain and Arg-14 and Gly-14 of the B β chain to expose polymerization sites, resulting in release of fibrinopeptide A and fibrinopeptide from their respective N-terminal regions.^{97,98} Fibrin monomers polymerize spontaneously, mediated by activated factor XIII (FXIIIa) that cross-link individual fibrin molecules to form a complex, stable, and insoluble branching network of fibrin fibers.⁹⁷ In the event of vascular injury, fibrinogen is digested by thrombin, and generates clots consisting of dense fibrin fiber networks.^{43,98} During systemic inflammatory states, characteristic of clinical RA, excessive fibrinogen generation and degradation occur.¹⁰¹

The role that fibrinogen and fibrin (in both native and citrullinated forms) play in RA disease development and progression will be extensively discussed further in this review, with specific focus on cardiovascular disease as a secondary risk in RA patients. Fibrinogen fulfils a unique role in rheumatic disease as it not only is involved in the conventional interplay between chronic local and systemic inflammation with dysregulated coagulation, but also actively contributes to the development of the autoimmune response. Insights into the role that fibrinogen/fibrin plays at each level of disease progression may provide the necessary knowledge to identify novel therapeutic targets and treatment strategies to alleviate severe disease symptoms.

As mentioned previously, there is a strong genetic link associated with RA, where the HLA-DR β 1 gene is the most well-known implicated gene.^{9,10} The extent to which citrullinated fibrinogen peptides may influence HLA-DR β 1-mediated immune responses in RA remains unclear. Several studies observed no significant T-cell response or arthritogenic induction to citrullinated-fibrinogen immunization of HLA-DR β 1 transgenic mice.^{100,102,103} However, the presence of antibodies to citrullinated fibrinogen (ACF) is associated with human HLA-DR β 1-0404* allele along with early disease onset and radiographic progression.¹⁰⁴ The arthritogenic potential of citrullinated-fibrinogen in HLA-DR4 transgenic mice has also been reported by Rajmakers et al.⁵⁵ Possible reasons for these discrepancies include increased ACPA heterogeneity in human RA, increased number of sites available for citrullination on human fibrinogen, and differences between the T-cell repertoire of humans and murine models.¹⁰³ The humoral response in murine models is also dependent on the strain, immunogen, and the immunization protocol used.¹⁰² Citrullinated fibrinogen has also been associated with the presence of protein tyrosine phosphatase, non-receptor type 22 (PTPN22) R620W risk alleles.¹⁰⁰ PTPN22 promotes the survival of B-cells in RA and other inflammatory conditions, such as type 2 diabetes, by evading tolerance checkpoints.^{105–107}

Fibrinogen and Fibrin Are Endogenous Sources of RA Autoantigens

As described previously, fibrin and fibrinogen are major targets for PAD citrullination and could therefore play a role

in the immunopathogenesis of RA.^{55,57,108} RA is characterized by excessive generation and breakdown of fibrinogen, in addition to deposition of fibrin within the inflamed RA synovium, even in cases of well-controlled disease.^{109,110} Increased plasma fibrinogen levels in RA have been reported to positively associate with disease activity scores and acute-phase markers such as CRP and serum amyloid A (SAA).¹¹⁰ The presence of immune complexes containing citrullinated fibrin(ogen) and autoantibodies directed toward the citrullinated forms of fibrin(ogen) has been confirmed in human RA sera and synovial fluid in various studies.^{55,57,100,101,108} The presence of circulating immune complexes containing citrullinated fibrinogen has been reported in half of anti-CCP⁺ RA patients in a study by Zhao et al.¹¹¹ ACF antibodies have predominantly nonoverlapping reactivity, and target citrullinated fibrinogen peptides more favorably than native fibrinogen.¹⁰⁰ ACF-ELISA detection tests display similar performance to commercial anti-CCP2 assays and are highly predictive of rapid radiographic progression in terms of RA specificity.^{112,113} The detection of both citrullinated fibrinogen antigens and autoantibodies directed toward such peptides, along with reported findings of arthritis development in certain murine models when immunized with both native and citrullinated forms of fibrinogen,^{102,114} strongly suggests a pathogenic role for fibrinogen and fibrin in human RA.

Deamination of arginine residues to citrulline leads to an increase in nominal molecular mass of 1 kDa per altered amino acid residue.⁵⁵ This leads to an overall molecular mass increase of 31 kDa in the α fibrinogen subunit and 21 kDa in the β fibrinogen subunit, respectively.⁹⁷ Two-thirds of arginine residues in fibrinogen are susceptible to citrullination by both PAD2 and PAD4 isoforms, with the most prominent citrullination sites located on the α (highest amount of target sites) and β fibrinogen subunits.^{46,97,115} Fibrinogen γ -chains and fibrinogen-derived degradation products are not significantly targeted for citrullination by PAD enzymes.⁹⁷ PAD2 and PAD4 enzymes recognize different epitopes for citrullination in fibrinogen and fibrin peptide chains.¹¹⁶ Studies have demonstrated that PAD2 is more efficient in deamination of arginine residues in fibrinogen and fibrin chains⁷⁵ and induces a significantly higher proinflammatory response¹¹⁷ than PAD4. The enhanced efficiency of PAD2 for deaminating arginine residues is due to the enzyme being less restricted by the amino acid composition surrounding acceptor arginine residues in fibrinogen and fibrin compared with PAD4.⁷⁵ There is no significant discrimination of ACF-directed reactivity toward citrullinated peptides generated by either PAD2 or PAD4, as there is a great degree of overlap in the citrullination sites targeted by both PAD2 and PAD4.⁷⁵ The mapping of citrullinated epitopes in fibrinogen and fibrin has been conducted by a large number of studies. The identification of 42 citrullinated residues in fibrinogen (27 in the α -chain, 11 in the β -chain and 4 in the γ -chain) currently represents the most comprehensive epitope list for citrullinated fibrinogen compiled thus far.¹¹⁵ Fibrin contains a decreased amount of susceptible arginine residues for citrullination compared with soluble fibrinogen.^{4,46} The major citrullination epitopes contained in fibrin have been reported as α _{36–50} and β _{60–74}, respectively.^{46,57}

Despite evidence of significantly increased levels of fibrin clot formation and fibrinolysis in various studies, limited research regarding the proteolysis of citrullinated fibrinogen peptides has been conducted. Paradoxical observations regarding the involvement of citrullinated fibrinogen in the thrombin-catalyzed fibrin polymerization reaction have been made in some literature. Studies by Nakayama-Hamada et al and Okumura et al found that citrullinated fibrinogen peptides are resistant to thrombin digestion under reverse phase high-performance liquid chromatography (HPLC) conditions.^{97,98} The inhibition of thrombin-catalyzed fibrinogen polymerization most likely occurs due to the citrullination of arginine residues within the thrombin cleavage sites in α and β fibrinogen chains, preventing thrombin binding.^{97,98,118} SEM analysis by Okumura et al of fibrin clot structure comparing samples containing native fibrinogen and native fibrinogen with citrullinated fibrinogen in combination indicated no significant difference in terms of fibrin fiber density and diameter between samples.⁹⁷ These findings confirm that citrullinated-fibrinogen does not participate in thrombin-catalyzed fibrin clot formation. Citrullinated fibrinogen could therefore be regarded as an uncompetitive inhibitor of the thrombin reaction in the coagulation process.⁹⁸ This impairment may suggest an effect of citrullination on the hemostatic balance in RA patients, but this directly conflicts with multiple observations of excessive fibrinogenesis and fibrinolysis in previous studies. Possibly explaining this phenomenon is that excessive levels of thrombin in RA synovial fluid increase the likelihood that high amount of fibrinogen is polymerized within the inflamed synovium, despite the concurrent presence of deaminating enzymes.^{35,119} The reduced fibrinolytic ability of plasmin has also been observed following fibrin citrullination.¹²⁰ Plasmin is a serine proteinase that cleaves peptide bonds near basic amino acid residues such as arginine, and therefore deamination will reduce the amount of potential degradation sites.¹²⁰

Possible Interaction between Protein Deamination and Amyloidogenic Clot Formation

As discussed earlier in this review, the deamination of arginine residues in plasma proteins such as fibrinogen is observable in RA patient sera and brings about a change in structural conformation. Arginine residues contribute to tertiary protein structure by forming noncovalent intramolecular bonds with acidic residues.¹²¹ Citrulline lacks the polarity of arginine, and the conversion of arginine to citrulline by PADs will therefore disrupt key structural features within protein molecules (**►Fig. 3**).¹²¹ It has been reported that 54 of the 81 arginine residues (66%) within human fibrinogen are susceptible to deamination by either PAD2 or PAD4,¹¹⁵ indicating that substantial protein structural modification could take place. The normal function of fibrinogen would inevitably also be altered, as discussed previously in the form of autoantibody-driven inflammatory responses and disrupted fibrin polymerization and degradation. The major sites of the fibrinogen molecule affected by citrullination are the paired α , β , and γ peptide chains within the D-regions that flank the central nodule of the

Fig. 3 Comparison of protein structure modification during citrullination and amyloidogenesis. Deamination of polar arginine residues interferes with intramolecular bonds, causing potential unfolding of tertiary protein structure. Under increasing mechanical force or pathological stress, α -helices transition to β -sheets in fibrin, resulting in a pathologically altered clot. (Original figure; created using Biorender.com).

fibrinogen molecule.¹¹⁵ Paired structures within the fibrinogen molecule are held in place by 29 disulfide bridges formed between cysteine amino acid residues.¹²² These individual peptide chains consist of an α -helical secondary protein structure, that wind around each other to form an α -helical coiled-coil structure.¹²³ The α -helical coiled-coil conformation comprises approximately 25% of the amino acid architecture of the entire fibrinogen molecule.¹²⁴ The normal elastic and plastic properties of fibrin(ogen) are highly dependent on the ability of coiled-coils to unwind, and the conversion of α -helical coiled-coils to a β -sheet formation under mechanical stress has been demonstrated.^{125,126} Staining of these β -sheet structures in mechanically altered fibrin(ogen) by Congo-red revealed that these structures closely resemble the β -sheet aggregation seen in amyloid proteins.¹²⁷

The relevant biophysical definition of amyloid is described as "unbranched protein fibers whose repeating substructure consists of β strands that run perpendicular to the fiber axis, forming a cross β -sheet of undefined length."¹²⁸ Amyloid proteins have been traditionally linked with conditions such as Alzheimer's disease (deposition of amyloid- β in senile plaques) and type 2 diabetes (islet amyloid polypeptide).¹²⁹ Amyloidosis are a category of protein-misfolding disorders in which soluble proteins undergo conformational changes and are deposited in extracellular locations as aberrant, insoluble fibrillar structures that disrupt tissue structure and function.¹³⁰ Fibrinogen is not generally considered amyloidogenic, except in rare cases of mutations in the *fibrinogen A α* genes, which seem to be the most common cause of hereditary renal amyloidosis.¹³¹ However, our group has demonstrated anomalous fibrin clot formation that is amyloid in nature in several

inflammatory diseases.^{132–134} The same aberrant clot formation has also been demonstrated in a purified fibrinogen model in the presence of certain inflammatory stimuli such as bacterial LPS^{27,135} and SAA.¹³⁶

It therefore begs the question whether any interaction may occur between the processes of citrullination and amyloid transformation of fibrin(ogen) as outlined in **Fig. 3**, and if it could play a major role in disease outcome in RA. This notion is substantiated by the occurrence of abnormal clot formation and presence of bacterial LPS from a dormant blood microbiome (seen as causal agents of amyloid fibrin formation) that seems intrinsically related to RA as well. However, limited research has been conducted with regards to the direct impact these two protein modification processes have upon each other. Osaki and Hiramatsu conducted research on the effect of citrullination on the single arginine residue conserved in the two isoforms of amyloid β protein ($A\beta_{1-40}$ and $A\beta_{1-42}$).¹³⁷ $A\beta$ is the major component of insoluble amyloid plaques in Alzheimer's disease (AD) neural tissue, with the 42-peptide isoform considered the most neurotoxic. The soluble, pre-fibrillar amyloid- β oligomers are also implicated in causing significant neurotoxicity and cytotoxicity in AD.¹³⁸ Citrullination of $A\beta_{1-40}$ resulted in an increased fraction of soluble amyloid oligomers, and citrullination of $A\beta_{1-42}$ caused in an increased fraction of β -sheet rich oligomers.¹³⁷ These findings suggest that citrullination causes structural changes bearing implication for the aggregative nature of amyloid proteins and providing some compelling additional research questions. Could citrullination of fibrin(ogen) α -helical coiled-coil regions that result in tertiary structural changes create favorable conditions for conversion to pathogenic, amyloid-like

β -sheet structures by LPS and SAA molecules? Conversely, could fibrin clots that become amyloid in nature provide suitable epitopes for PAD enzymes, thus resulting in enhanced autoantibody driven inflammatory responses? Further investigation to assess the extent that the processes of citrullination and amyloidogenesis affect each other could further enhance current knowledge on the nature of aberrant fibrin clot formation seen in RA.

Acute Phase Reaction

The acute phase response is highly prominent in RA as both a result and a contributor to the chronic inflammatory state in affected individuals. Acute SAA (A-SAA) is of particular significance, as it is a highly sensitive marker of acute inflammation (increases 1,000-fold above normal) and is therefore highly elevated in RA patients, even within individuals that present with improved clinical symptoms.^{139–142} A-SAA is the circulating precursor of amyloid A protein, a fibrillar, insoluble apolipoprotein that is deposited in major organs (amyloidosis) leading to the severe risk of organ damage and possible death.^{143–145} Despite being primarily produced by the liver, extrahepatic sources of SAA have also been identified.¹⁴⁶ A-SAA is expressed by synovial fibroblasts,^{146,147} monocytes/macrophages,^{146,148} and endothelial cells.¹⁴⁶ A-SAA synthesis is also stimulated by various proinflammatory cytokines such as IL-6, IL-1, TNF- α , and IFN- γ .¹⁴² A-SAA is also able to induce several proinflammatory processes contributing to RA immunopathogenesis. A-SAA promotes the expression of vascular cell adhesion molecule 1 (VCAM-1) and intercellular adhesion molecule-1 (ICAM-1) and the adhesion of peripheral blood mononuclear cells to synovial fibroblasts and thus accelerating pannus formation.¹⁴⁹ A-SAA also mediates the production of MMPs by synovial fibroblasts, thus contributing to joint destruction.¹⁴⁹

Recently our group has demonstrated the ability of SAA to bind to fibrin(ogen), and that relatively low concentrations of SAA-induced amyloid fibrillation of fibrin clots.¹³⁶ Amyloidogenic formation of proteins occurs when α -helices uncoil resulting in β -sheet structure¹²⁵ with misfolded proteins leading to an increased thrombotic tendency.¹⁵⁰ With previous reported findings of significantly elevated levels of A-SAA in RA patients, including patients with improved clinical symptoms, A-SAA may significantly contribute to coagulopathy seen in RA individuals and should thus be closely monitored in clinical settings. LPS-mediated β -amyloid formation of fibrin networks through the ability of LPS to bind human fibrin(ogen) has also been demonstrated.²⁷

Involvement of Bacterial Components in Immunopathogenesis and Hemostatic Complications of Rheumatoid Arthritis

Porphyromonas gingivalis as Environmental Source of (Fibrinogen) Citrullination

As mentioned in the introduction, gingivitis and periodontitis are now increasingly implicated in development and progression of RA. *Porphyromonas gingivalis*, the major etiological cause of periodontitis, has received significant

attention as a possible risk exposure for RA development. *P. gingivalis* is a gram-negative, asaccharolytic, and highly proteolytic bacterium,^{151,152} and is the only known prokaryote to produce deaminating enzymes similar to that of human isoforms, termed *P. gingivalis* peptidylarginine deaminase (PPAD).¹⁵² Human PAD enzymes, particularly PAD2 and PAD4, are the major sources of citrullinated antigens that drive the development of RA autoimmunity. It is therefore suggested that an external source of citrullination through bacterial PPAD activity could significantly contribute to the generation of RA-specific autoantibodies. Previous studies have demonstrated the ability of PPAD to citrullinate conventional RA-specific self-peptide targets, such as human fibrin(ogen), vimentin, and α -enolase.^{153,154} The mode of citrullination by PPAD is distinct from human PAD isoforms in that it targets C-terminal arginine residues for deamination in proteins, whereas human PAD preferentially deaminates internal arginine residues.¹⁵⁵ PPAD is also not Ca^{2+} dependent like human PAD isoforms.¹⁵⁶

The major bacterial co-virulence factor of PPAD secreted by *P. gingivalis* is a class of proteolytic enzymes termed gingipains. Gingipains catalyze the cleavage of proteins or peptides at either lysine or arginine residues.¹⁵⁵ Arginine-specific gingipains (RgpA or RgpB) yield suitable peptide fragments with C-terminal Arginine residues for citrullination by PPAD.^{155–157} RgpB has been shown to effectively cleave arginine residues in fibrinogen, with PPAD able to execute citrullination of the majority of resulting peptide fragments.¹⁵⁸ PPAD and Rgp are both expressed on the outer membrane of *P. gingivalis*, and operate in concert to elicit autoimmune inflammatory responses in the human host.^{154,156}

Positive correlations between serum levels of anti-PPAD antibodies with CRP and IL-6 indicate a role of *P. gingivalis* infection in producing systemic inflammation.⁸⁰ Yamakawa et al demonstrated that *P. gingivalis* infection of the spontaneous keratoconus (SKC) mouse model produces rapid RA development, with significant increases in arthritis score, APCAs, proinflammatory signaling molecules, and a decrease in bone density observed.¹⁵⁹ PPAD citrullination of histone proteins aids *P. gingivalis* in evading neutrophil extracellular traps (NETs).¹⁵² *P. gingivalis* in turn can induce NETosis.¹⁶⁰ The development of arthritis through *P. gingivalis*-induced inflammatory lesions has been demonstrated in an animal model.¹⁶¹

The activity of *P. gingivalis* and its virulence factors may also contribute to the increased cardiovascular disease risk in RA and Parkinson's disease (PD) patients.^{162,163} It has been demonstrated that Arginine-specific gingipains (RgpA, RgpB) are able to activate the coagulation cascade at multiple levels. Gingipains proteolytically stimulate coagulation factors IX,¹⁶⁴ X,¹⁶⁵ and II (prothrombin).¹⁶⁶ However, unstable fibrin clots are formed as a result of concurrent fibrinogen degradation by lysine-gingipains (Kgp).¹⁶⁷ The proteolytic activities of the virulence factors of *P. gingivalis* have been shown to degrade apolipoproteins essential for normal lipoprotein metabolism, producing proinflammatory and proarthritogenic events that could account for increased risk of atherosclerosis.¹⁶² The proteolysis of human fibrinogen through the combined effect of PPAD and RgpB inhibits

fibrin polymerization, which results in localized bleeding and a potentially dysregulated coagulation cascade.¹⁶⁸ Rgp is able to activate platelets via platelet activation receptors one (PAR-1) and four (PAR-4).¹⁶⁹ Activation of PARs by gingipains is similar to platelet activation by thrombin, as gingipains target N-terminal regions of PARs that contain cleavage sites for thrombin.¹⁵¹

Recently, LPS and gingipains from *P. gingivalis* were also shown to cause amyloid formation in purified fibrinogen.¹⁷⁰ The presence of gingipains in whole blood clots as detected by a fluorescent anti-RgpA antibody was confirmed in a PD patient population, and this was associated with increased levels of circulating proinflammatory cytokines, hypercoagulation, amyloid fibrin formation, and altered platelet ultrastruc-

ture.¹⁷⁰ These findings could therefore also be observable within RA patient sera, as the presence of *P. gingivalis* and its associated gingipains has a long-established relationship with disease as well. ► **Fig. 4** illustrates the human coagulation cascade, indicating the effects RA-specific inflammatory and autoimmune events may have on overall hemostatic outcome.

RA Inflammatory Pathways: Interactions between Citrullinated Fibrinogen and Prominent Immune Cells

The chronic inflammatory state that is a hallmark of RA is characterized by an accumulation of inflammatory leukocytes into the synovium and articular spaces.^{54,171} The

Chart key

PPAD: *Porphyromonas gingivalis* peptidylarginine deaminase **PAD:** peptidylarginine deaminase

B: B lymphocyte **Mφ:** Macrophage **FLS:** Fibroblast-like synoviocyte **LPS:** lipopolysaccharide

————— Physiological hemostasis reactions

----- RA-specific interferences with hemostasis

Fig. 4 Chronic synovitis and invasive pannus formation, characteristic of clinical rheumatoid arthritis (RA), may result in the amplification of coagulation by increased stimulation of clotting components in both the contact pathway (heightened activated platelet phenotype) and tissue factor pathway (enhanced tissue factor expression by immune cells). Insoluble fibrin in return acts as a structural scaffold agent for pannus formation in RA synovial joints. Activation of coagulation may also be proteolytically propagated by *Porphyromonas gingivalis* virulence factors (gingipains) at various levels. However, previous studies have demonstrated that citrullinated fibrinogen is not able to undergo thrombin digestion and subsequent fibrin polymerization. It has also been shown that LPS from gram-negative bacteria (such as *P. gingivalis*) is able to inflict fibrin alteration that are amyloid in nature, resulting in structurally unstable clots. (Original figure; created with Biorender.com).

cellular composition of inflamed RA synovium consists of immune cells from both innate and humoral systems, fibroblast-like synoviocytes, chondrocytes, and bone-remodeling cells such as osteoclasts and osteoblasts.⁵⁴

The RA autoimmune response is likely elicited due to an abnormal humoral response to the presence of citrullinated proteins in affected individuals.¹⁷² The major genetic risk factor for RA ACPA generation, HLA-DR β 1 “shared epitope” alleles, exerts major influence on $\alpha\beta$ CD4⁺ T-cell selection.^{11,173} The altered CD4⁺ T-cell repertoire is responsible for inflammatory signaling and generation of anticitrullinated autoantibodies by synovial B-cells through the self-presentation of citrullinated proteins by MHC molecules (►Fig. 5).¹¹ Deaminating enzymes responsible for generation of citrullinated antigens (PAD2 and PAD4) are also expressed by inflammatory leukocytes within the synovial microenvironment, which include synovial T and B cells, macrophages, neutrophils, and fibroblast-like synoviocytes.^{174,175}

Human fibrin(ogen) is capable of directly stimulating a variety of leukocyte activities (►Fig. 6). Fibrin(ogen) engagement with macrophages via TLR-4 and other integrins induces expression of proinflammatory chemokines,¹⁷⁶ and cytokines such as TNF- α .¹⁷⁷ Fibrinogen stimulates the proinflammatory activity of neutrophils by binding to the α -subunit of CD11b/CD18.¹⁷⁸ Fibrinogen regulates NF κ B signaling and inflammatory chemokine expression of endothelial cells.¹⁷⁹ Human synovial fibroblasts express ICAM-1 and inflammatory chemokines under the influence of fibrinogen.¹⁸⁰ These findings indicate that fibrin(ogen), which is present in high amounts within RA articular spaces, may fulfil a pronounced role in mediating various RA-related inflammatory processes. Fibrin displays even stronger involvement in propagation of inflammatory processes in RA than its soluble precursor.¹¹⁷ Fibrin deposits are present within the RA synovial tissue lining, and thus enable the proliferation of RA synovial fibroblasts and local cytokine release.^{44,181} The insoluble nature of the protein increases the difficulty of its clearance from articular spaces, thus leading to increased likelihood of posttranslational modifications and generation of an autoimmune inflammatory response.¹¹⁷

Macrophages/Monocytes

ACPAs directly contribute to initiating and sustaining the state of synovial inflammation by their ability to activate monocytes and macrophages, to produce inflammatory cytokines.¹⁸² Citrullinated-fibrinogen containing immune complexes stimulate TNF α secretion by macrophages through Fc γ R-TLR4 engagement.^{108,183} The presence of ACPAs causes a dysregulated M1/M2 macrophage subset polarization, favoring M1 (classical) activation.¹⁸⁴ The M1 subset is characterized by its inflammatory, microbicidal, and tumoricidal properties.¹⁸⁴ There is an overexpression of TLR2 and TLR4 receptors by macrophages and synovial fibroblasts in RA.¹¹⁷ TLRs recognize disease-associated molecular patterns (DAMPs) and PAMPs, resulting in innate immune responses.¹⁸⁵ PAMPs are microbial-derived molecules such as peptidoglycan, bacterial and viral deoxyribonucleic acid (DNA) that are structurally distinguish-

able from host molecules.^{185,186} DAMPs are derived from clinical symptoms associated with the pathophysiological responses, with fibrinogen/fibrin and dsRNA from apoptotic cells present in the RA joint serving as DAMPs.^{185,186}

RA Synovial Fibroblasts

The porous structural arrangement of the synovial lining that lacks epithelial cells, basement membranes, tight junctions, or desmosomes allows accumulation of immune complexes and bacterial cell wall components within rheumatic joints.¹⁸⁷ The synovial intimal lining layer consists of two distinct cell types—macrophage-like synoviocytes (Type A) and fibroblast-like synoviocytes (Type B).¹⁸⁷ The inflammatory milieu of the RA synovium transforms the surrounding tissues to become increasingly hyperplastic, invasive, and immunogenic.¹⁸⁷ Transcription factor NF κ B is ubiquitously expressed in monocytes and synovial fibroblasts, and regulates the expression of proinflammatory cytokines, chemokines, and adhesion molecules in RA.¹⁸⁸ There is an overexpression of TLRs in RA synovial fibroblasts (RASFs), with both fibrin deposits in the RA joint and soluble fibrinogen acting as candidate TLR ligands.^{44,186} TLR ligation potentiates the inflammatory cytokine production of RASFs, with prominent secreted mediators including IL-6, IL-8, TNF α , MMPs, and vascular endothelial growth factor (VEGF).¹⁸⁶ Citrullination of peptides further amplifies the proinflammatory potential of RASFs, with a significant increase in cytokine secretion.¹¹⁷

Tumor Necrosis Factor: Apical RA Cytokine

TNF α is considered the most prominent cytokine involved in the RA inflammatory response, as it in turn stimulates further monocyte and macrophage activity and proinflammatory cytokine release.^{189–191} TNF α has been demonstrated to be increased by 62% in RA patient sera compared with controls.¹⁹² The suggested importance of TNF α in RA pathogenesis has been highlighted by the relative efficacy of TNF α inhibitor (infliximab) treatment in the attenuation of disease symptoms and synovial inflammation.^{193,194} TNF α is produced by T cells within the synovial lining, local macrophages, and monocytes resulting in chronic synovitis and activates osteoclasts and MMPs leading to cartilage and bone destruction.^{195,196} TNF α is also able to activate platelets through the arachidonic pathway, with platelets further propagating RA synovial inflammation and the associated risk of thrombotic cardiovascular events.¹⁹⁷ The chronic exposure of synovial T-cells to TNF α leads to decreased immune tolerance and unresponsiveness to TNF α inhibitor treatment (prevalent in 30–40% of RA patients).¹⁹¹

Neutrophils and NETosis

The presence of hyperactive neutrophils in RA synovial fluid and pannus tissue, in particular NETosis (formation of neutrophil extracellular traps or NETs), has been confirmed by multiple studies.^{198–200} NETs are chromatin-derived extracellular snares that are externalized from neutrophils in response to microbial infections or inflammation.²⁰¹ Neutrophils and NETosis are key components in the perpetual inflammatory response encountered in RA patients.

Chart key

B: B lymphocyte **T_h :** Naïve T-helper lymphocyte **DC:** Dendritic cell
M ϕ : Macrophage **NET:** Neutrophil extracellular trap **PC:** Plasma cell
FLS: Fibroblast-like synoviocyte **OC:** Osteoclast

Fig. 5 The rheumatoid arthritis (RA) "shared epitope" risk gene results in citrullinated proteins being presented as self-antigens via MHC-II complexes to CD4⁺ T-helper cells, resulting in a breakdown of immune tolerance to the presence of citrullinated proteins, with T-lymphocyte differentiation resulting in the production of RA-specific autoantibodies (ACPAs and RFs), as well as direct stimulation of inflammatory effector cells via cytokine signaling. ACPAs, anti-citrullinated protein antibodies; MHC, major histocompatibility complex; RF, rheumatoid factor (Original figure; created with Biorender.com).

Fig. 6 The prominent cell surface receptors recognized as ligating with human fibrinogen, citrullinated fibrinogen, and citrullinated-fibrinogen immune complexes and their downstream signaling pathways are summarized. (Original figure; created with Biorender.com).

Neutrophils express multiple cell surface receptors, and NETosis is induced by proinflammatory mediators such as bacterial LPS,²⁰² platelet TLR-4,²⁰³ and cytokines such as TNFα, IL-1β, and IL-8.²⁰⁴ Enhanced NETosis is associated with local and peripheral presence of ACPAs, with NETs prominent sources of citrullinated autoantigens that express high levels of PAD2 and PAD4 enzymes.²⁰⁵ NETs along with their individual components (DNA and histones) have also been shown to possess significant prothrombotic and antifibrinolytic properties, and have been discussed at length in a recent review by Varjú and Kolev.²⁰⁶ NETs provide a platform for the accumulation of integral hemostatic components such as platelets, erythrocytes, von Willebrand factor, and fibrinogen.²⁰⁷ Histones promote fibrin deposition through its ability to bind to and inactivate antithrombin.²⁰⁸ This protective mechanism against the inhibitory effects of antithrombin is also achieved by the citrullination of antithrombin by PAD4.²⁰⁹ Integration of DNA–histone complexes into fibrin networks results in thicker fibrin fibers that consequently improve clot stability and rigidity but decreases clot permeability.^{208,210} The presence of these NET-derived complexes within the fibrin network also impedes fibrinolysis, through the ability of both histones and DNA to bind large fibrin

degradation products (such as plasminogen) and inhibit its activity.^{208,210}

Conclusion

In this review, we have discussed in detail the role that specific plasma proteins play at every stage of RA immunopathogenic progression. All major themes and the clinical relevance considered in this review are summarized in **Table 1**. Endogenous and exogenous sources of protein citrullination, coupled with the genetic propensity to present citrullinated antigens to CD4⁺ T-lymphocytes present the most convincing current paradigm to explain the development of RA autoimmunity. The insoluble clotting protein fibrin, along with its soluble precursor fibrinogen, has been demonstrated as prime targets for PAD citrullination and the generation of fibrin(ogen)-specific autoantibodies in RA. The excessive rate of fibrin generation and breakdown in the RA synovium actively contributes to RA pannus structure and inflammatory signaling. Both native and citrullinated forms of fibrin(ogen) have been shown to interact with and regulate the proinflammatory activities of various immune cells in RA, including macrophages,

Table 1 Key points of this review

<ul style="list-style-type: none"> • To date, a combination of genetic risk, the presence of infectious agents, including those that become dormant within the host, along with certain environmental exposures present the most plausible etiological explanation for the development of clinical rheumatoid arthritis (RA) disease.
<ul style="list-style-type: none"> • Ebringer's theory (with experiments) states that <i>Proteus</i> infection from the urinary tract is a major cause of RA. Also, the periodontal pathogen <i>Porphyromonas gingivalis</i> possibly presents a major exogenous contributor to RA autoimmune generation through the expression of its unique virulence factors.
<ul style="list-style-type: none"> • In addition to destructive synovitis, a high risk of cardiovascular disease as a result of dysregulated thromboinflammatory function is well established in RA patients.
<ul style="list-style-type: none"> • The presence of autoantibodies directed toward citrullinated proteins (ACPAs) is highly correlated with disease development and influences various proinflammatory processes.
<ul style="list-style-type: none"> • Human fibrin(ogen), a key component of the coagulation process, is a prime candidate ligand for citrullination by peptidylarginine deaminase (PAD) enzymes.
<ul style="list-style-type: none"> • Further research is necessary to define the involvement of citrullination with regards to the formation of abnormal fibrin clots observed in RA patients.
<i>Clinical relevance</i>
<ul style="list-style-type: none"> • Infections of certain mucosal tissues and bacterial components are important factors for consideration in the possible prevention, diagnosis, and treatment of RA.
<ul style="list-style-type: none"> • Acute phase proteins, such as fibrinogen and serum amyloid A, present significant clinical value due to their involvement at various levels of disease development.
<ul style="list-style-type: none"> • By monitoring highly sensitive markers of thrombotic function in RA patients through an individualized medicine approach may represent an improved diagnostic and treatment strategy for systemic inflammation and secondary cardiovascular disease.

synovial fibroblasts, and neutrophils. The increased CVD risk in RA is independent of traditional risk factors, and the interactions between dysregulated inflammatory and hemostatic processes present a major contribution to this secondary disease complication. The extent to which the protein modification processes of citrullination and amyloidogenesis of fibrinogen may interact and influence the other represents an area of novel research that could explain the presence of excessive and aberrant clot formation in RA patients. β -amyloid formation of fibrin fibers as a result of increased A-SAA levels and bacterial LPS constitutes additional coagulopathic complications in RA patients. Despite observations of increased thrombotic tendency coupled with structurally abnormal fibrin clot formation in RA patients, the exact role that citrullinated proteins fulfil in this regard remains controversial and requires further investigation. The contribution of bacterial LPS to the generation of autoantibodies in RA is also yet to be determined. Whether LPS could directly stimulate increased rates of protein citrullination as a result of perhaps autocrine TNF α -induced apoptosis of immune cells,²¹¹ or through indirect mechanisms certainly warrants further investigation. Furthermore, it is yet to be determined how free LPS in circulation might interact directly with plasma proteins like fibrinogen which can lead to citrullination, amyloid formation, and anomalous blood clot formation.

To conclude, it is suggested that specific proteins involved in chronic synovitis and thrombotic tendency in RA patients, such as fibrin(ogen) and amyloid A proteins be considered more intensively in research. The extensive signaling processes these proteins are involved with regards to disease

development may present novel areas for identifying potential diagnostic biomarkers and therapeutic interventions.

Conflicts of Interest

None.

References

- 1 Scott DL, Wolfe F, Huizinga TW. Rheumatoid arthritis. *Lancet* 2010;376(9746):1094–1108
- 2 Silman AJ. Epidemiology of rheumatoid arthritis. *APMIS* 1994; 102(10):721–728
- 3 Tedeschi SK, Bermas B, Costenbader KH. Sexual disparities in the incidence and course of SLE and RA. *Clin Immunol* 2013;149(02): 211–218
- 4 Pratesi F, Panza F, Paolini I, et al. Fingerprinting of anti-citrullinated protein antibodies (ACPA): specificity, isotypes and subclasses. *Lupus* 2015;24(4-5):433–441
- 5 van der Woude D, Catrina AI. HLA and anti-citrullinated protein antibodies: building blocks in RA. *Best Pract Res Clin Rheumatol* 2015;29(06):692–705
- 6 Smolen JS, Breedveld FC, Eberl G, et al. Validity and reliability of the twenty-eight-joint count for the assessment of rheumatoid arthritis activity. *Arthritis Rheum* 1995;38(01):38–43
- 7 Sokka T. Work disability in early rheumatoid arthritis. *Clin Exp Rheumatol* 2003;21(05, Suppl 31):S71–S74
- 8 Klareskog L, Rönnelid J, Lundberg K, Padyukov L, Alfredsson L. Immunity to citrullinated proteins in rheumatoid arthritis. *Annu Rev Immunol* 2008;26:651–675
- 9 Ärlestig L, Mullazehi M, Kokkonen H, Rocklöv J, Rönnelid J, Dahlqvist SR. Antibodies against cyclic citrullinated peptides of IgG, IgA and IgM isotype and rheumatoid factor of IgM and IgA isotype are increased in unaffected members of multicase rheumatoid arthritis families from northern Sweden. *Ann Rheum Dis* 2012;71(06):825–829

- 10 Kolfenbach JR, Deane KD, Derber LA, et al. A prospective approach to investigating the natural history of preclinical rheumatoid arthritis (RA) using first-degree relatives of probands with RA. *Arthritis Rheum* 2009;61(12):1735–1742
- 11 Roudier J. Association of MHC and rheumatoid arthritis. Association of RA with HLA-DR4: the role of repertoire selection. *Arthritis Res* 2000;2(03):217–220
- 12 Picchianti-Diamanti A, Rosado MM, D'Amelio R. Infectious agents and inflammation: the role of microbiota in autoimmune arthritis. *Front Microbiol* 2018;8:2696
- 13 Salmi M, Andrew DP, Butcher EC, Jalkanen S. Dual binding capacity of mucosal immunoblasts to mucosal and synovial endothelium in humans: dissection of the molecular mechanisms. *J Exp Med* 1995;181(01):137–149
- 14 Rashid T, Ebringer A. Rheumatoid arthritis is linked to Proteus—the evidence. *Clin Rheumatol* 2007;26(07):1036–1043
- 15 Tishler M, Caspi D, Almog Y, Segal R, Yaron M. Increased incidence of urinary tract infection in patients with rheumatoid arthritis and secondary Sjögren's syndrome. *Ann Rheum Dis* 1992;51(05):604–606
- 16 Senior BW, Anderson GA, Morley KD, Kerr MA. Evidence that patients with rheumatoid arthritis have asymptomatic 'non-significant' *Proteus mirabilis* bacteriuria more frequently than healthy controls. *J Infect* 1999;38(02):99–106
- 17 Ebringer A, Cunningham P, Ahmadi K, Wrigglesworth J, Hosseini R, Wilson C. Sequence similarity between HLA-DR1 and DR4 subtypes associated with rheumatoid arthritis and proteus/serratia membrane haemolysins. *Ann Rheum Dis* 1992;51(11):1245–1246
- 18 Rashid T, Tiwana H, Wilson C, Ebringer A. Rheumatoid arthritis as an autoimmune disease caused by Proteus urinary tract infections: a proposal for a therapeutic protocol. *Isr Med Assoc J* 2001;3(09):675–680
- 19 Alexander C, Rietschel ET. Bacterial lipopolysaccharides and innate immunity. *J Endotoxin Res* 2001;7(03):167–202
- 20 Benoit R, Rowe S, Watkins SC, et al. Pure endotoxin does not pass across the intestinal epithelium in vitro. *Shock* 1998;10(01):43–48
- 21 Guo S, Al-Sadi R, Said HM, Ma TY. Lipopolysaccharide causes an increase in intestinal tight junction permeability in vitro and in vivo by inducing enterocyte membrane expression and localization of TLR-4 and CD14. *Am J Pathol* 2013;182(02):375–387
- 22 Maeda Y, Kumanogoh A, Takeda K. Altered composition of gut microbiota in rheumatoid arthritis patients. *Nihon Rinsho Meneki Gakkai Kaishi* 2016;39(01):59–63
- 23 Zhang X, Zhang D, Jia H, et al. The oral and gut microbiomes are perturbed in rheumatoid arthritis and partly normalized after treatment. *Nat Med* 2015;21(08):895–905
- 24 Lu YC, Yeh WC, Ohashi PS. LPS/TLR4 signal transduction pathway. *Cytokine* 2008;42(02):145–151
- 25 Koch L, Hofer S, Weigand MA, Frommhold D, Poeschl J. Lipopolysaccharide-induced activation of coagulation in neonatal cord and adult blood monitored by thrombelastography. *Thromb Res* 2009;124(04):463–467
- 26 Landsem A, Fure H, Christiansen D, et al. The key roles of complement and tissue factor in *Escherichia coli*-induced coagulation in human whole blood. *Clin Exp Immunol* 2015;182(01):81–89
- 27 Pretorius E, Mbotwe S, Bester J, Robinson CJ, Kell DB. Acute induction of anomalous and amyloidogenic blood clotting by molecular amplification of highly substoichiometric levels of bacterial lipopolysaccharide. *J R Soc Interface* 2016;13(122):20160539
- 28 Pretorius E, Akeredolu OO, Soma P, Kell DB. Major involvement of bacterial components in rheumatoid arthritis and its accompanying oxidative stress, systemic inflammation and hypercoagulability. *Exp Biol Med* (Maywood) 2017;242(04):355–373
- 29 Aviña-Zubieta JA, Choi HK, Sadatsafavi M, Etmann M, Esdaile JM, Lacaille D. Risk of cardiovascular mortality in patients with rheumatoid arthritis: a meta-analysis of observational studies. *Arthritis Rheum* 2008;59(12):1690–1697
- 30 Maradit-Kremers H, Nicola PJ, Crowson CS, Ballman KV, Gabriel SE. Cardiovascular death in rheumatoid arthritis: a population-based study. *Arthritis Rheum* 2005;52(03):722–732
- 31 McEntegart A, Capell HA, Czeran D, Rumley A, Woodward M, Lowe GD. Cardiovascular risk factors, including thrombotic variables, in a population with rheumatoid arthritis. *Rheumatology (Oxford)* 2001;40(06):640–644
- 32 del Rincón ID, Williams K, Stern MP, Freeman GL, Escalante A. High incidence of cardiovascular events in a rheumatoid arthritis cohort not explained by traditional cardiac risk factors. *Arthritis Rheum* 2001;44(12):2737–2745
- 33 Gallistl S, Mangge H, Neuwirth G, Muntean W. Activation of the haemostatic system in children with juvenile rheumatoid arthritis correlates with disease activity. *Thromb Res* 1998;92(06):267–272
- 34 Kwasny-Krochin B, Glusko P, Undas A. Unfavorably altered fibrin clot properties in patients with active rheumatoid arthritis. *Thromb Res* 2010;126(01):e11–e16
- 35 So AK, Varisco PA, Kemkes-Matthes B, et al. Arthritis is linked to local and systemic activation of coagulation and fibrinolysis pathways. *J Thromb Haemost* 2003;1(12):2510–2515
- 36 Habets KL, Trouw LA, Levarht EW, et al. Anti-citrullinated protein antibodies contribute to platelet activation in rheumatoid arthritis. *Arthritis Res Ther* 2015;17(01):209
- 37 Henn V, Slupsky JR, Gräfe M, et al. CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial cells. *Nature* 1998;391(6667):591–594
- 38 Amelot AA, Tagzirt M, Ducouret G, Kuen RL, Le Bonniec BF. Platelet factor 4 (CXCL4) seals blood clots by altering the structure of fibrin. *J Biol Chem* 2007;282(01):710–720
- 39 Wohner N. Role of cellular elements in thrombus formation and dissolution. *Cardiovasc Hematol Agents Med Chem* 2008;6(03):224–228
- 40 Chen J, Bierhaus A, Schiekofer S, et al. Tissue factor—a receptor involved in the control of cellular properties, including angiogenesis. *Thromb Haemost* 2001;86(01):334–345
- 41 Chen L, Lu Y, Chu Y, Xie J, Ding W, Wang F. Tissue factor expression in rheumatoid synovium: a potential role in pannus invasion of rheumatoid arthritis. *Acta Histochem* 2013;115(07):692–697
- 42 Mantovani A, Sozzani S, Vecchi A, Introna M, Allavena P. Cytokine activation of endothelial cells: new molecules for an old paradigm. *Thromb Haemost* 1997;78(01):406–414
- 43 Pretorius E, Oberholzer HM, van der Spuy WJ, Swanepoel AC, Soma P. Scanning electron microscopy of fibrin networks in rheumatoid arthritis: a qualitative analysis. *Rheumatol Int* 2012;32(06):1611–1615
- 44 Sánchez-Pernaute O, Largo R, Calvo E, Alvarez-Soria MA, Egido J, Herrero-Baumont G. A fibrin based model for rheumatoid synovitis. *Ann Rheum Dis* 2003;62(12):1135–1138
- 45 Masson-Bessière C, Sebbag M, Durieux JJ, et al. In the rheumatoid pannus, anti-filaggrin autoantibodies are produced by local plasma cells and constitute a higher proportion of IgG than in synovial fluid and serum. *Clin Exp Immunol* 2000;119(03):544–552
- 46 Sebbag M, Moinard N, Auger I, et al. Epitopes of human fibrin recognized by the rheumatoid arthritis-specific autoantibodies to citrullinated proteins. *Eur J Immunol* 2006;36(08):2250–2263
- 47 Falkenburg WJ, van Schaardenburg D, Ooijevaar-de Heer P, Wolbink G, Rispens T. IgG subclass specificity discriminates restricted IgM rheumatoid factor responses from more mature anti-citrullinated protein antibody-associated or isotype-switched IgA responses. *Arthritis Rheumatol* 2015;67(12):3124–3134
- 48 Egeland T, Munthe E. The role of the laboratory in rheumatology. Rheumatoid factors. *Clin Rheum Dis* 1983;9(01):135–160
- 49 Trier NH, Holm BE, Heiden J, et al. The use of synthetic peptides for detection of anti-citrullinated protein antibodies in rheumatoid arthritis. *J Immunol Methods* 2018;454:6–14

- 50 Nishimura K, Sugiyama D, Kogata Y, et al. Meta-analysis: diagnostic accuracy of anti-cyclic citrullinated peptide antibody and rheumatoid factor for rheumatoid arthritis. *Ann Intern Med* 2007;146(11):797–808
- 51 Lingampalli N, Sokolove J, Lahey LJ, et al. Combination of anti-citrullinated protein antibodies and rheumatoid factor is associated with increased systemic inflammatory mediators and more rapid progression from preclinical to clinical rheumatoid arthritis. *Clin Immunol* 2018;195:119–126
- 52 Bugatti S, Bogliolo L, Vitolo B, Manzo A, Montecucco C, Caporali R. Anti-citrullinated protein antibodies and high levels of rheumatoid factor are associated with systemic bone loss in patients with early untreated rheumatoid arthritis. *Arthritis Res Ther* 2016;18(01):226
- 53 Trela M, Perera S, Sheeran T, Rylance P, Nelson PN, Attridge K. Citrullination facilitates cross-reactivity of rheumatoid factor with non-IgG1 Fc epitopes in rheumatoid arthritis. *Sci Rep* 2019; 9(01):12068
- 54 Nevius E, Gomes AC, Pereira JP. Inflammatory cell migration in rheumatoid arthritis: a comprehensive review. *Clin Rev Allergy Immunol* 2016;51(01):59–78
- 55 Rajmakers R, van Beers JJ, El-Azzouny M, et al. Elevated levels of fibrinogen-derived endogenous citrullinated peptides in synovial fluid of rheumatoid arthritis patients. *Arthritis Res Ther* 2012;14(03):R114
- 56 Tarcsa E, Marekov LN, Mei G, Melino G, Lee SC, Steinert PM. Protein unfolding by peptidylarginine deiminase. Substrate specificity and structural relationships of the natural substrates trichohyalin and filaggrin. *J Biol Chem* 1996;271(48):30709–30716
- 57 Masson-Bessière C, Sebbag M, Girbal-Neuhausser E, et al. The major synovial targets of the rheumatoid arthritis-specific anti-filaggrin autoantibodies are deiminated forms of the alpha- and beta-chains of fibrin. *J Immunol* 2001;166(06):4177–4184
- 58 Hagiwara T, Hidaka Y, Yamada M. Deimination of histone H2A and H4 at arginine 3 in HL-60 granulocytes. *Biochemistry* 2005; 44(15):5827–5834
- 59 Vossenaar ER, Després N, Lapointe E, et al. Rheumatoid arthritis specific anti-Sa antibodies target citrullinated vimentin. *Arthritis Res Ther* 2004;6(02):R142–R150
- 60 Kinloch A, Tatzer V, Wait R, et al. Identification of citrullinated alpha-enolase as a candidate autoantigen in rheumatoid arthritis. *Arthritis Res Ther* 2005;7(06):R1421–R1429
- 61 Burkhardt H, Sehnert B, Bockermann R, Engström A, Kalden JR, Holmdahl R. Humoral immune response to citrullinated collagen type II determinants in early rheumatoid arthritis. *Eur J Immunol* 2005;35(05):1643–1652
- 62 Suzuki A, Yamada R, Ohtake-Yamanaka M, Okazaki Y, Sawada T, Yamamoto K. Anti-citrullinated collagen type I antibody is a target of autoimmunity in rheumatoid arthritis. *Biochem Biophys Res Commun* 2005;333(02):418–426
- 63 Hill JA, Southwood S, Sette A, Jevnikar AM, Bell DA, Cairns E. Cutting edge: the conversion of arginine to citrulline allows for a high-affinity peptide interaction with the rheumatoid arthritis-associated HLA-DRB1*0401 MHC class II molecule. *J Immunol* 2003;171(02):538–541
- 64 Suwannaalai P, Trouw LA, Toes RE, Huizinga TW. Anti-citrullinated protein antibodies (ACPA) in early rheumatoid arthritis. *Mod Rheumatol* 2012;22(01):15–20
- 65 Rantapää-Dahlqvist S, de Jong BA, Berglin E, et al. Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the development of rheumatoid arthritis. *Arthritis Rheum* 2003;48(10):2741–2749
- 66 Nielsen MM, van Schaardenburg D, Reesink HW, et al. Specific autoantibodies precede the symptoms of rheumatoid arthritis: a study of serial measurements in blood donors. *Arthritis Rheum* 2004;50(02):380–386
- 67 Klareskog L, Catrina AI, Paget S. Rheumatoid arthritis. *Lancet* 2009;373(9664):659–672
- 68 Vossenaar ER, Zendman AJ, van Venrooij WJ, Puijn GJ. PAD, a growing family of citrullinating enzymes: genes, features and involvement in disease. *BioEssays* 2003;25(11):1106–1118
- 69 Asaga H, Yamada M, Senshu T. Selective deimination of vimentin in calcium ionophore-induced apoptosis of mouse peritoneal macrophages. *Biochem Biophys Res Commun* 1998;243(03):641–646
- 70 Schwab BL, Guerini D, Didszun C, et al. Cleavage of plasma membrane calcium pumps by caspases: a link between apoptosis and necrosis. *Cell Death Differ* 2002;9(08):818–831
- 71 Szekecz Z, Soós L, Szabó Z, et al. Anti-citrullinated protein antibodies in rheumatoid arthritis: as good as it gets? *Clin Rev Allergy Immunol* 2008;34(01):26–31
- 72 Puszczewicz M, Iwaszkiewicz C. Role of anti-citrullinated protein antibodies in diagnosis and prognosis of rheumatoid arthritis. *Arch Med Sci* 2011;7(02):189–194
- 73 Farid SSH, Azizi G, Mirshafiey A. Anti-citrullinated protein antibodies and their clinical utility in rheumatoid arthritis. *Int J Rheum Dis* 2013;16(04):379–386
- 74 Chang X, Xia Y, Pan J, Meng Q, Zhao Y, Yan X. PAD2 is significantly associated with rheumatoid arthritis. *PLoS One* 2013;8(12): e81259
- 75 Damgaard D, Bawadekar M, Senolt L, Stensballe A, Shelef MA, Nielsen CH. Relative efficiencies of peptidylarginine deiminase 2 and 4 in generating target sites for anti-citrullinated protein antibodies in fibrinogen, alpha-enolase and histone H3. *PLoS One* 2018;13(08):e0203214
- 76 Holers VM. Autoimmunity to citrullinated proteins and the initiation of rheumatoid arthritis. *Curr Opin Immunol* 2013;25 (06):728–735
- 77 Bang SY, Lee KH, Cho SK, Lee HS, Lee KW, Bae SC. Smoking increases rheumatoid arthritis susceptibility in individuals carrying the HLA-DRB1 shared epitope, regardless of rheumatoid factor or anti-cyclic citrullinated peptide antibody status. *Arthritis Rheum* 2010;62(02):369–377
- 78 Mikuls TR, Payne JB, Yu F, et al. Periodontitis and *Porphyromonas gingivalis* in patients with rheumatoid arthritis. *Arthritis Rheumatol* 2014;66(05):1090–1100
- 79 Mikuls TR, Payne JB, Reinhardt RA, et al. Antibody responses to *Porphyromonas gingivalis* (*P. gingivalis*) in subjects with rheumatoid arthritis and periodontitis. *Int Immunopharmacol* 2009; 9(01):38–42
- 80 Shimada A, Kobayashi T, Ito S, et al. Expression of anti-*Porphyromonas gingivalis* peptidylarginine deiminase immunoglobulin G and peptidylarginine deiminase-4 in patients with rheumatoid arthritis and periodontitis. *J Periodontol Res* 2016;51(01):103–111
- 81 Okada M, Kobayashi T, Ito S, et al. Periodontal treatment decreases levels of antibodies to *Porphyromonas gingivalis* and citrulline in patients with rheumatoid arthritis and periodontitis. *J Periodontol* 2013;84(12):e74–e84
- 82 Reparón-Schuijt CC, van Esch WJ, van Kooten C, et al. Secretion of anti-citrulline-containing peptide antibody by B lymphocytes in rheumatoid arthritis. *Arthritis Rheum* 2001;44(01):41–47
- 83 Ioan-Facsinay A, el-Bannoudi H, Scherer HU, et al. Anti-cyclic citrullinated peptide antibodies are a collection of anti-citrullinated protein antibodies and contain overlapping and non-overlapping reactivities. *Ann Rheum Dis* 2011;70(01):188–193
- 84 Snir O, Widhe M, von Spee C, et al. Multiple antibody reactivities to citrullinated antigens in sera from patients with rheumatoid arthritis: association with HLA-DRB1 alleles. *Ann Rheum Dis* 2009;68(05):736–743
- 85 Li S, Yu Y, Yue Y, et al. Autoantibodies from single circulating plasmablasts react with citrullinated antigens and *Porphyromonas gingivalis* in rheumatoid arthritis. *Arthritis Rheumatol* 2016; 68(03):614–626
- 86 Schellekens GA, de Jong BA, van den Hoogen FH, van de Putte LB, van Venrooij WJ. Citrulline is an essential constituent of antigenic determinants recognized by rheumatoid arthritis-specific autoantibodies. *J Clin Invest* 1998;101(01):273–281

- 87 Chibnik LB, Mandl LA, Costenbader KH, Schur PH, Karlson EW. Comparison of threshold cutpoints and continuous measures of anti-cyclic citrullinated peptide antibodies in predicting future rheumatoid arthritis. *J Rheumatol* 2009;36(04):706–711
- 88 van de Stadt LA, van der Horst AR, de Koning MH, et al. The extent of the anti-citrullinated protein antibody repertoire is associated with arthritis development in patients with seropositive arthralgia. *Ann Rheum Dis* 2011;70(01):128–133
- 89 van der Woude D, Rantapää-Dahlqvist S, Ioan-Facsinay A, et al. Epitope spreading of the anti-citrullinated protein antibody response occurs before disease onset and is associated with the disease course of early arthritis. *Ann Rheum Dis* 2010;69(08):1554–1561
- 90 Derksen VF, Ajeganova S, Trouw LA, et al. Rheumatoid arthritis phenotype at presentation differs depending on the number of autoantibodies present. *Ann Rheum Dis* 2017;76(04):716–720
- 91 van der Woude D, Syversen SW, van der Voort EI, et al. The ACPA isotype profile reflects long-term radiographic progression in rheumatoid arthritis. *Ann Rheum Dis* 2010;69(06):1110–1116
- 92 Aletaha D, Neogi T, Silman AJ, et al. 2010 rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum* 2010;62(09):2569–2581
- 93 Nienhuis RL, Mandema E. A new serum factor in patients with rheumatoid arthritis; the antiperinuclear factor. *Ann Rheum Dis* 1964;23:302–305
- 94 Young BJ, Mallya RK, Leslie RD, Clark CJ, Hamblin TJ. Anti-keratin antibodies in rheumatoid arthritis. *BMJ* 1979;2(6182):97–99
- 95 Schellekens GA, Visser H, de Jong BA, et al. The diagnostic properties of rheumatoid arthritis antibodies recognizing a cyclic citrullinated peptide. *Arthritis Rheum* 2000;43(01):155–163
- 96 van Venrooij WJ, Hazes JM, Visser H. Anticitrullinated protein/peptide antibody and its role in the diagnosis and prognosis of early rheumatoid arthritis. *Neth J Med* 2002;60(10):383–388
- 97 Okumura N, Haneishi A, Terasawa F. Citrullinated fibrinogen shows defects in FPA and FPB release and fibrin polymerization catalyzed by thrombin. *Clin Chim Acta* 2009;401(1–2):119–123
- 98 Nakayama-Hamada M, Suzuki A, Furukawa H, Yamada R, Yamamoto K. Citrullinated fibrinogen inhibits thrombin-catalysed fibrin polymerization. *J Biochem* 2008;144(03):393–398
- 99 Yermolenko IS, Lishko VK, Ugarova TP, Magonov SN. High-resolution visualization of fibrinogen molecules and fibrin fibers with atomic force microscopy. *Biomacromolecules* 2011;12(02):370–379
- 100 Joshua V, Schobers L, Titcombe PJ, et al. Antibody responses to de novo identified citrullinated fibrinogen peptides in rheumatoid arthritis and visualization of the corresponding B cells. *Arthritis Res Ther* 2016;18(01):284
- 101 Zhao Y, Tian X, Li Z. Prevalence and clinical significance of antibodies to citrullinated fibrinogen (ACF) in Chinese patients with rheumatoid arthritis. *Clin Rheumatol* 2007;26(09):1505–1512
- 102 Cantaert T, Teitsma C, Tak PP, Baeten D. Presence and role of anti-citrullinated protein antibodies in experimental arthritis models. *Arthritis Rheum* 2013;65(04):939–948
- 103 Shin K, Hong S, Choi E-H, et al. Role of citrullinated fibrinogen peptides in the activation of CD4 T cells from patients with rheumatoid arthritis. *Immune Netw* 2013;13(04):116–122
- 104 Šenolt L, Grassi W, Szodoray P. Laboratory biomarkers or imaging in the diagnostics of rheumatoid arthritis? *BMC Med* 2014;12:49
- 105 Giancchetti E, Crinò A, Giorda E, et al. Altered B cell homeostasis and toll-like receptor 9-driven response in type 1 diabetes carriers of the C1858T PTPN22 allelic variant: implications in the disease pathogenesis. *PLoS One* 2014;9(10):e110755
- 106 Giltaiy NV, Chappell CP, Clark EA. B-cell selection and the development of autoantibodies. *Arthritis Res Ther* 2012;14(Suppl 4):S1
- 107 Menard L, Saadoun D, Isnardi I, et al. The PTPN22 allele encoding an R620W variant interferes with the removal of developing autoreactive B cells in humans. *J Clin Invest* 2011;121(09):3635–3644
- 108 Ho PP, Lee LY, Zhao X, et al. Autoimmunity against fibrinogen mediates inflammatory arthritis in mice. *J Immunol* 2010;184(01):379–390
- 109 Ingegnoli F, Fantini F, Favalli EG, et al. Inflammatory and pro-thrombotic biomarkers in patients with rheumatoid arthritis: effects of tumor necrosis factor- α blockade. *J Autoimmun* 2008;31(02):175–179
- 110 Rooney T, Scherzer R, Shigenaga JK, Graf J, Imboden JB, Grunfeld C. Levels of plasma fibrinogen are elevated in well-controlled rheumatoid arthritis. *Rheumatology (Oxford)* 2011;50(08):1458–1465
- 111 Zhao X, Okeke NL, Sharpe O, et al. Circulating immune complexes contain citrullinated fibrinogen in rheumatoid arthritis. *Arthritis Res Ther* 2008;10(04):R94
- 112 Nielsen MM, van der Horst AR, van Schaardenburg D, et al. Antibodies to citrullinated human fibrinogen (ACF) have diagnostic and prognostic value in early arthritis. *Ann Rheum Dis* 2005;64(08):1199–1204
- 113 Vander Cruyssen B, Cantaert T, Nogueira L, et al. Diagnostic value of anti-human citrullinated fibrinogen ELISA and comparison with four other anti-citrullinated protein assays. *Arthritis Res Ther* 2006;8(04):R122
- 114 Hill JA, Bell DA, Brintnell W, et al. Arthritis induced by post-translationally modified (citrullinated) fibrinogen in DR4-IE transgenic mice. *J Exp Med* 2008;205(04):967–979
- 115 van Beers JJBC, Rajmakers R, Alexander L-E, et al. Mapping of citrullinated fibrinogen B-cell epitopes in rheumatoid arthritis by imaging surface plasmon resonance. *Arthritis Res Ther* 2010;12(06):R219–R219
- 116 Nakayama-Hamada M, Suzuki A, Kubota K, et al. Comparison of enzymatic properties between hPAD12 and hPAD14. *Biochem Biophys Res Commun* 2005;327(01):192–200
- 117 Sanchez-Pernaute O, Filkova M, Gabucio A, et al. Citrullination enhances the pro-inflammatory response to fibrin in rheumatoid arthritis synovial fibroblasts. *Ann Rheum Dis* 2013;72(08):1400–1406
- 118 Wolberg AS. Thrombin generation and fibrin clot structure. *Blood Rev* 2007;21(03):131–142
- 119 Ohba T, Takase Y, Ohhara M, Kasukawa R. Thrombin in the synovial fluid of patients with rheumatoid arthritis mediates proliferation of synovial fibroblast-like cells by induction of platelet derived growth factor. *J Rheumatol* 1996;23(09):1505–1511
- 120 Sebbag M, Chapuy-Regaud S, Auger I, et al. Clinical and pathophysiological significance of the autoimmune response to citrullinated proteins in rheumatoid arthritis. *Joint Bone Spine* 2004;71(06):493–502
- 121 Yamada R, Suzuki A, Chang X, Yamamoto K. Citrullinated proteins in rheumatoid arthritis. *Front Biosci* 2005;10:54–64
- 122 Henschen A, Lottspeich F, Kehl M, Southan C. Covalent structure of fibrinogen. *Ann N Y Acad Sci* 1983;408:28–43
- 123 Cohen C, Parry DA. Alpha-helical coiled coils and bundles: how to design an alpha-helical protein. *Proteins* 1990;7(01):1–15
- 124 Kollman JM, Pandi L, Sawaya MR, Riley M, Doolittle RF. Crystal structure of human fibrinogen. *Biochemistry* 2009;48(18):3877–3886
- 125 Litvinov RI, Faizullin DA, Zuev YF, Weisel JW. The α -helix to β -sheet transition in stretched and compressed hydrated fibrin clots. *Biophys J* 2012;103(05):1020–1027
- 126 Zhmurov A, Kononova O, Litvinov RI, Dima RI, Barsegov V, Weisel JW. Mechanical transition from α -helical coiled coils to β -sheets in fibrin(ogen). *J Am Chem Soc* 2012;134(50):20396–20402
- 127 Purohit PK, Litvinov RI, Brown AE, Discher DE, Weisel JW. Protein unfolding accounts for the unusual mechanical behavior of fibrin networks. *Acta Biomater* 2011;7(06):2374–2383

- 128 Greenwald J, Riek R. Biology of amyloid: structure, function, and regulation. *Structure* 2010;18(10):1244–1260
- 129 Eisenberg D, Jucker M. The amyloid state of proteins in human diseases. *Cell* 2012;148(06):1188–1203
- 130 Pepys MB. Amyloidosis. *Annu Rev Med* 2006;57:223–241
- 131 Stangou AJ, Banner NR, Hendry BM, et al. Hereditary fibrinogen A alpha-chain amyloidosis: phenotypic characterization of a systemic disease and the role of liver transplantation. *Blood* 2010;115(15):2998–3007
- 132 Pretorius E, Bester J, Page MJ, Kell DB. The potential of LPS-binding protein to reverse amyloid formation in plasma fibrin of individuals with Alzheimer-type dementia. *Front Aging Neurosci* 2018;10:257
- 133 Pretorius E, Page MJ, Engelbrecht L, Ellis GC, Kell DB. Substantial fibrin amyloidogenesis in type 2 diabetes assessed using amyloid-selective fluorescent stains. *Cardiovasc Diabetol* 2017;16(01):141
- 134 Pretorius E, Page MJ, Mbotwe S, Kell DB. Lipopolysaccharide-binding protein (LBP) can reverse the amyloid state of fibrin seen or induced in Parkinson's disease. *PLoS One* 2018;13(03):e0192121
- 135 Pretorius E, Page MJ, Hendricks L, Nkosi NB, Benson SR, Kell DB. Both lipopolysaccharide and lipoteichoic acids potentially induce anomalous fibrin amyloid formation: assessment with novel Amytracker™ stains. *J R Soc Interface* 2018;15(139):20170941
- 136 Page MJ, Thomson GJA, Nunes JM, et al. Serum amyloid A binds to fibrin(ogen), promoting fibrin amyloid formation. *Sci Rep* 2019;9(01):3102
- 137 Osaki D, Hiramatsu H. Citrullination and deamidation affect aggregation properties of amyloid β -proteins. *Amyloid* 2016;23(04):234–241
- 138 Sakono M, Zako T. Amyloid oligomers: formation and toxicity of A β oligomers. *FEBS J* 2010;277(06):1348–1358
- 139 Connolly M, Marrelli A, Blades M, et al. Acute serum amyloid A induces migration, angiogenesis, and inflammation in synovial cells in vitro and in a human rheumatoid arthritis/SCID mouse chimera model. *J Immunol* 2010;184(11):6427–6437
- 140 Cunnane G, Grehan S, Geoghegan S, et al. Serum amyloid A in the assessment of early inflammatory arthritis. *J Rheumatol* 2000;27(01):58–63
- 141 Hwang YG, Balasubramani GK, Metes ID, Levesque MC, Bridges SL Jr, Moreland LW. Differential response of serum amyloid A to different therapies in early rheumatoid arthritis and its potential value as a disease activity biomarker. *Arthritis Res Ther* 2016;18(01):108
- 142 Targońska-Stępnik B, Majdan M. Serum amyloid A as a marker of persistent inflammation and an indicator of cardiovascular and renal involvement in patients with rheumatoid arthritis. *Mediators Inflamm* 2014;2014:793628
- 143 Cunnane G, Whitehead AS. Amyloid precursors and amyloidosis in rheumatoid arthritis. *Best Pract Res Clin Rheumatol* 1999;13(04):615–628
- 144 Husby G, Husebekk A, Skogen B, et al. Serum amyloid A (SAA)—the precursor of protein AA in secondary amyloidosis. *Adv Exp Med Biol* 1988;243:185–192
- 145 Steel DM, Whitehead AS. The major acute phase reactants: C-reactive protein, serum amyloid P component and serum amyloid A protein. *Immunol Today* 1994;15(02):81–88
- 146 O'Hara R, Murphy EP, Whitehead AS, FitzGerald O, Bresnihan B. Acute-phase serum amyloid A production by rheumatoid arthritis synovial tissue. *Arthritis Res* 2000;2(02):142–144
- 147 Kumon Y, Suehiro T, Hashimoto K, Nakatani K, Sipe JD. Local expression of acute phase serum amyloid A mRNA in rheumatoid arthritis synovial tissue and cells. *J Rheumatol* 1999;26(04):785–790
- 148 Urieli-Shoval S, Meek RL, Hanson RH, Eriksen N, Benditt EP. Human serum amyloid A genes are expressed in monocyte/macrophage cell lines. *Am J Pathol* 1994;145(03):650–660
- 149 Mullan RH, Bresnihan B, Golden-Mason L, et al. Acute-phase serum amyloid A stimulation of angiogenesis, leukocyte recruitment, and matrix degradation in rheumatoid arthritis through an NF-kappaB-dependent signal transduction pathway. *Arthritis Rheum* 2006;54(01):105–114
- 150 Kell DB, Pretorius E. Proteins behaving badly. Substoichiometric molecular control and amplification of the initiation and nature of amyloid fibril formation: lessons from and for blood clotting. *Prog Biophys Mol Biol* 2017;123:16–41
- 151 Naito M. *Porphyromonas gingivalis*-induced platelet aggregation in human plasma. *Journal of Oral Biosciences*. 2007;49(03):173–179
- 152 Stobernack T, du Teil Espina M, Mulder LM, et al. A secreted bacterial peptidylarginine deiminase can neutralize human innate immune defenses. *MBio* 2018;9(05):e01704–e01718
- 153 Abdullah S-N, Farmer E-A, Spargo L, Logan R, Gully N. *Porphyromonas gingivalis* peptidylarginine deiminase substrate specificity. *Anaerobe* 2013;23:102–108
- 154 Wegner N, Wait R, Sroka A, et al. Peptidylarginine deiminase from *Porphyromonas gingivalis* citrullinates human fibrinogen and α -enolase: implications for autoimmunity in rheumatoid arthritis. *Arthritis Rheum* 2010;62(09):2662–2672
- 155 Potempa J, Pike R, Travis J. The multiple forms of trypsin-like activity present in various strains of *Porphyromonas gingivalis* are due to the presence of either Arg-gingipain or Lys-gingipain. *Infect Immun* 1995;63(04):1176–1182
- 156 McGraw WT, Potempa J, Farley D, Travis J. Purification, characterization, and sequence analysis of a potential virulence factor from *Porphyromonas gingivalis*, peptidylarginine deiminase. *Infect Immun* 1999;67(07):3248–3256
- 157 Quirke A-M, Lugli EB, Wegner N, et al. Heightened immune response to autocitrullinated *Porphyromonas gingivalis* peptidylarginine deiminase: a potential mechanism for breaching immunologic tolerance in rheumatoid arthritis. *Ann Rheum Dis* 2014;73(01):263–269
- 158 Montgomery AB, Kopec J, Shrestha L, et al. Crystal structure of *Porphyromonas gingivalis* peptidylarginine deiminase: implications for autoimmunity in rheumatoid arthritis. *Ann Rheum Dis* 2016;75(06):1255–1261
- 159 Yamakawa M, Ouhara K, Kajiyama M, et al. *Porphyromonas gingivalis* infection exacerbates the onset of rheumatoid arthritis in SKG mice. *Clin Exp Immunol* 2016;186(02):177–189
- 160 Khandpur R, Carmona-Rivera C, Vivekanandan-Giri A, et al. NETs are a source of citrullinated autoantigens and stimulate inflammatory responses in rheumatoid arthritis. *Sci Transl Med* 2013;5(178):178ra40
- 161 Bartold PM, Marino V, Cantley M, Haynes DR. Effect of *Porphyromonas gingivalis*-induced inflammation on the development of rheumatoid arthritis. *J Clin Periodontol* 2010;37(05):405–411
- 162 Lönn J, Ljunggren S, Klarström-Engström K, Demirel I, Bengtsson T, Karlsson H. Lipoprotein modifications by gingipains of *Porphyromonas gingivalis*. *J Periodontal Res* 2018;53(03):403–413
- 163 Bale BF, Doneen AL, Vigerust DJ. High-risk periodontal pathogens contribute to the pathogenesis of atherosclerosis. *Postgrad Med J* 2017;93(1098):215–220
- 164 Imamura T, Tanase S, Hamamoto T, Potempa J, Travis J. Activation of blood coagulation factor IX by gingipains R, arginine-specific cysteine proteinases from *Porphyromonas gingivalis*. *Biochem J* 2001;353(Pt 2):325–331
- 165 Imamura T, Potempa J, Tanase S, Travis J. Activation of blood coagulation factor X by arginine-specific cysteine proteinases (gingipain-Rs) from *Porphyromonas gingivalis*. *J Biol Chem* 1997;272(25):16062–16067
- 166 Imamura T, Banbula A, Pereira PJ, Travis J, Potempa J. Activation of human prothrombin by arginine-specific cysteine proteinases (Gingipains R) from *Porphyromonas gingivalis*. *J Biol Chem* 2001;276(22):18984–18991
- 167 Guo Y, Nguyen KA, Potempa J. Dichotomy of gingipains action as virulence factors: from cleaving substrates with the precision of a surgeon's knife to a meat chopper-like brutal degradation of proteins. *Periodontol* 2000 2010;54(01):15–44

- 168 Imamura T, Potempa J, Pike RN, Moore JN, Barton MH, Travis J. Effect of free and vesicle-bound cysteine proteinases of *Porphyromonas gingivalis* on plasma clot formation: implications for bleeding tendency at periodontitis sites. *Infect Immun* 1995;63(12):4877–4882
- 169 Lourbakos A, Yuan YP, Jenkins AL, et al. Activation of protease-activated receptors by gingipains from *Porphyromonas gingivalis* leads to platelet aggregation: a new trait in microbial pathogenicity. *Blood* 2001;97(12):3790–3797
- 170 Adams B, Nunes JM, Page MJ, et al. Parkinson's disease: a systemic inflammatory disease accompanied by bacterial inflammagens. *Front Aging Neurosci* 2019;11(210):210
- 171 McInnes IB, Schett G. The pathogenesis of rheumatoid arthritis. *N Engl J Med* 2011;365(23):2205–2219
- 172 Vossenaar ER, Smeets TJ, Kraan MC, Raats JM, van Venrooij WJ, Tak PP. The presence of citrullinated proteins is not specific for rheumatoid synovial tissue. *Arthritis Rheum* 2004;50(11):3485–3494
- 173 van der Helm-van Mil AH, Verpoort KN, Breedveld FC, Huizinga TW, Toes RE, de Vries RR. The HLA-DRB1 shared epitope alleles are primarily a risk factor for anti-cyclic citrullinated peptide antibodies and are not an independent risk factor for development of rheumatoid arthritis. *Arthritis Rheum* 2006;54(04):1117–1121
- 174 Vossenaar ER, Radstake TR, van der Heijden A, et al. Expression and activity of citrullinating peptidylarginine deiminase enzymes in monocytes and macrophages. *Ann Rheum Dis* 2004;63(04):373–381
- 175 Chang X, Yamada R, Suzuki A, et al. Localization of peptidylarginine deiminase 4 (PADI4) and citrullinated protein in synovial tissue of rheumatoid arthritis. *Rheumatology (Oxford)* 2005;44(01):40–50
- 176 Smiley ST, King JA, Hancock WW. Fibrinogen stimulates macrophage chemokine secretion through toll-like receptor 4. *J Immunol* 2001;167(05):2887–2894
- 177 Hsieh JY, Smith TD, Meli VS, Tran TN, Botvinick EL, Liu WF. Differential regulation of macrophage inflammatory activation by fibrin and fibrinogen. *Acta Biomater* 2017;47:14–24
- 178 Rubel C, Fernández GC, Dran G, Bompadre MB, Isturiz MA, Palermo MS. Fibrinogen promotes neutrophil activation and delays apoptosis. *J Immunol* 2001;166(03):2002–2010
- 179 Guo M, Sahni SK, Sahni A, Francis CW. Fibrinogen regulates the expression of inflammatory chemokines through NF-kappaB activation of endothelial cells. *Thromb Haemostasis* 2004;92(04):858–866
- 180 Liu X, Piela-Smith TH. Fibrin(ogen)-induced expression of ICAM-1 and chemokines in human synovial fibroblasts. *J Immunol* 2000;165(09):5255–5261
- 181 Weinberg JB, Phippen AM, Greenberg CS. Extravascular fibrin formation and dissolution in synovial tissue of patients with osteoarthritis and rheumatoid arthritis. *Arthritis Rheum* 1991;34(08):996–1005
- 182 Clavel C, Ceccato L, Anquetil F, Serre G, Sebbag M. Among human macrophages polarised to different phenotypes, the M-CSF-oriented cells present the highest pro-inflammatory response to the rheumatoid arthritis-specific immune complexes containing ACPA. *Ann Rheum Dis* 2016;75(12):2184–2191
- 183 Sokolove J, Zhao X, Chandra PE, Robinson WH. Immune complexes containing citrullinated fibrinogen costimulate macrophages via Toll-like receptor 4 and Fcγ receptor. *Arthritis Rheum* 2011;63(01):53–62
- 184 Zhu W, Li X, Fang S, et al. Anti-citrullinated protein antibodies induce macrophage subset disequilibrium in RA patients. *Inflammation* 2015;38(06):2067–2075
- 185 Bianchi ME. DAMPs, PAMPs and alarmins: all we need to know about danger. *J Leukoc Biol* 2007;81(01):1–5
- 186 Hu F, Li Y, Zheng L, et al. Toll-like receptors expressed by synovial fibroblasts perpetuate Th1 and Th17 cell responses in rheumatoid arthritis. *PLoS One* 2014;9(06):e100266
- 187 Bartok B, Firestein GS. Fibroblast-like synoviocytes: key effector cells in rheumatoid arthritis. *Immunol Rev* 2010;233(01):233–255
- 188 Makarov SS. NF-kappa B in rheumatoid arthritis: a pivotal regulator of inflammation, hyperplasia, and tissue destruction. *Arthritis Res* 2001;3(04):200–206
- 189 Smolen JS, Redlich K, Zwerina J, Aletaha D, Steiner G, Schett G. Pro-inflammatory cytokines in rheumatoid arthritis: pathogenetic and therapeutic aspects. *Clin Rev Allergy Immunol* 2005;28(03):239–248
- 190 Siebert S, Tsoukas A, Robertson J, McInnes I. Cytokines as therapeutic targets in rheumatoid arthritis and other inflammatory diseases. *Pharmacol Rev* 2015;67(02):280–309
- 191 Bystrom J, Clanchy FI, Taher TE, et al. TNFα in the regulation of Treg and Th17 cells in rheumatoid arthritis and other autoimmune inflammatory diseases. *Cytokine* 2018;101:4–13
- 192 Rho YH, Chung CP, Oeser A, et al. Inflammatory mediators and premature coronary atherosclerosis in rheumatoid arthritis. *Arthritis Rheum* 2009;61(11):1580–1585
- 193 Kawashima M, Miossec P. Effect of treatment of rheumatoid arthritis with infliximab on IFN gamma, IL4, T-bet, and GATA-3 expression: link with improvement of systemic inflammation and disease activity. *Ann Rheum Dis* 2005;64(03):415–418
- 194 Pawlik A, Ostanek L, Brzosko I, et al. Therapy with infliximab decreases the CD4+CD28-T cell compartment in peripheral blood in patients with rheumatoid arthritis. *Rheumatol Int* 2004;24(06):351–354
- 195 Abbas S, Zhang YH, Clohisy JC, Abu-Amer Y. Tumor necrosis factor-alpha inhibits pre-osteoblast differentiation through its type-1 receptor. *Cytokine* 2003;22(1-2):33–41
- 196 Lam J, Takeshita S, Barker JE, Kanagawa O, Ross FP, Teitelbaum SL. TNF-alpha induces osteoclastogenesis by direct stimulation of macrophages exposed to permissive levels of RANK ligand. *J Clin Invest* 2000;106(12):1481–1488
- 197 Pignatelli P, De Biase L, Lenti L, et al. Tumor necrosis factor-alpha as trigger of platelet activation in patients with heart failure. *Blood* 2005;106(06):1992–1994
- 198 Chen W, Wang Q, Ke Y, Lin J. Neutrophil function in an inflammatory milieu of rheumatoid arthritis. *J Immunol Res* 2018;2018:8549329
- 199 Khawaja AA, Pericleous C, Ripoll VM, Porter JC, Giles IP. Autoimmune rheumatic disease IgG has differential effects upon neutrophil integrin activation that is modulated by the endothelium. *Sci Rep* 2019;9(01):1283
- 200 Thieblemont N, Wright HL, Edwards SW, Witko-Sarsat V. Human neutrophils in auto-immunity. Paper presented at: Seminars in Immunology; 2016
- 201 Brinkmann V, Reichard U, Goosmann C, et al. Neutrophil extracellular traps kill bacteria. *Science* 2004;303(5663):1532–1535
- 202 Zou Y, Chen X, Xiao J, et al. Neutrophil extracellular traps promote lipopolysaccharide-induced airway inflammation and mucus hypersecretion in mice. *Oncotarget* 2018;9(17):13276–13286
- 203 Clark SR, Ma AC, Tavener SA, et al. Platelet TLR4 activates neutrophil extracellular traps to ensnare bacteria in septic blood. *Nat Med* 2007;13(04):463–469
- 204 Keshari RS, Jyoti A, Dubey M, et al. Cytokines induced neutrophil extracellular traps formation: implication for the inflammatory disease condition. *PLoS One* 2012;7(10):e48111
- 205 Spengler J, Lugonja B, Ytterberg AJ, et al. Release of active peptidyl arginine deiminases by neutrophils can explain production of extracellular citrullinated autoantigens in rheumatoid arthritis synovial fluid. *Arthritis Rheumatol* 2015;67(12):3135–3145
- 206 Varjú I, Kolev K. Networks that stop the flow: a fresh look at fibrin and neutrophil extracellular traps. *Thromb Res* 2019;182:1–11
- 207 Fuchs TA, Brill A, Duerschmied D, et al. Extracellular DNA traps promote thrombosis. *Proc Natl Acad Sci U S A* 2010;107(36):15880–15885

- 208 Varjú I, Longstaff C, Szabó L, et al. DNA, histones and neutrophil extracellular traps exert anti-fibrinolytic effects in a plasma environment. *Thromb Haemost* 2015;113(06):1289–1298
- 209 Chang X, Yamada R, Sawada T, Suzuki A, Kochi Y, Yamamoto K. The inhibition of antithrombin by peptidylarginine deiminase 4 may contribute to pathogenesis of rheumatoid arthritis. *Rheumatology (Oxford)* 2005;44(03):293–298
- 210 Longstaff C, Varjú I, Sótónyi P, et al. Mechanical stability and fibrinolytic resistance of clots containing fibrin, DNA, and histones. *J Biol Chem* 2013;288(10):6946–6956
- 211 Xaus J, Comalada M, Valledor AF, et al. LPS induces apoptosis in macrophages mostly through the autocrine production of TNF-alpha. *Blood* 2000;95(12):3823–3831