Joint Guidance from the Society of Gastrointestinal Endoscopy of India (SGEI), Indian Society of Gastroenterology (ISG), and Indian National Association for Study of the Liver (INASL) for Gastroenterologists and Gastrointestinal Endoscopists on the Prevention, Care, and Management of Patients with COVID-19

Mathew Philip¹ Sundeep Lakhtakia² Rakesh Aggarwal³ Kaushal Madan⁴ Vivek Saraswat⁵
Govind Makharia⁶

¹President SGEI, HOD, Lisie Institute of Gastroenterology Lisie Hospital, Kochi

²Secretary SGEI, Director Endoscopy, Asian Institute of Gastroenterology, Hyderabad

³President INASL, Director, JIPMER, Puducherry

⁴Secretary, INASL, Director & Head, Gastroenterology & Hepatology, Max Smart Super Speciality Hospital, Saket, New Delhi

⁵President ISG, Professor & Head, Gastroenterology, Sanjay Gandhi Postgraduate Institute of Medical Sciences (SGPGIMS), Lucknow

⁶Secretary ISG, Professor, Gastroenterology, AIIMS, Delhi

J Digest Endosc 2019;10:201–205

Background

Coronavirus disease 2019 (COVID-19), which is caused by infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2, known also as novel coronavirus 2019), is currently occurring as a pandemic. It first appeared in December 2019 in Wuhan city, located in the Hubei region of China, and was soon followed by a quick spread to nearby provinces of China and to its neighboring countries. As of March 26, 2020, the infection has been reported from 198 countries and has affected more than 471,000 people worldwide, with more than 21,000 deaths (<https://www.worldometers.info/coronavirus/>).

COVID-19 most often presents with a recent-onset fever, dry cough, weakness, and sore throat. Up to 50% of patients may report shortness of breath, and a few develop acute respiratory distress syndrome. Nasal symptoms are infrequent. Asymptomatic infection can also occur; however, in the absence of a serological test, its frequency remains unclear. The case fatality rate has been reported between 1 and 3.5%, but may depend on case definition; for instance, if milder cases or asymptomatic persons are tested, diagnosed, and included in the case count, the mortality rate would

appear to be low. Human-to-human transmission occurs primarily through direct contact through air droplets. The mean incubation period is 5 days (range: 0–14 days). Spread from asymptomatic persons in the late incubation period can occur; however, most of the viral spread appears to occur from symptomatic persons. Older people and the immunocompromised individuals are at particular risk of severe disease and death.

Gastrointestinal (GI) symptoms including nausea and/or diarrhea have been reported to occur in 5 to 50% of infected individuals in various series. Liver enzymes are abnormal in a quarter of cases. Viral RNA is detectable in stool and may persist for longer than the acute illness; however, whether this represents the presence of viable virus and the risk of transmission remains unclear. Meanwhile, it appears prudent to consider GI secretions as infective, capable of causing fecal–oral transmission, and associated with a potential for transmission of the virus during endoscopic procedures from patient to patient or from a patient to health care workers (HCWs).

In GI endoscopy units, several staff members including physicians and other HCW often work at a very short physical distance from patients. Furthermore, they are frequently

Address for correspondence

Dr. Mathew Philip MD, DNB, DM,
DNB (GE) President, SGEIHOD &
Senior Consultant, Lisie Institute of
Gastroenterology Lisie Hospital, Kochi
(e-mail: drmathewphilip@gmail.com)

DOI <https://doi.org/10.1055/s-0040-1709837>
ISSN 0976-5042.

©2020 Society of Gastrointestinal
Endoscopy of India

License terms

exposed to splashes, air droplets, mucus, or saliva during GI endoscopy procedures. Endoscopy is potentially an aerosol producing procedure and the risk of exposure may be particularly high during intubation with an endoscope that can occasionally induce coughing or violent retching. Or, if unexpected respiratory adverse event occurs during endoscopy with or without the need of placement of an endotracheal tube.

The best personal protection techniques currently recommended at all times are as follows:

- Frequent and thorough handwashing (with soap and water or antiseptic handwash solutions, preferably those containing 60% alcohol).
- Avoiding touching one's face, mouth, or nose with unwashed hands.
- Following cough and sneezing etiquette.
- Maintaining physical distance from other people and avoiding crowds.

In addition, in health care settings including in endoscopy suites, wearing surgical masks by HCWs may help prevent exposure to infectious material from an infected patient source such as splashes, saliva, or mucus. Though this practice is very useful, it may not be sufficient enough to provide complete protection from exposure to the virus and other contaminants to the wearer.

With an increasing number of COVID-19 cases in India (673 cases including 13 deaths on March 26, 2020), it is felt that GI health professionals need to be aware of the disease and how to prevent COVID-19 transmission and manage patients during the ongoing COVID-19 pandemic.

Keeping this in view, the three Indian professional bodies in the field of GI disease, namely the Society of Gastrointestinal Endoscopy of India (SGEI), Indian Society of Gastroenterology (ISG), and Indian National Association for the Study of the Liver (INASL), have come up with this guidance for gastroenterologists and GI endoscopists who are involved in providing care to patients with GI and liver disease.

Since the available scientific evidence on the disease is scanty, these recommendations are mostly based on expert opinion and knowledge derived from other pathogens with similar characteristics. However, the guidance represents what is believed to be the best current understanding and prudent clinical practice and should generally serve the gastroenterology community well.

These recommendations are divided into two sections, namely (1) those related to endoscopic procedures and (2) other important aspects of patient care in the face of the COVID-19 pandemic.

Recommendations Related to Endoscopic Procedures

Scheduling of Endoscopic Procedures

Endoscopy procedures can be divided into three categories based on their urgency as follows:

- **Emergency endoscopic procedures:** procedures for patients with life-threatening conditions, for example, diagnostic or therapeutic endoscopic procedures in patients with acute upper GI or lower GI bleeding, removal of impacted foreign body, and therapeutic endoscopy in patients with cholangitis or GI perforations.
- **Urgent endoscopic procedures:** diseases/conditions in which the treating clinician feels that an endoscopic procedure will have a significant beneficial impact on clinical outcome over the next 1 month. Examples include drainage of an infected pancreatic fluid collection, diagnosis and staging of GI cancers, placement of a nasojejunal or percutaneous gastrostomy tube for nutritional support, drainage of malignant biliary obstruction, and placement of a stent for malignant luminal obstruction of the esophagus, colon, or duodenum.
- **Routine endoscopic procedures:** endoscopic procedures that do not fall in either of the aforementioned two categories, for example, all routine referrals for diagnostic endoscopy procedures, and endoscopic procedures for screening or surveillance.

It is recommended that only emergency and urgent endoscopy procedures may be undertaken for the next 4 weeks or until the current threat of COVID-19 lasts or further evidence becomes available. Routine endoscopy procedures can usually be safely postponed for 1 month, though such patients must be closely monitored for any change in clinical status that may change the need for endoscopy to "urgent" or "emergency." In such cases, alternative approaches (e.g., a radiological investigation or procedure) for diagnosis or treatment may also be explored since are less risky options.

All the three Indian gastroenterology societies (SGEI, ISG, and INASL) jointly recommend to consider only emergency and urgent endoscopy procedures for the next 1 month or till the current threat due to COVID-19 is over. Routine endoscopic procedures can be postponed for the next 4 weeks unless a change in a patient's clinical status mandates an emergency or urgent endoscopy in the intervening period.

Endoscopic Procedures

For any patient scheduled for endoscopy, the following steps are recommended during the preprocedure, procedure, and postprocedure phases.

Preprocedure Screening

In each patient scheduled for an endoscopic procedure, history of fever or respiratory symptoms, contact with a confirmed case of COVID-19, and a recent history of travel to or of living in an area with higher rate of transmission of COVID-19 disease should be obtained. Furthermore, for each such person, body temperature should be measured as a routine. Based on these parameters, the person should be categorized into one of the following three categories of risk of harboring SARS-CoV-2 infection: low risk, intermediate risk, and high risk.

Low Risk

- No symptom suggestive of COVID-19 (cough, fever, breathlessness, or diarrhea).
- No history of travel to or stay in a high-risk area* in the past 14 days (*a “high-risk area” implies an area where more than 1,000 cases have been confirmed till date; this is changing over time).
- No contact with a COVID-19 patient.

Intermediate Risk

- Symptoms present but no history of travel to or stay in a high-risk area during the past 14 days or of contact with a COVID-19 patient; or
- No symptom, but history of contact with a confirmed COVID-19 patient or stay in or travel to a high-risk area in the last 14 days.

High Risk

- At least one symptom present; and
- either contact with a confirmed COVID-19 patient or of stay in a high-risk area

In case of a possibility of intermediate or high risk of exposure to coronavirus, the need and urgency of the procedure must be reconsidered. In such cases, the procedure should generally be postponed unless there is an indication for emergency endoscopy. Furthermore, for persons with high-risk exposure or the presence of symptoms, follow the protocol recommended by the Ministry of Health and Family Welfare (MoHFW), Government of India.

In the Procedure Room

- The number of staff members present in the endoscopy area during the procedure should be reduced to the minimum required.
- All members of the endoscopy team should wear appropriate personal protective equipment (PPE), such as gloves, mask, eye shield/goggles, face shields, and gown, as appropriate, based on risk assessment and stratification and undertake adequate handwashing before and after handling the patients.
- For high-risk cases, ensure that appropriate PPE is available and worn by all members of the endoscopy team. In such cases, the sequence of wearing (donning) and removal (doffing) of PPE must follow the prescribed standard protocol.
- Data on the efficacy of commonly used chemical disinfection agents against SARS-CoV-2 are currently not available. However, since most of the other coronaviruses are inactivated by the commonly used disinfectants, it appears that no additional steps beyond those currently recommended for endoscope cleaning and reprocessing are needed. However, the recommended protocols for disinfection techniques for endoscope reprocessing must be strictly adhered to.
- As far as possible, only disposable endoscopic accessories should be used.

- Standard endoscopy room disinfection policy should be followed for non-COVID-19 or low-risk patients undergoing endoscopy.
- For patients with intermediate or high risk of COVID-19 infection, noncritical environmental surfaces frequently touched by hand (e.g., bedside tables, bed rails, cell phones, computers) and endoscopy furniture and floor should be disinfected at the end of each procedure.
- With a COVID-19 positive or very high-risk case with respiratory symptoms, the endoscopy may be performed in a negative-pressure room, if available.

Postprocedure Observation

- During patient observation in the postprocedure area or a recovery room, adequate spacing between beds (at least 6 feet) should be ensured.
- Surgical masks should be provided for patients with respiratory symptoms.

Other Recommendations Relevant to Gastroenterology Practice

Outpatient Clinics

- Nonurgent consultations and outpatient visits may be postponed or rescheduled for 4 weeks later (unless change in symptoms or clinical situation warrants an earlier visit during the intervening period).
- The policy of having only one accompanying person per patient should be insisted in consultation rooms, waiting areas, and for inpatients to prevent crowding.
- Information about COVID-19 must be displayed in the outpatient and other patient waiting areas with visuals recommending the dos and the don'ts.
- An appointment system should be instituted and followed so that the patients do not have to wait for a long time or to crowd in the outpatient or endoscopy waiting area.
- The electronic means of communications or telemedicine (such as phone calls, text messaging, WhatsApp, or other video calling applications) can be used for resolving minor queries and may help obviate a visit to the hospital or clinic, thereby reducing the risk of transmission of infection.

Academic Activities and Work Schedule of the Department

- It is ideal to follow the institutional policy regarding holding academic activities and the work schedule of the department.
- Rescheduling of department meetings or academic sessions involving more than 10 persons till the COVID-19 crisis is over should be considered.
- Fellow students and doctors should consider the use of text-messaging tools (e.g., WhatsApp) or social media

tools for communication and academic interaction between members of the gastroenterology team as well as other specialists. While sharing information about patients over such tools, the issues related to patient confidentiality must receive due attention.

- In the event of an outbreak in the department/hospital, it is most appropriate to follow the institutional guidance. It seems appropriate for each unit/department to have more than one team of doctors and other staff working on a rotation basis to ensure that it is able to provide uninterrupted service. A schedule may be drawn whereby one group attends the hospital for a specified number of days and the other group follows the next days. This may help avoid the risk of the whole department needing quarantine in case of a high-risk exposure to a patient or another HCW in the hospital, resulting in the entire department closing down.

Actions in Case of Exposure to a Health Care Worker to COVID-19

- If an HCW is exposed to a person at high risk of or a confirmed COVID-19 case, the hospital's infection control team should be informed immediately, and the guidelines set up by the MoHFW, Government of India, should be followed.
- Such workers may need quarantine for 14 days with self-monitoring and/or supervised guidance based on the risk stratification of the exposure.
- For asymptomatic HCWs involved in the care of suspected or confirmed cases of COVID-19, prophylactic treatment with hydroxychloroquine may be considered, as per the guidelines put forward by the Indian Council of Medical Research. The recommended dosage for this purpose is 400 mg (taken with meals) twice a day on day 1 followed by 400 mg once weekly for the next 7 weeks.

However, it is pertinent to point out that data to support this recommendation are limited to a French study in treatment (and not prophylaxis) setting, which had a nonrandomized nonblinded design with a small sample size (treated cohort of 26 and untreated cohort of 16 derived from different hospitals and hence not necessarily comparable) and different dropout rates (6/26 and 0/16, respectively) in the two cohorts. Of the 20 patients who received hydroxychloroquine, 7 also took azithromycin. Furthermore, it compared with a surrogate outcome (absence of viral RNA on day 6), and it was unclear whether this was decided *a priori* (before the study started).

Also, the use of chemoprophylaxis carries the risk of adverse events and instilling a false sense of security with reduced adherence to safety precautions. The recommendations for quarantine may change over time if the community spread of coronavirus becomes common.

Patients with Preexisting Digestive Diseases

- Patients on specific immunosuppressive treatment such as corticosteroids or cancer chemotherapy (e.g., in patients with inflammatory bowel disease, autoimmune

liver disease, transplant recipients) should contact their treating doctors for advice about the need to continue their treatment and for updated information.

- In patients with inflammatory bowel disease, there is no recommendation to pause the immunosuppressive treatment at the moment. Often, the risk of flare-up of the original disease may outweigh the chance of contracting COVID-19, necessitating the continuation of such drugs.
- All such patients should follow the guidelines of the MoHFW, Government of India, for the general public, which are meant to minimize exposure to the coronavirus disease, especially social distancing and frequent handwashing.
- Patients with cirrhosis (even Child A) and those with prior liver transplantation should be discouraged from visiting a clinic or hospital, unless absolutely essential.
- Patients with decompensated cirrhosis should be considered for inpatient treatment only if there is a pressing indication for admission, such as acute GI bleed, hepatic encephalopathy, tense ascites causing respiratory distress, or liver cancer requiring locoregional therapy or liver transplantation.
- Endoscopic variceal ligation as primary prophylaxis should be postponed till 4 to 6 weeks later or till the threat of COVID-19 infection has passed.
- Liver transplant recipients with COVID-19 infection should be monitored for drug-drug interactions, if they are prescribed lopinavir/ritonavir antiviral therapy (see AST Guidance).¹⁶

Each hospital or clinic should adopt measures, as locally suitable and acceptable and as per the regional or state policies and the local risk of occurrence of the COVID-19 outbreak.

As gastroenterologists, we should adopt steps to prevent the spread of this virus and to protect ourselves, our staff, coworkers, and their family members, and the population at large, while imparting quality care to our patients.

Conflict of Interest

None declared.

Bibliography

- 1 Government of India, Ministry of Health & Family Welfare, Directorate General of Health Services (EMR Division). Guidelines on Clinical Management of COVID-19. Available at: <https://www.mohfw.gov.in/pdf/GuidelinesonClinicalManagementofCOVID1912020.pdf>. Accessed March 16, 2020
- 2 American Gastroenterological Association. Joint GI society message: COVID-19 clinical insights for our community of gastroenterologists and gastroenterology care providers. Available at: <https://www.gastro.org/press-release/joint-gi-society-message-covid-19-clinical-insights-for-our-community-of-gastroenterologists-and-gastroenterology-care-providers>. Accessed March 26, 2020
- 3 Repici A, Maselli R, Matteo, R, et al. Coronavirus (COVID-19) outbreak: what the department of endoscopy

- should know. *Gastrointest Endosc* 2020 (e-pub ahead of print). doi:10.1016/j.gie.2020.03.019
- 4 Gu J, Han B, Wang J. COVID-19: gastrointestinal manifestations and potential fecal-oral transmission. *Gastroenterology* 2020 (e-pub ahead of print). doi:10.1053/j.gastro.2020.02.054
 - 5 Xiao F, Tang M, Zheng X, et al. Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterology*, 2020. pii: S0016-5085(20)30282-1. <https://doi.org/10.1053/j.gastro.2020.02.05>
 - 6 Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; 395(10223):497–506
 - 7 World Endoscopy Organization. WEO ALERT: Wuhan proposal for Safety in Digestive Endoscopy. Available at: <http://www.worldendo.org/2020/02/05/weo-alert-wuhan-proposal-for-safety-in-digestive-endoscopy/>. Accessed March 26, 2020
 - 8 Centers for Disease Control and Prevention. Interim U.S. Guidance for Risk Assessment and Public Health Management of Healthcare Personnel with Potential Exposure in a Healthcare Setting to Patients with Coronavirus Disease (COVID-19). Available at: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assessment-hcp.html>. Accessed March 26, 2020
 - 9 British Society of Gastroenterology. Advice for Endoscopy Teams during COVID-19. Available at: <https://www.bsg.org.uk/wp-content/uploads/2020/03/Advice-for-Endoscopy-Teams-during-COVID-ver-2-4-published-22032020FINAL-1.pdf>. Accessed March 26, 2020
 - 10 World Endoscopy Organization. Advice to Endoscopists. Available at: http://www.worldendo.org/wp-content/uploads/2020/03/WEO_Advice_To_Endoscopists_COVID-19_032020.pdf. Accessed March 26, 2020
 - 11 Mao R, Liang J, Shen J, et al; Chinese Society of IBD, Chinese Elite IBD Union; Chinese IBD Quality Care Evaluation Center Committee. Implications of COVID-19 for patients with pre-existing digestive diseases. *Lancet Gastroenterol Hepatol* 2020 (e-pub ahead of print). doi:10.1016/S2468-1253(20)30076-5
 - 12 Das Gastroenterologie Portal. 2nd Interview COVID-19 ECCO Taskforce. Available at: http://dasgastroenterologieportal.de/Interview_der_ECCO_zu_COVID_19.html. Accessed March 26, 2020
 - 13 Indian Council of Medical Research. Recommendation for empiric use of hydroxy-chloroquine for prophylaxis of SARS-CoV-2 infection. Available at: https://icmr.nic.in/sites/default/files/upload_documents/HCO_Recommendation_22March_final_MM.pdf. Accessed March 26, 2020
 - 14 Danese S, Cecconi M, Spinelli A. Management of IBD during the COVID-19 outbreak: resetting clinical priorities. *Nat Rev Gastroenterol Hepatol* 2020 (e-pub ahead of print). doi:10.1038/s41575-020-0294-8
 - 15 Gautreta P, Lagiera J-C, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an openlabel non-randomized clinical trial. *Int J Antimicrob Agents* 2020 (e-pub ahead of print). doi:10.1016/j.ijantimicag.2020.105949
 - 16 Information for transplant professionals and community members regarding 2019 novel coronavirus at the American Society of Transplantation website: <https://www.myast.org/information-transplant-professionals-and-community-members-regarding-2019-novel-coronavirus>

From the Editor-in-Chief's Desk

This is a position paper prepared in March 2020 jointly by three Gastroenterology societies of India as per available evidence on COVID. However as more data pours in, there may be some changes in our position in near future. Please visit our website www.sgei.co.in for latest information.

Dr Mahesh Goenka.

Editor-in-Chief.