A Rare Cause of Hand Infection: *Streptococcus Constellatus*

Una rara causa de infección de la mano: *Streptococcus Constellatus*

Rúben Malcata Nogueira¹ Carolina Vasconcelos² Nelson Teixeira¹

¹ Department of Reconstructive and Aesthetic Surgery, Centro Hospitalar Lisboa Ocidental, Lisboa, Portugal

² Department of Reconstructive and Aesthetic Surgery, Hospital Garcia de Orta, Almada, Portugal

Address for correspondence Rúben Malcata Nogueira, MD, Rua 25 de Abril, 8, 2600-541, Alhandra, Portugal (e-mail: malcatanogueira@gmail.com).

Rev Iberam Cir Mano 2020;48:53–55.

Abstract

Hand infections represent common medical and surgical challenges that endanger delicate structures with severe consequences if not promptly addressed. Early identification and management are essential to achieve optimal outcomes. To the best of our knowledge this is the first reported case caused by *Streptococcus constellatus*, in which a severe periungueal infection evolved rapidly with septic thrombosis of digital vessels and culminated in amputation. This microorganism belongs to a group of commensal bacteria, *Streptococcus milleri*, that causes dental, peritonsillar and sinus abscesses. When bacteriemia outbursts, distant abscesses may form or endocarditis may ensue. A missed diagnosis and treatment can induce important morbidities, often delayed by the difficult isolation of the agent in the laboratory and its complex mechanisms of antibacterial resistance. This article focus on the importance of identifying serious hand infections requiring urgent or emergent treatment, since delayed or inadequate identification and management can lead to important and permanent deficits.

Keywords

- hand infection
- streptococcus
- *streptococcus constellatus*
- milleri

Resumen

Las infecciones de las manos representan desafíos médicos y quirúrgicos comunes que ponen en peligro estructuras delicadas con graves consecuencias si no se abordan con prontitud. La identificación y el manejo tempranos son esenciales para lograr resultados óptimos. Hasta donde sabemos, este es el primer caso reportado causado por *Streptococcus constellatus*, en el cual una infección periungueal severa evolucionó rápidamente con trombosis séptica de vasos digitales y culminó en amputación. Este microorganismo pertenece a un grupo de bacterias comensales, *Streptococcus milleri*, que causa abscesos dentales, periamigdalinos y sinusales. Cuando estalla la bacteriemia, pueden formarse abscesos distantes o puede producirse endocarditis. Un diagnóstico y tratamiento fallados pueden inducir morbilidades importantes, a menudo retrasadas por el difícil aislamiento del agente en el laboratorio y sus complejos mecanismos de resistencia antibacteriana. Este artículo se centra en la importancia de identificar infecciones graves de las manos que requieren tratamiento urgente o de emergencia, ya que la identificación y el manejo retrasados o inadecuados pueden conducir a déficits importantes y permanentes.

Palabras clave

- infección de la mano
- estreptococo
- streptococcus constellatus
- Milleri

received
September 8, 2019
accepted
March 10, 2020

DOI <https://doi.org/10.1055/s-0040-1712093>.
ISSN 1698-8396.

Copyright © 2020 Thieme Revinter
Publicações Ltda, Rio de Janeiro, Brazil

License terms

Introduction

Hand infections represent common medical and surgical challenges that often endanger delicate anatomical structures, with severe consequences if not treated promptly. Alexander Fleming's discovery of Penicillin in 1929 had a revolutionary impact on the way this entity is addressed. The early institution of appropriate antibiotherapy has considerably reduced short- and long-term morbidity, and virtually eliminated mortality. However, medical treatment is often not enough, and timely debridement and surgical drainage must always be considered.¹

In our Emergency Department, hand infections are increasingly frequent, mainly in the form of eponychia, cellulite, felons or tenosynovitis. The most common mechanism is direct inoculation, either caused by animal or human bites, stings or use of intravenous substances. The occurrence of infection depends not only from individual agent virulence factors, but also from the host susceptibility and immune status. Human immunodeficiency virus infection, uncontrolled or undiagnosed diabetes mellitus, immunosuppressor drugs use and toxic substance abuse are among the most frequent comorbidities found in our patient population. Besides their increased susceptibility, they also present frequently with polymicrobial infections or antibiotic-resistant agents, or even with subtle initial presentations, delaying the diagnosis and management or leading to a poorer result.

The most frequently isolated microorganism in hand infections is *Staphylococcus*. However, whenever possible, hand infections should have a microbiological analysis, with the identification of the causative organism, as well as its antibiotic susceptibility profile. Some bacteria, like *S. constellatus*, cause an extremely aggressive clinical course, that can lead to disastrous outcomes if not timely diagnosed and treated.² The occurrence of these rare infections underline the need to institute culture-based antibiotherapy and to complement it with thorough surgical debridement.

Case Report

A 55-year-old man with previously undiagnosed type 2 diabetes mellitus and heavy smoking habits presented with a 2-week history of painful and swollen third digit of his left hand. There was no history of a penetrating injury to that finger, paresthesias or systemic signs such as fever or chills. The patient reported only a mild periungueal infection some days before these symptoms ensued. At presentation, there was necrotic tissue both on distal and middle phalanxes, especially on the dorsal aspect, and serous fluid drained spontaneously from scattered punctiform wounds (►Fig. 1). White blood cell count was 20,000/u/l with neutrophilia, C-Reactive Protein was 14-mg/dl, and glucose level was 275mg/dl. X-rays excluded the presence of foreign bodies or osteomyelitis.

A large amount of mobile debris communicated with the volar subcutaneous tissues and a comprehensive surgical debridement was performed on the day of admission, and collected material was sent for microbiological analysis. The patient was admitted to the Plastic Surgery Department, and

Fig. 1 Hand at presentation.

rigorous glycemic control and empiric broad-spectrum intravenous antibiotic treatment were instituted. Areas of necrosis progressively defined with daily chemical and mechanical debridement (►Fig. 2). On day four of inpatient admission, microbiological results revealed a *Streptococcus constellatus* infection, and antibiotherapy was switched to Ampicillin and Clindamycin, based on susceptibility test. Despite the undertaken measures, necrosis progressed and affected also the proximal phalanx. One week after presentation, amputation of digital ray was performed and healed uneventfully.

Discussion

To the authors' best knowledge, this is the first reported case of *Streptococcus constellatus* hand infection leading to digital amputation. This microorganism belongs to a group of commensal bacteria, *Streptococcus milleri* (eponym for Willoughby Dayton Miller), which has undergone several changes in its taxonomic classification over the last years. Originally named *Streptococcus anginosus* (SAG) group, it consists of three species: *Streptococcus anginosus*, *Streptococcus constellatus* and *Streptococcus intermedius*. They are gram-positive organisms of the microaerophilic C-β hemolytic group, often found in the oral cavity and in the gastrointestinal and urogenital tracts. Although much less frequent than other subspecies, *Streptococcus constellatus* can be found in dental biofilms, which can at least in part explain its role in periungueal infection in nail bitters.³⁻⁵

During the last decade, there has been a growing number of infections caused by these agents, a fact that emphasizes their relevance and high pathogenicity, particularly as cause of abscesses, bacteremia and in patients with cystic fibrosis, especially in those with immunological commitment.^{1,6,7}

Fig. 2 After first surgical debridement.

A missed diagnosis can induce important morbidities, and it is often delayed by the difficult isolation of the agent in the laboratory, aggravated by the frequent assumption of contamination instead of infection, particularly in polymicrobial infections, that leads to undiagnosed and inappropriately treated cases. Several virulence factors, already described for other related agents, seem to also play a role in the pathogenicity of *Streptococcus constellatus*. Its adhesion capacity allows the attachment to components of extracellular matrix, such as fibrinogen or fibronectin, and the connection to other bacteria; the capsule production mechanism, which is linked to the formation of abscesses; and the production of β -hemolysins, all contribute to its high pathogenicity.^{1,7,8}

To date, hand infections caused by *S. constellatus* are described infrequently, and we could not find a similar

case with such aggressivity and rapid and ramping evolution. The case presented was caused, not only by the pathogenicity and virulence of the microorganism, but also by the host susceptibility, which the authors believe contributed enormously to the adverse outcome. Despite appropriate and aggressive medical and surgical treatment, we could not avoid the digital amputation, which emphasizes the need to adequately and timely diagnose and treat such infections.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Siegman-Igra Y, Azmon Y, Schwartz D. Milleri group streptococcus—a stepchild in the viridans family. *Eur J Clin Microbiol Infect Dis* 2012;31(09):2453–2459
- 2 Rashid RM, Salah W, Parada JP. 'Streptococcus milleri' aortic valve endocarditis and hepatic abscess. *J Med Microbiol* 2007;56(Pt 2):280–282
- 3 Tsunashima H, Miyake K, Motono M, Iijima S. Organization of the capsule biosynthesis gene locus of the oral streptococcus *Streptococcus anginosus*. *J Biosci Bioeng* 2012;113(03):271–278
- 4 Jensen A, Hoshino T, Kilian M. Taxonomy of the Anginosus group of the genus *Streptococcus* and description of *Streptococcus anginosus* subsp. *whileyi* subsp. nov. and *Streptococcus constellatus* subsp. *viborgensis* subsp. nov. *Int J Syst Evol Microbiol* 2013; 63(Pt 7):2506–2519
- 5 Fuentes Olmo J, Ruiz Mariscal M, Uribarrena Amezaña R, Omiste Sanvicente T. [Spontaneous bacterial peritonitis due to *Streptococcus constellatus*]. *Gastroenterol Hepatol* 2006;29(09): 595–596
- 6 Reissmann S, Friedrichs C, Rajkumari R, et al. Contribution of *Streptococcus anginosus* to infections caused by groups C and G streptococci, southern India. *Emerg Infect Dis* 2010;16(04): 656–663
- 7 Claridge JE III, Attorri S, Musher DM, Hebert J, Dunbar S. *Streptococcus intermedius*, *Streptococcus constellatus*, and *Streptococcus anginosus* ("Streptococcus milleri group") are of different clinical importance and are not equally associated with abscess. *Clin Infect Dis* 2001;32(10):1511–1515
- 8 Deutschmann MW, Livingstone D, Cho JJ, Vanderkooi OG, Brookes JT. The significance of *Streptococcus anginosus* group in intracranial complications of pediatric rhinosinusitis. *JAMA Otolaryngol Head Neck Surg* 2013;139(02):157–160