Aortoiliac Occlusive Disease: When the Development of Arterial Collateral Network Takes Over

Efstratios Georgakarakos, MD, MSc, PhD¹ Ioannis Katsaros, MD² Aliko Fiska, MD, PhD²

¹Department of Vascular Surgery, "Democritus" University of Thrace, University Hospital of Alexandroupolis, Alexandroupolis, Greece

²Department of Anatomy, Medical School of Alexandroupolis, Democritus University of Thrace, Alexandroupolis, Greece

Address for correspondence Efstratios Georgakarakos, MD, MSc, PhD, Department of Vascular Surgery, Democritus University of Thrace, 68100 Dragana, Alexandroupolis, Greece (e-mail: efstratiosgeorg@gmail.com).

AORTA 2020;8:116–117.

Abstract

Keywords

- aortoiliac occlusive disease
- Leriche's syndrome
- collateral circulation
- anatomy

This report describes the collateral pathways that restore arterial circulation in cases of aortoiliac occlusive disease and discusses the clinical and surgical importance of these systemic-systemic, visceral-systemic, and visceral-visceral anastomoses.

An 85-year old male presented with severe intermittent claudication bilaterally. Computed tomography angiography showed complete occlusion of the common and external iliac artery on the left and occlusion of the external iliac artery on the right (► **Fig. 1**). In such cases, the perfusion of limbs is attributed to an extensive collateral network between the systemic-systemic, visceral-systemic, and visceral-visceral pathways.^{1,2} The celiac, superior, and inferior mesenteric arteries form collateral pathways (white arrowhead) that perfuse the hypogastric arteries through the superior rectal and the middle sacral arteries (white arrow) or the branches (yellow arrowhead) of the common and deep femoral arteries via the pelvis transverse collateral

pathway (obturator, internal pudendal, and symphyseal arteries). The adequacy of this mechanism depends on the enlargement of preexisting vessels, although individuals differ in their capacity to develop collateral vessels.³ Age, diabetes, and hypertension suppress collateral development. Oxidative stress and endothelial dysfunction influence the capacity for collateral growth, while increased wall shear stress and circumferential wall tension due to dilation can lead to the luminal expansion and medial thickening observed in collateral arteries.³ These factors can be modified by a supervised training/walking program and pharmaceutical agents, such as statins and angiotensin-converting enzyme-inhibitors.

received
January 2, 2019
accepted after revision
May 24, 2020

DOI <https://doi.org/10.1055/s-0040-1714078>.
ISSN 2325-4637.

Copyright © 2020 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.
Tel: +1(212) 760-0888.

License terms

Fig. 1 Computed tomography angiography of the patient with complete occlusion of the left common and external iliac artery and occlusion of the right external iliac artery. A complex network of systemic and visceral anastomotic vessels restores the flow to the lower limbs (for details see in the text).

Funding

None.

Conflict of Interest

The authors declare no conflict of interest related to this article.

Acknowledgments

None.

References

- 1 Hardman RL, Lopera JE, Cardan RA, Trimmer CK, Josephs SC. Common and rare collateral pathways in aortoiliac occlusive disease: a pictorial essay. *AJR Am J Roentgenol* 2011;197(03):W519-24
- 2 Ahmed S, Raman SP, Fishman EK. CT angiography and 3D imaging in aortoiliac occlusive disease: collateral pathways in Leriche syndrome. *Abdom Radiol (NY)* 2017;42(09):2346–2357
- 3 Unthank JL, Sheridan KM, Dalsing MC. Collateral growth in the peripheral circulation: a review. *Vasc Endovascular Surg* 2004;38(04):291–313