Identification and Sensitivity of Vaginal and Probiotic *Lactobacillus* species to Urinary Antibiotics

Lipika Singhal¹ Varsha Gupta¹ Menal Gupta¹ Poonam Goel² Jagdish Chander¹

¹Department of Microbiology, Government Medical College and Hospital, Chandigarh, India

²Department of Obstetrics and Gynaecology, Government Medical College and Hospital, Chandigarh, India

Address for correspondence Menal Gupta, MBBS, MD, Department of Microbiology, Government Medical College and Hospital, Sector 32, Chandigarh 160030, India (e-mail: m.gupta1013@gmail.com).

J Lab Physicians:2020;12:111–114

Abstract

Objectives Healthy vaginal microbiota is mainly dominated by *Lactobacillus* species namely *L. crispatus*, *L. gasseri*, *L. jensenii*, and *L. iners*. Lactobacilli are thought to play an important role in the prevention of urogenital infections, and *Lactobacillus* probiotics to restore and/or maintain vaginal health has been advocated. These can interfere with the adherence, growth, and colonization by uropathogenic bacteria, thus reducing the risk of urinary tract infection (UTI). This study aims to isolate and evaluate the susceptibility of healthy vaginal and probiotic *Lactobacillus* spp. to urinary antibiotics.

Materials and methods A total of 50 premenopausal, nonmenstruating women with no symptoms of vaginal infection or UTI or antimicrobial use in the past 2 weeks were enrolled. Two high vaginal swabs were collected for Nugent's scoring and anaerobic culture. Colonies yielding gram-positive rods were confirmed by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS). *Lactobacillus* strains in a vaginal probiotic tablet were also isolated, and the sensitivity of both sets to urinary antibiotics was determined.

Results A total of 29 *Lactobacillus* spp. were isolated, including *L. crispatus* (15), *L. gasseri* (5), *L. vaginalis* (4), *L. paracasei* (2), *L. curvatus* (1), *L. fermentum* (1), and *L. paraplantarum* (1). All strains were susceptible to nitrofurantoin and resistant to norfloxacin, gentamicin, and cotrimoxazole (except *L. paracasei*). The probiotic strains were pan-sensitive.

Conclusions Prophylactic antibiotics are capable of eliminating the normal vaginal inhabitants, which may increase the probability of UTI. The administration of vaginal probiotics as an alternate or multidrug therapy can restore vaginal microbiota and help prevent recurrent UTI.

Keywords

- *Lactobacillus*
- probiotics
- UTI
- MALDI-TOF

Introduction

The vaginal microbiome has evolved to protect women against a variety of urogenital infections. Healthy vaginal microbiota is mainly dominated by *Lactobacillus* species, namely *L. crispatus*, *L. gasseri*, *L. jensenii*, and *L. iners*.¹ Depletion of *Lactobacilli* may result in bacterial vaginosis (BV), which is associated with urinary tract infections (UTIs), increased risk of sexually transmitted infections, and pelvic inflammatory disease.² UTI is a common problem among young

healthy women of reproductive age group, with approximately 25 to 30% reporting recurrence. Recurrent UTI (RUTI) is defined as at least three episodes of UTI in 12 months or at least two episodes in 6 months. *Escherichia coli* is the predominant pathogen in UTIs followed by *Staphylococcus saprophyticus*, *Enterococcus faecalis*, *Klebsiella pneumoniae*, and *Proteus mirabilis*. Antibiotics are effective in the treatment of UTIs but lead to increased antibiotic resistance in microorganisms.³ Antibiotic prophylaxis with agents such as cotrimoxazole, nitrofurantoin, and fluoroquinolones has been

associated with a decreased rate of recurrence, from 2 to 3 cases per patient-year to 0.1 to 0.2 cases per patient-year, but it disrupts normal flora of urinary tract and increases development of resistance in uropathogens.⁴ The emergence of extended-spectrum β -lactamases and carbapenem-resistant *E. coli* emphasize the need to explore novel agents that have fewer side effects and promise long-term benefit.⁵ In case of RUTIs, disruption of normal vaginal flora has been shown to predispose patients to chronic intermittent urogenital infections.⁶ In this context, researchers have reported that the use of probiotics as an adjunct to antibiotic therapy may provide higher antimicrobial activity and reduce the dose of antibiotics required in addition to replenishing the intestinal flora and decreasing antibiotic side effects.⁵ Use of *Lactobacillus*-containing probiotics to restore commensal vaginal flora has been proposed for the treatment and prophylaxis of bacterial urogenital infections. Florisia (CD Pharma India Pvt. Ltd.) is an intravaginal tablet containing at least 10^9 CFU (colony-forming unit) of lyophilized *L. brevis*, *L. salivarius* subsp. *salicinus*, and *L. plantarum*. The three strains have been selected for their ability to adhere to vaginal epithelial cells, production of H_2O_2 , and coaggregation with pathogens.² This study aims to isolate and identify the *Lactobacillus* species from the vagina of healthy females and probiotic tablet and to evaluate their susceptibility to urinary antibiotics.

Materials and Methods

This study was conducted from December 2016 to December 2017 with approval of the Institutional Ethical Committee and informed consent of all participants. A total of 50 premenopausal, nonmenstruating females (18–45 years old) who had no symptoms of UTI or vaginal infection were enrolled. Women with chronic autoimmune or inflammatory conditions or on oral/topical antimicrobials in the past 2 weeks were excluded. Two high vaginal swabs were collected from each female. Gram stain of direct smear was performed and graded on a 10-point scale based on the presence of *Lactobacilli* and other anaerobes as described by Nugent. The vaginal flora was defined as “healthy” if the score was 0 to 3, “intermediate” if 4 to 6, and “BV” if > 7 .⁷ Only women diagnosed as “healthy” were included in the analyses. The second swab from these “healthy” women was inoculated onto de Man, Rogosa, and Sharpe (MRS) and Brain Heart Infusion agar plates (HiMedia Laboratories, Mumbai, India). Plates were incubated anaerobically for 24 hours at 37°C in anaerobic jars with gaspak.⁸ White, mucoid, catalase-negative colonies on MRS agar yielding gram-positive rods were isolated for confirmation by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS, Bruker Daltonics, Hamburg, Germany). The probiotic tablet was incubated anaerobically at 37°C in MRS broth for 48 hours and then isolation on MRS agar was performed. Antimicrobial susceptibility of *Lactobacillus* isolates from healthy females and probiotic strains to antibiotics co-trimoxazole, norfloxacin, nitrofurantoin and gentamicin was determined by the method of Bauer et al using MRS agar under anaerobic conditions.⁹

Results

Of the participants, 66% ($n = 33$) had Nugent's scores of 0 to 3 (► **Table 1**). A total of 29 species of *Lactobacillus* were isolated from these: most common being *L. crispatus* ($n = 15$) followed by *L. gasseri* ($n = 5$) and *L. vaginalis* ($n = 4$). The other species isolated were *L. paracasei* ($n = 2$), *L. curvatus* ($n = 1$), *L. fermentum* ($n = 1$), and *L. paraplantarum* ($n = 1$). All these isolates were susceptible to nitrofurantoin and resistant to norfloxacin and gentamicin. All the isolates were resistant to cotrimoxazole except *L. paracasei*. The probiotic tablet yielded *L. brevis*, *L. salivarius*, and *L. plantarum*, which were susceptible to all antibiotics.

Discussion

Lactobacilli are the predominant bacteria in vaginal flora and possess antimicrobial properties that regulate other urogenital microbiota.⁶ Most UTIs are caused by intestinal bacteria that ascend through the urethra to the bladder and, sometimes, kidneys.³ The *Lactobacillus*-dominated vaginal flora in premenopausal women impedes colonization of uropathogens due to competitive exclusion and maintaining low vaginal pH.⁵ Inverse association has been reported between H_2O_2 -producing *Lactobacilli* and vaginal *E. coli* colonization in women with RUTIs. Specific *Lactobacilli* strains can interfere with the adherence, growth, and colonization of uropathogenic bacteria and thus reduce the risk of ascent into the bladder.¹⁰ There are only a few studies characterizing the vaginal *Lactobacillus* species in healthy women of reproductive age in India (► **Table 2**). More studies in India are warranted as the species distribution in Indian women may be different. This would also guide as to whether replacement by a single species is sufficient or a balanced mixture depending on the local flora is needed in a probiotic.

The species most frequently isolated in our study was *L. crispatus*, which is a strong H_2O_2 producer and is associated with healthy vaginal microbiota. In addition, *L. vaginalis* was isolated, which has been associated with healthy flora. The other species was *L. gasseri*, and females carrying this are reported to have higher Nugent scores, though asymptomatic.¹⁴ Women in different geographical settings face different environmental conditions, dietary habits, and lifestyle.¹ The results of our study resemble the most prevalent vaginal species in other Indian studies (► **Table 2**). Several authors have reported varying observations of vaginal *Lactobacilli* elsewhere in the world.^{14–17} Despite reports of high incidence of *L. iners* in human vaginal microbiota, we did not obtain isolates belonging to this species probably because of stringent nutritional requirements, very low oxygen tolerance, and molecular-based identification.¹⁵

Table 1 Nugent score of participants

Nugent score	Category	Participants (%)
0–3	Healthy	33 (66%)
4–6	Intermediate	10 (20%)
> 7	Bacterial vaginosis	7 (14%)

Table 2 Comparison of *Lactobacillus* spp. isolated in various studies from India

	Garg et al ¹¹	Madhivanan et al ⁸	Madhivanan et al ¹²	Pramanick et al ¹³	Das Purkayastha et al ¹
	n = 80	n = 11	n = 39	n = 107	n = 26
<i>Lactobacillus reuteri</i>	33%	9%	23%	22%	1%
<i>Lactobacillus fermentum</i>	25%	9%	15%	–	6%
<i>Lactobacillus salivarius</i>	16%	9%	–	–	–
<i>Lactobacillus crispatus</i>	5%	27%	41%	27%	–
<i>Lactobacillus gasseri</i>	3%	–	46%	15%	4%
<i>Lactobacillus jensenii</i>	4%	27%	18%	16%	2%
<i>Lactobacillus iners</i>	–	–	–	65%	–
<i>Lactobacillus mucosae</i>	–	–	5%	–	16%

Several in vitro and in vivo studies support the beneficial effect of *Lactobacilli* on the restoration of vaginal flora and the prevention of RUTIs. The rationale for the use of probiotics is based on the regulatory role played by commensal microflora in the gastrointestinal and genitourinary tracts.⁶ In their meta-analysis, Grin et al concluded that probiotic strains of *Lactobacillus* are safe and effective in preventing RUTI in adult women.¹⁸ Barrons and Tassone concluded that intravaginal *Lactobacillus* suppositories are effective against uropathogens and show the greatest efficacy for UTI prophylaxis.⁶ In another study, the use of *Lactobacillus* vaginal suppository in women resulted in the reduction of *E. coli* positive cultures from 5.0 ± 1.6 episodes to 1.3 ± 1.2 episodes ($p < 0.0007$) over a 12-month period.¹⁹

Prolonged low-dose chemotherapeutic agents such as nitrofurantoin, ciprofloxacin, trimethoprim, and cotrimoxazole have been traditionally used as prophylaxis in reducing UTI.³ All *Lactobacillus* isolates were found resistant to cotrimoxazole except the *L. paracasei* strains. All strains were susceptible to nitrofurantoin and resistant to norfloxacin and gentamicin. The three probiotic strains were found susceptible to all the tested antibiotics. This suggests that prophylactic antibiotics are capable of eliminating the normal vaginal inhabitants, which are the main inhibitors of urinary pathogens. This way they counteract the benefits of antibacterial effect and increase the probability of UTI.

Conclusion

We suggest that the administration of vaginal probiotics can restore vaginal microbiota and help prevent recurrence of UTI, particularly as part of an alternate therapy or multidrug treatment. The limitation of our study is the small sample size. Also, as our methodology was culture-based, only those *Lactobacillus* species that can be cultured were identified. To the best of our knowledge, very few similar studies have been performed in India, and large-scale studies in future would help in generating conclusive evidence for alternative therapies.

Ethical Approval

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or

national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Funding

This study was part of a research project funded by the Department of Science and Technology, Government of India.

Presentation at a Meeting

North-West Chapter of the Indian Association of Medical Microbiologists, Rohtak, Haryana, India.

Conflict of Interest

Dr. Chander reports grants from the Department of Science and Technology, Government of India, during the conduct of the study. Dr. V. Gupta reports grants from the Department of Science and Technology, Government of India, during the conduct of the study. Dr. Goel reports grants from the Department of Science and Technology, Government of India, during the conduct of the study. Dr. Singhal reports grants from the Department of Science and Technology, Government of India, during the conduct of the study. Dr. M. Gupta reports grants from the Department of Science and Technology, Government of India, during the conduct of the study.

Acknowledgment

The authors gratefully acknowledge the Department of Medical Microbiology, Postgraduate Institute of Medical Education and Research, Chandigarh, India, for their help in species identification by MALDI-TOF MS.

References

- 1 Das Purkayastha S, Bhattacharya MK, Prasad HK, et al. Contrasting diversity of vaginal *Lactobacilli* among the females of Northeast India. *BMC Microbiol* 2019;19(1):198
- 2 Mastromarino P, Hemalatha R, Barbonetti A, et al. Biological control of vaginosis to improve reproductive health. *Indian J Med Res* 2014;140(Suppl):S91–S97
- 3 Al-Badr A, Al-Shaikh G. Recurrent urinary tract infections management in women - a review. *Sultan Qaboos Univ Med J* 2013;13(3):359–367

- 4 Reid G, Bruce AW, Cook RL, Llano M. Effect on urogenital flora of antibiotic therapy for urinary tract infection. *Scand J Infect Dis* 1990;22(1):43–47
- 5 Gupta V, Nag D, Garg P. Recurrent urinary tract infections in women: how promising is the use of probiotics? *Indian J Med Microbiol* 2017;35(3):347–354
- 6 Barrons R, Tassone D. Use of *Lactobacillus probiotics* for bacterial genitourinary infections in women: a review. *Clin Ther* 2008;30(3):453–468
- 7 Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol* 1991;29(2):297–301
- 8 Madhivanan P, Raphael E, Rumphs A, et al. Characterization of culturable vaginal *Lactobacillus* species among women with and without bacterial vaginosis from the United States and India: a cross-sectional study. *J Med Microbiol* 2014;63(Pt 7) : 931–935
- 9 Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 1966;45(4):493–496
- 10 Gupta K, Stapleton AE, Hooton TM, Roberts PL, Fennell CL, Stamm WE. Inverse association of H2O2-producing *Lactobacilli* and vaginal *Escherichia coli* colonization in women with recurrent urinary tract infections. *J Infect Dis* 1998;178(2): 446–450
- 11 Garg KB, Ganguli I, Das R, Talwar GP. Spectrum of *Lactobacillus* species present in healthy vagina of Indian women. *Indian J Med Res* 2009;129(6):652–657
- 12 Madhivanan P, Alleyn HN, Raphael E, et al. Identification of culturable vaginal *Lactobacillus* species among reproductive age women in Mysore, India. *J Med Microbiol* 2015;64(6):636–641
- 13 Pramanick R, Parab S, Mayadeo N, Warke H, Aranha C. Cross sectional analysis of vaginal *Lactobacillus* in asymptomatic women of reproductive age in Mumbai, India. *J Infect Dev Ctries* 2018;12(12):1096–1104
- 14 Vásquez A, Jakobsson T, Ahrné S, Forsum U, Molin G. Vaginal *Lactobacillus* flora of healthy Swedish women. *J Clin Microbiol* 2002;40(8):2746–2749
- 15 Pino A, Bartolo E, Caggia C, Cianci A, Randazzo CL. Detection of vaginal *Lactobacilli* as probiotic candidates. *Sci Rep* 2019;9(1):3355
- 16 Zhang R, Daroczy K, Xiao B, Yu L, Chen R, Liao Q. Qualitative and semiquantitative analysis of *Lactobacillus* species in the vaginas of healthy fertile and postmenopausal Chinese women. *J Med Microbiol* 2012;61(Pt 5) :729–739
- 17 Pendharkar S, Magopane T, Larsson PG, et al. Identification and characterisation of vaginal *Lactobacilli* from South African women. *BMC Infect Dis* 2013;13:43
- 18 Grin PM, Kowalewska PM, Alhazzan W, Fox-Robichaud AE. *Lactobacillus* for preventing recurrent urinary tract infections in women: meta-analysis. *Can J Urol* 2013;20(1):6607–6614
- 19 Uehara S, Monden K, Nomoto K, Seno Y, Kariyama R, Kumon H. A pilot study evaluating the safety and effectiveness of *Lactobacillus* vaginal suppositories in patients with recurrent urinary tract infection. *Int J Antimicrob Agents* 2006;28(Suppl 1) :S30–S34