Psychological Follow-up During Prenatal Care of Pregnant Women: Insights During the covid-19 pandemic

Magda Spinello Consul Da Silva¹ Edward Araujo Júnior¹ Julio Elito Júnior¹

¹ Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, SP, Brazil

Address for correspondence Edward Araujo Júnior, PhD, Rua Napoleão de Barros, 875, Vila Clementino, São Paulo, SP, 04024-002, Brazil (e-mail: araujojred@terra.com.br).

Rev Bras Ginecol Obstet 2021;43(1):72–73.

Dear Editor,

The puerperal pregnancy cycle is considered a period of crisis marked by emotional changes in women and men, and may evolve healthily or present psychological disorders that will compromise psychological, physical and marital health. Psychological prenatal care is the monitoring of the pregnant and/or of the couple during pregnancy, the postpartum period and the puerperium until at least the 3rd month. The follow-up is for any and all pregnancies in obstetric physiological, pathological, fetal medicine and pregnancy losses. The main objective of psychological support involves preparing the couple to experience motherhood and fatherhood, to develop an active posture of the pregnant and the father, the strengthening of the marital relationship, the facilitation of the maternal-paternal-filial bond and the prevention of postpartum depression and puerperal psychosis.¹

As for the specific objectives, these will be elaborated according to the physical emotional experiences of each patient's pregnancy. It is a personalized interdisciplinary psychological monitoring that aims to provide humanized assistance to women at such a special time and with so many changes.² Pregnancy is contemplated by different feelings and transformations each quarter, which means that the stages of psychological prenatal care are adapted to each moment of the evolution of pregnancy. The use of an initial interview and the investigation through psychological anamnesis are important instruments for the knowledge of the psychic functioning of the pregnant woman and the couple. Anguish and anxiety mark the gestational and postpartum 1st quarter. Stimulating the free expression of these feelings is one of the most important aspects to be highlighted for facilitating the experience of the puerperal pregnancy cycle. The need for information regarding physical and emotional changes, fetal development, childbirth, neonatal care and breastfeeding

are extremely relevant during the approach with the pregnant woman. During psychological counselling, it is important to encourage appropriate body activities according to each case. Meditation, massage and abdominal breathing are also tools taught to the woman and the couple to encourage motherhood, facilitating the moment of delivery and favoring the emotional strengthening of parents with the unborn child.³

Pregnancy is the pinnacle of a female psychosexual development. The changes that may occur are inherent to the puerperal pregnancy cycle, for both men and women; they lead to increased expectations and anxiety, feelings that can be aggravated during this Covid-19 pandemic period.^{4,5} Social distance is mandatory in the midst of the need for health care to the future parents; however, considering the technological advances that have been happening in the last decades, psychological prenatal care remains viable through the use of videoconferencing applications and software such as WhatsApp, Skype and Zoom.⁶

Videoconferences allow the obstetric psychologist to perform the service in all its completeness, and the only difference is in the adaptation of some dynamics – visual resources such as video tutorials or illustrated documents can be used to assist the pregnant woman and the couple in performing physical exercises, massages and meditation, for instance. The current moment sets a great challenge in the hands of professionals, who suddenly see themselves in the need of finding new protocols to reinvent the way of dealing with patients. With the use of virtual resources, obstetric psychology sees itself as an opportunity to experience a new way of guiding pregnant women during the day of setting up their maternity.

Conflict of Interests

The authors have no conflict of interests to declare.

received
June 30, 2020
accepted
August 14, 2020

DOI <https://doi.org/10.1055/s-0040-1718451>.
ISSN 0100-7203.

© 2021. Federação Brasileira das Associações de Ginecologia e Obstetrícia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

References

- 1 Benute GR, Nozzella DC, Prohaska C, Liao A, de Lucia MC, Zugaib M. Twin pregnancies: evaluation of major depression, stress, and social support. *Twin Res Hum Genet.* 2013;16(02):629–633. Doi: 10.1017/thg.2012.153
- 2 Kliemann A, Böing E, Crepaldi MA. [Factors for anxiety and depression during pregnancy: systematic review of empiric articles]. *Mudanças Psicol Saúde.* 2017;25(02):69–76
- 3 Araújo WS, Romero WG, Zandonade E, Amorim MHC. Effects of relaxation on depression levels in women with high-risk pregnancies: a randomised clinical trial. *Rev Lat Am Enfermagem.* 2016;24(00):e2806. Doi: 10.1590/1518-8345.1249.2806
- 4 Silva IL, Lucena Filho EL. [Mental health and social assistance: challenges during COVID - 19]. *Rev Psicol Saberes..* 2020;9(19): 138–146
- 5 Castro P, Narciso C, Matos AP, Werner H, Araujo Júnior E. Pregnant, uninfected, stressed, and confined in the COVID-19 period: what can we expect in the near future? *Rev Assoc Med Bras (1992).* 2020;66(04):386–387. Doi: 10.1590/1806-9282.66.4.386
- 6 Estrela FM, Silva KKA, Cruz MA, Gomes NP. Gestantes no contexto da pandemia da Covid-19: reflexões e desafios. *Physis.* 2020;30(02):e300215. Doi: 10.1590/s0103-73312020300215