The German–Austrian S3 Guideline “Cardiogenic Shock Due to Myocardial Infarction: Diagnosis, Monitoring, and Treatment”

Kevin Pilarczyk¹ Karl Werdan² Martin Russ³ Holger Thiele⁴ Guido Michels⁵ Udo Boeken⁶ Matthias Thielmann⁷

¹Imland Hospital Rendsburg, Department for Intensive Care Medicine, Rendsburg, Schleswig-Holstein, Germany

²Department of Internal Medicine III, Heart Center, Martin-Luther-University Halle-Wittenberg, University Hospital, Germany

³Department of Cardiology, Traunstein Hospital, Traunstein, Germany

⁴University of Leipzig-Heart Center, Department of Internal Medicine/Cardiology, Leipzig, Germany

⁵Department of Acute and Emergency Care, St.-Antonius-Hospital gGmbH, Eschweiler, Germany

⁶Department of Cardiac Surgery, University Hospital, Moorenstrasse 5, Duesseldorf, Germany

⁷Department of Thoracic and Cardiovascular Surgery, West German Heart and Vascular Center Essen, University Hospital Essen, Essen, Germany

Address for correspondence Udo Boeken, MD, Department of Cardiac Surgery, University Hospital, Moorenstrasse 5, Duesseldorf 40225, Germany (e-mail: boeken@uni-duesseldorf.de).

Thorac Cardiovasc Surg 2021;69:684–692.

Abstract

Keywords

- ▶ cardiogenic shock
- ▶ myocardial infarction
- ▶ guideline
- ▶ percutaneous coronary intervention
- ▶ coronary artery bypass graft
- ▶ extracorporeal
- ▶ multiorgan dysfunction syndrome
- ▶ extracorporeal membrane oxygenation
- ▶ intra-aortic balloon counterpulsation
- ▶ ventricular assist device

Despite advances in the treatment of acute myocardial infarction with subsequent mortality reduction, which are mainly caused by the early timing of revascularization, cardiogenic shock still remains the leading cause of death with mortality rates still approaching 40 to 50%. Cardiogenic shock is characterized by a multiorgan dysfunction syndrome, often complicated by a systemic inflammatory response syndrome that affects the outcome more than the reduction of the cardiac contractile function. However, both European and American guidelines on myocardial infarction focus on interventional or surgical aspects only. Therefore, experts from eight German and Austrian specialty societies including the German Society for Thoracic and Cardiovascular Surgery published the German–Austrian S3 guideline “cardiogenic shock due to myocardial infarction: diagnosis, monitoring, and treatment” to provide evidence-based recommendations for the diagnosis and treatment of infarction-related cardiogenic shock in 2010 covering the topics of early revascularization, revascularization techniques, intensive care unit treatment including ventilation, transfusion regimens, adjunctive medical therapy, and mechanical support devices. Within the last 3 years, this guideline was updated as some major recommendations were outdated, or new evidence had been found. This review will therefore outline the management of patients with cardiogenic shock complicating acute myocardial infarction according to the updated guideline with a major focus on evidence-based recommendations which have been found relevant for cardiac surgery.

received

September 20, 2020

accepted after revision

September 25, 2020

published online

December 24, 2020

© 2020. Thieme. All rights reserved.
Georg Thieme Verlag KG,
Rüdigerstraße 14,
70469 Stuttgart, Germany

DOI <https://doi.org/10.1055/s-0040-1719155>.
ISSN 0171-6425.

Introduction

Provided that they reach the hospital, patients with acute myocardial infarction (MI) have more than 90% probability of surviving. However, if cardiogenic shock occurs, either initially or in the course of infarction, only one in two survives.¹ One main cause of the high mortality among patients with inhaled corticosteroids (ICS) is the development of multi-organ dysfunction syndrome (MODS) despite reperfusion of the infarcted vessel as early as possible.^{2,3} Consequently, ICS is not just a disease of the heart, but affects all organs of the patient, who therefore requires intensive care. The current European and American MI guidelines focus their recommendations mostly on “interventional aspects” in the treatment of the coronary artery disease; the “intensive care” aspect of the treatment of MODS, however, is insufficiently considered.^{4,5} This deficit prompted the German and Austrian cardiologists, intensivists, cardiac surgeons, anesthesiologists, and rehabilitation specialists, together with their professional societies and associations, to develop a S3 guideline for “infarction-related cardiogenic shock,” under the auspices of the Association of Scientific Medical Societies in Germany (AWMF, Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften).

The aim of this German–Austrian guideline is to provide an adequate concept of both the cardiological/cardiosurgical and the intensive care aspects of ICS, since the prognosis of patients in this condition depends not only on the impaired cardiac function, but, even more, on the resulting impairment of organ blood supply and microcirculation with consequent MODS. Evidence on diagnosis, monitoring, and therapy of ICS was collected and recommendations compiled in a nominal group process by members of the German and Austrian Societies for Cardiology, Medical and General Intensive Care Medicine, Thoracic Cardiac and Vascular Surgery, Anaesthesiology and Intensive Care Medicine as well as Cardiac Preventive and Rehabilitative Medicine under the auspices of the German Guideline Working Group of Medical Scientific Societies (AWMF). A total of 95 recommendations—including two statements—and seven algorithms were assembled. The most relevant alterations/modifications as well as new recommendations in the updated guideline are given in ►Table 1. The full version and the guideline report are available at www.leitlinien.net (in German).⁶

Medical Societies Involved in Guideline Development

Deutsche Gesellschaft für Kardiologie—Herz und Kreislauforschung (DGK; German Cardiac Society; lead society) (K. Werdan as coordinator, M. Ruß as secretary with M. Kelm, H. Thiele, S. Willems, U. Zeymer, M. Ferrari, H. Figulla, G. Hindricks, P.M. Pieske, and J. Bauersachs).

- Deutsche Gesellschaft für Internistische Intensivmedizin und Notfallmedizin (DGIIN) German Society of Medical Intensive Care and Emergency Medicine (M. Buerke, U. Janssens, G. Michels, and R. Prondzinsky).

Deutsche Gesellschaft für Thorax-, Herz- und Gefäßchirurgie (DGTHG; German Society for Thoracic and Cardiovascular Surgery) (U. Boeken, K. Pilarczyk, and M. Thielmann).

- Österreichische Gesellschaft für Internistische und Allgemeine Intensivmedizin (ÖGIAIM) (Austrian Society of Internal and General Intensive Care Medicine) (A. Geppert).
- Deutsche Interdisziplinäre Vereinigung für Intensivmedizin (DIVI; German Interdisciplinary Association of Intensive Care and Emergency Medicine) (A. Heller).
- Österreichische Kardiologische Gesellschaft (ÖKG; Austrian Society of Cardiology) (G. Delle-Karth, E. Pichler-Cetin).
- Deutsche Gesellschaft für Anästhesie und Intensivmedizin (DGA; German Society of Anaesthesiology and Intensive Care Medicine) (B. Zwißler, J. Briegel).
- Deutsche Gesellschaft für Prävention und Rehabilitation von Herz-Kreislaufkrankungen (DGPR) (A. Schlitt).
- Moderated by Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Gesellschaften (AWMF) (I. Kopp).

Contents of the German–Austrian S3 Guideline “Infarction-Related Cardiogenic Shock: Diagnosis, Monitoring, and Treatment”

- Introduction
- Method
- Synopsis: Diagnosis, monitoring, and treatment of infarction-related cardiogenic shock
- Definition, diagnosis, and monitoring
- Earliest possible coronary revascularization
- Cardiovascular support
- Treatment of complications of infarction-related cardiogenic shock
- Supportive therapy for multiorgan dysfunction syndrome (MODS)
- Nutrition and insulin therapy, red cell substitution and prophylaxis, considerations regarding limitation of treatment
- Aftercare and rehabilitation
- Recommendations “Gemeinsam Klug Entscheiden”
- Need for research

The Aims of the Guideline and Who the Guideline Is for

The aim of the S3 guideline “Infarktbedingter kardiogener Schock: Diagnose, Monitoring, und Therapie” (Infarction-Related Cardiogenic Shock: Diagnosis, Monitoring, and Treatment) is to improve the quality of care of patients with ICS, by publishing evidence-based recommendations. It also presents the “state of the art” in diagnostics, monitoring, and treatment, thus representing a starting point for comparative studies. This is particularly to be emphasized, because many of the recommendations in this S3 guideline are based on expert opinions due to the lack of high-quality evidence.

The recommendations in the S3 guideline are directed at physicians managing patients with shock and acute MI: that

Table 1 Comparison of the original 2010 and the updated version from 2019

Original guideline 2010		Updated guideline 2019	
IABP			
IABP with primary fibrinolysis: In patients with primary fibrinolysis, IABP should be performed adjunctively.	3/4↑	IABP with primary PCI: Routine use of IABPs in patients with cardio- genic shock due to MI is not recommended.	↓↓/1++
IABP with primary PCI: In patients with primary PCI, IABP may be considered, but the evidence is unclear.	3/4↔	IABP with CABG, fibrinolysis, or transfer: No recommendation can be made for patients undergoing revascularization with CABG or fibrinolysis and patients that have to be transferred.	EO
Patient transfer: In patients that have to be transferred to an intervention center, IABP should be used.	3/4↑	Mechanical complications of MI: IABP may be used for hemodynamic stabili- zation in patients with mechanical compli- cations of myocardial infarction including ventricular septal rupture, papillary muscle rupture.	
Temporary mechanical support system			
No recommendation	–	Short-term mechanical circulatory support can be considered in selected patients with MI and cardiogenic shock that cannot be quickly stabilized with conservative management.	⇔/EO
Culprit-lesion-only PCI vs. multivessel PCI			
In selected patients with coronary multivessel disease, complete revascularization during the index primary percutaneous coronary intervention (PCI) apart from the infarct-related artery (IRA) can be performed.	⇔ ↑ 3/4	In patients with coronary multivessel disease and more than one significant stenosis, cul- prit-lesion-only-lesion preferred during the index PCI.	↑↑/1++
Vascular access for PCI			
No recommendation	–	It is possible to use the transradial as well as the transfemoral access in patients with car- diogenic shock. It is recommended to choose the operators standard access in patients without acute coronary syndromes and car- diogenic shock.	⇔/EO
Post-infarction—ventricular septal defect			
Patients with a ventricular septal rupture should undergo urgent surgery after hemodynamic stabilization.	↑, 3/4	Patients with a ventricular septal rupture should undergo urgent surgery or percuta- neous intervention.	↑/EO

Abbreviations: CABG, coronary artery bypass graft surgery; IABP, intra-aortic balloon counterpulsation; MI, myocardial infarction.

is, in particular, cardiologists and specialists in internal medicine, intensivists, heart surgeons, anesthetists, physicians working in interdisciplinary emergency admission services, emergency physicians, and rehabilitation specialists, along with the care staff working with them.

Data Acquisition and Evaluation of Recommendations and Evidence

A systematic search of international guidelines was conducted to generate a statement of the thematic areas and questions on which there was consensus (“source guidelines”; see Leitlinien report at www.leitlinien.net/). In addition, a primary systematic literature search was performed and publications from January 1, 1990 to September 30, 2009

(3,546 results) were included in the first version of this guideline. Publications from October 1, 2009 to January 31, 2019 were also included in this update. Evaluation of study quality and consecutive evidence as well as the assigning of recommendation grades was done using the nominal group process in accordance with the recommendation grades and evidence levels listed in ► **Tables 2 and 3.**

Diagnosis

Early Diagnosis of Cardiogenic Shock—It Is a Clinical Diagnosis

Cardiogenic shock after acute myocardial infarction (CS-AMI) develops within 6 hours in approximately 50% of the patients and within 24 hours in approximately 75% of the

Table 2 Level of evidence (LOE)

Level of evidence		Description
1	++	High-quality meta-analyses, systematic reviews of randomized controlled trials (RCTs), or RCTs with a very low risk of bias.
	+	Well-conducted meta-analyses, systematic reviews, or RCTs with a low risk of bias.
2	++	High-quality systematic reviews of case–control or cohort studies. High-quality case–control or cohort studies with a very low risk of confounding or bias and a high probability that the relationship is causal.
	+	Well-conducted case–control or cohort studies with a low risk of confounding or bias and a moderate probability that the relationship is causal.
3		Analytic studies without a concurrent comparison group, e.g., before-and-after studies, interrupted time series nonanalytic studies, e.g., case reports, case series.
4		Expert opinion (EO), e.g., editorial commentaries, guidelines without a clear methodology.

Table 3 Grade of recommendation (GoR)

GoR	Description
↑↑	Strongly recommended: “shall” (usually based on studies with evidence level 1 ++ or 1 +)
↑	Recommended: “should” (usually based on studies with evidence level 2 ++ or 2 +)
↔	No recommendation: “may” (no confirmed study results exist that demonstrate either a beneficial or a harmful effect).
↓	Rejected: “should not” (negative recommendation).
↓↓	Strongly rejected: “shall not” (strong negative recommendation).

patients. Consequently, shock can already occur in the pre-hospital phase, and the emergency physician has to recognize cardiogenic shock as soon as possible mainly as a clinical diagnosis. Cornerstones of the diagnosis are the 12-lead electrocardiogram (STEMI, seldom Non-ST elevation myocardial infarction [NSTEMI]), hypotension (systolic blood pressure <90 mm of mercury [mm Hg] for at least 30 minutes [min] in the absence of volume depletion), and clinical signs of reduced organ perfusion (cold extremities, oliguria, altered mental status [e.g., agitation]). In those, one-fourth of CS-AMI patients do not present with hypotension and diagnosis of shock must exclusively rely on those clinical signs.

The diagnosis of cardiogenic shock can be made on the basis of clinical and hemodynamic parameters^{7,8}:

Clinical Signs of Cardiogenic Shock

1. Systemic hypoperfusion
 - (a) Oligo-/anuria <30 mL/h.
 - (b) Cyanotic extremities.
 - (c) Signs of cerebral hypoperfusion with somnolence and confusion.

Hemodynamic Signs/Interventions of Cardiogenic Shock

1. Blood pressure <90 mm Hg for more than 90 minutes.
2. Systemic hypotension.
3. Administration of catecholamines to stabilize the patient.
4. Use of intra-aortic counterpulsation.
5. Cardiac index <2.2 L/min/m².
6. Pulmonary capillary wedge pressure of >15 mm Hg.

Revascularization

Coronary reperfusion is the mainstay evidence-based therapeutic intervention for patients with acute MI presenting with CS.⁹ In this section, recommendations for reperfusion and revascularization techniques and other adjunctive therapies used in the management of CS are reviewed.

Timing

Early coronary revascularization is an essential therapeutic intervention for patients with ACS complicated by CS. However, the SHOCK (Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock) trial and (S)MASH (Swiss) Multicenter Trial of Angioplasty for Shock, are the only randomized trials that examined the benefit of early revascularization in the setting of CS.^{9–11} Regrettably, (S) MASH was terminated prematurely due to insufficient patient enrollment.¹¹

The SHOCK trial demonstrated that, in patients with cardiogenic shock complicating AMI, emergency revascularization with PCI or coronary artery bypass graft (CABG) surgery improved long-term survival, compared with initial intensive medical therapy.^{9,10} All-cause mortality at 6 months was lower in the group assigned to revascularization than in the medically treated patients (50.3 vs. 63.1%, respectively; relative risk 0.80, 95% confidence interval 0.65–0.98, $p = 0.03$). A recent analysis of 12,675 STEMI patients in the FITT-STEMI (Feedback Intervention and Treatment Times in ST-Elevation Myocardial Infarction) trial emphasizes the strong impact of time delays on mortality, particularly in STEMI patients with cardiogenic shock or out-of-hospital cardiac arrest.^{12,13} In shock without out-of-hospital cardiac

arrest, every 10-minute treatment delay between 60 to 180 minutes from the first medical contact resulted in 3.3 additional deaths per 100 PCI-treated patients and in 1.3 additional deaths after out-of-hospital cardiac arrest without cardiogenic shock.

Type of Revascularization

Coronary stenting is the technique of choice during primary PCI.¹⁴ Compared with balloon angioplasty alone, stenting with a bare-metal stent (BMS) is associated with a lower risk of reinfarction and target vessel revascularization. However, it is not associated with a reduction in the mortality rate. In primary PCI, drug-eluting stents reduce the risk of repeated target vessel revascularization compared with BMS. A sub-analysis of the SHOCK trial comparing patients treated with CABG or PCI showed similar survival rates between the two subgroups. There were more patients with diabetes (48.9 vs. 26.9%; *p* = 0.02), three-vessel disease (80.4 vs. 60.3%; *p* = 0.03), and LM coronary disease (41.3 vs. 13.0%; *p* = 0.001) in the CABG group. The results of this nonrandomized comparison suggest that CABG should be considered in patients with cardiogenic shock who have suitable anatomy, particularly if successful PCI is not feasible.

Recommendation 5.1.2.A. (Revascularization/PCI) Emergency PCI of the culprit lesion is indicated for patients with cardiogenic shock due to STEMI or NSTEMI-ACS, independent of time delay of symptom onset, if coronary anatomy is amenable to PCI.	↑↑/1+ ^a
Recommendation 5.1.2.B. (PCI) In patients presenting with ICS, prompt intervention with primary PCI within 90 minutes of first medical contact should be performed. ^a DGIIN: 90 minutes only for patients with cardiogenic shock due to a STEMI.	↑/EC

PCI Strategy

Culprit Lesion or Total Strategy

More than three-fourths of the patients in CS present with multivessel disease. Complete revascularization, addressing both culprit and hemodynamically significant non-culprit lesions, has historically been the preferred strategy in patients with acute MI and CS and was recommended in recent guidelines. However, this paradigm has recently been challenged. The CULPRIT-SHOCK (Culprit Lesion Only PCI vs. Multivessel PCI in Cardiogenic Shock) trial randomized 706 patients with STEMI/NSTEMI and an identifiable culprit lesion to multivessel or culprit lesion-only PCI.¹⁵ The composite primary end point was death or renal failure requiring dialysis at 30 days. The trial demonstrated a 9.5% absolute risk reduction of the composite primary end point in the culprit lesion-only group (7.3% of which was attributable to an absolute risk reduction in all-

cause mortality). Of note, the culprit-lesion only cohort had the option for staged revascularization of nonculprit lesions and almost 20% of patients underwent further staged or urgent PCI. Additionally, 75 patients crossed over from culprit lesion-only to multivessel PCI raising the possibility of including more complex and comorbid patients in the multivessel PCI group, thus overestimating the benefit of culprit lesion-only PCI. Also, greater dye loads in multivessel PCI may partially account for observed differences observed. Another limitation of the study was that low rates of MCS device were used in the multivessel PCI group. One-year follow-up showed no mortality difference between the culprit lesion-only and multivessel PCI groups (50 vs. 56.9%, respectively). The CULPRIT-SHOCK trial contradicts widespread current practice and prior studies in nonshock patients (DANAMI-3-PRIMULTI, PRAMI, CvLPRIT) which suggested that there may be a benefit from complete revascularization.^{16–18}

Recommendation 5.2.4.A (Culprit lesion PCI) In patients with coronary multivessel disease and more than one significant stenosis, culprit-lesion-only-lesion during the index PCI is preferred.	↑↑/1+ ^a
---	--------------------

Fibrinolysis

Fibrinolytic therapy is an important reperfusion strategy in settings where primary PCI cannot be offered in a timely manner and prevents 30 early deaths per 1,000 patients treated within 6 hours after symptom onset. The largest absolute benefit is seen among patients at highest risk, including the elderly, and when treatment is offered <2 hours after symptom onset.¹⁹ Fibrinolytic therapy is recommended if primary PCI cannot be performed within 6 hours from MI diagnosis and there are no contraindications. If the patient is presented later (particularly after 3 hours), more consideration should be given to transfer for primary PCI (as opposed to administering fibrinolytic therapy) because the efficiency and clinical benefit of fibrinolysis decrease as the time from symptom onset increases. In the presence of contraindications for fibrinolytic treatment, it is important to weigh the potentially life-saving effect of fibrinolysis against potentially life-threatening side effects, taking into account alternative treatment options such as delayed primary PCI.

Recommendation 5.4.1.A. (Systemic fibrinolysis) Fibrinolysis should be performed in patients with ICS if early coronary angiography and revascularization cannot be performed within 6 hours. Coronary angiography should be performed as soon as possible.	↑/EO
---	------

Surgical Revascularization (Coronary Artery Bypass Graft Surgery)

Emergent CABG surgery should be considered for patients with cardiogenic shock with a patent IRA but with unsuitable anatomy for PCI. In patients with MI-related mechanical complications who require coronary revascularization, CABG is recommended at the time of repair. In MI patients with failed PCI or coronary occlusion not amenable to PCI, emergent CABG is infrequently performed because the benefits of surgical revascularization in this setting are uncertain. As the delay to reperfusion is long, the probabilities of myocardial salvage affecting prognosis are low and the surgical risks are elevated.

In a propensity-matched comparison from the AUCITY trial, moderate- and high-risk patients with ACS and multi-vessel disease treated with PCI rather than CABG had lower rates of periprocedural stroke, MI, major bleeding, and renal injury, with comparable 1-month and 1-year rates of mortality, but more frequently developed recurrent ischemia requiring repeated revascularization procedures during follow-up.^{20,21} However, recently published studies indicate a reasonable survival rate in patients with acute MI complicated by cardiogenic shock. Therefore, early CABG should be performed in patients with a coronary anatomy not suitable for PCI and should be discussed as alternative to PCI in patients with complex coronary anatomy after interdisciplinary evaluation of the Heart Team.

Recommendation 5.2.5.A. (Complex coronary anatomy or failed percutaneous PCI) In patients with complex coronary anatomy emergency surgical or interventional revascularization should be performed as decided by the interdisciplinary Heart Team. Emergency CABG is recommended for patients with cardiogenic shock if the coronary anatomy is not amenable to PCI.	↑↑/1+
Recommendation 5.3.3.B (Complex coronary anatomy or pathology and mechanical complication of myocardial infarction) In patients with complex coronary anatomy and mechanical complications of MI, emergency surgical or interventional therapy is indicated, as decided by the interdisciplinary Heart Team.	↑↑/EO

Mechanical Circulatory Support

Escalating doses of vasopressors and inotropes are associated with significant limitations including arrhythmias, increased myocardial oxygen consumption, and inadequate circulatory support as well as increased mortality. Therefore, mechanical circulatory support is an essential part of the management of cardiogenic shock and is commonly utilized as a bridge-to-decision, whether it is recovery, palliation, heart transplantation, or a durable mechanical circulatory support

device. Options for acute percutaneous MCS include the intra-aortic balloon pump (IABP), axial flow pumps (Impella LP 2.5, Impella CP), left atrial-to-femoral arterial ventricular assist devices (Tandem Heart), and venous-arterial extracorporeal life support (ECLS).

IABP

Since its introduction into clinical practice >50 years ago, intra-aortic balloon counterpulsation has been used empirically to provide hemodynamic support in patients undergoing coronary revascularization in the setting of MI and cardiogenic shock. In the landmark SHOCK trial, conducted between 1993 and 1998, IABPs were implanted in 86% of the participants, irrespective of the assigned management strategy. However, despite its widespread use and high recommendations in international guidelines, the last version of the German–Austrian S3 guideline gave only a weak recommendation (↑) for the use of IABP in ICS patients treated with systemic fibrinolysis, and only “may” information for patients treated with PCI (↔) as there were no randomized controlled trials showing a benefit of aortic counterpulsation.

The IABP-SHOCK II (Intra-aortic Balloon Pump in Cardiogenic Shock II) trial randomly assigned 600 participants planned for early revascularization of acute MI complicated by cardiogenic shock to either IABP placement or no IABP placement.²² The primary end point was 30-day all-cause mortality. After 30 days or at day 30, all-cause mortality was 40%, with no difference between patients randomized to receive an IABP versus those who were not. There were no differences between treatment groups in secondary outcomes, including bleeding, ischemic complications, stroke, time to hemodynamic stabilization, intensive care unit length of stay, and the dose and duration of catecholamine therapy. At 6 years of follow-up, all-cause mortality was high and did not differ between the IABP and control groups (66.3 vs. 67.0%) in intention-to-treat, per-protocol, and as-treated analyses. No signal for benefit associated with IABP use was observed in any prespecified or post hoc subgroups. There were no differences in the frequency of recurrent MI, repeat revascularization, stroke, or cardiovascular rehospitalization between the two groups. Quality of life, measured by the EuroQol 5D questionnaire and New York Heart Association classification, was favorable in survivors of cardiogenic shock. Four of five survivors had New York Heart Association Class I or II symptoms, with no difference between patients randomly assigned to IABP and no IABP therapy.²³ Based on the current evidence, IABP placement cannot be recommended routinely in patients with ICS.

Recommendation 7.3.7.A. (IABP in patients with PCI) Routine use of IABPs in patients with cardiogenic shock due to ACS is not recommended.	↓/1++
Recommendation 7.3.8.A. (IABP in patients with mechanical	↔/EO

(Continued)

(Continued)

complications of MI) IABP may be used for hemodynamic stabilization in patients with mechanical complications of myocardial infarction including ventricular septal rupture and papillary muscle rupture.	
Recommendation 7.3.9.A. (IABP in patients with CABG, fibrinolysis, transfer) No recommendation can be made for patients undergoing revascularization with CABG or fibrinolysis and patients that have to be transferred.	Statement

Other Mechanical Circulatory Support Systems

Alternative percutaneous left ventricular assist devices (LVADs)—centrifugal pumps without oxygenator (Tandem Heart) and with oxygenator (venoarterial extracorporeal membrane oxygenation, LifeBridge) as well as axial flow pumps (Impella Recover LP2.5 and Impella Recover LP5.0) are hemodynamically more effective than the IABP.¹⁹

A recent meta-analysis on MCS in cardiogenic shock included four randomized trials investigating the efficacy and safety of percutaneous LVADs versus IABP, and demonstrated similar short-term mortality despite initial beneficial effects on arterial blood pressure and peripheral perfusion, measured by serum lactate levels.²⁰ In all trials, a higher rate of bleeding from vascular access sites and a significantly higher incidence of limb ischemia following percutaneous LVAD was notable. In summary, the evidence for mechanical circulatory systems is insufficient to provide a recommendation on its routine clinical use in cardiogenic shock.

A recently published meta-analysis comparing mortality in patients treated with and without ECLS support included 13 studies in which nine studies dealt with cardiac arrest patients ($n = 3098$) and four studies with patients suffering from cardiogenic shock after acute MI ($n = 235$).²⁴

In cardiac arrest, the use of ECLS was associated with an absolute increase of 30 days survival of 13% compared with patients in which ECLS was not used and a higher rate of favorable neurological outcome was achieved at 30 days. In cardiogenic shock, ECLS showed a 33% higher 30-day survival compared with IABP but no difference when compared with TandemHeart/Impella.

Therefore, the current guideline suggests that short-term mechanical circulatory support can be considered in selected patients with ACS and cardiogenic shock that cannot be rapidly stabilized with conservative management

if

- MCS is implanted before occurrence of irreversible end-organ-damage.

- MCS implantation does not delay emergency revascularization.
- There is a clear patient-specific therapeutic goal as evaluated and documented by the Heart Team.
- There is a structured cooperation with a heart-failure center for early destination therapy.
- Patients are enrolled in a registry for MCS of the participating medical societies.

The choice of which MCS device to be used is based on many factors, including type of shock (right heart failure, left heart failure) as well as operator abilities and institutional resources.

Mechanical Complications of MI

Ventricular Septal Rupture

Ventricular septal rupture may occur within 24 hours to several days after MI and usually presents as rapid-onset clinical deterioration with acute heart failure or cardiogenic shock. The diagnosis is confirmed by echocardiography. IABP may stabilize patients in preparation for angiography and surgery.

Surgical repair may be required urgently, but there is no consensus on the optimal timing for surgery. Early surgery is associated with a high mortality rate, reported as 20 to 40%, and a high risk of recurrent ventricular rupture, while delayed surgery allows easier septal repair in scarring tissue but carries the risk of rupture extension and death while waiting for surgery.²⁵ Therefore, early surgery should be performed in all patients with severe heart failure that does not respond rapidly to aggressive therapy, but delayed elective surgical repair may be considered in patients who respond well to aggressive heart failure therapy. Percutaneous closure of the defect with appropriately designed devices may soon become an alternative to surgery.²⁶

Recommendation 8.4.2.A. (Ventricular septal rupture) Patients with a ventricular septal rupture should undergo urgent surgery or percutaneous intervention.	↑/EO
---	------

Papillary Muscle Rupture

Acute mitral regurgitation may occur 2 to 7 days after AMI due to rupture of the papillary muscle or chordae tendineae.²⁷ The rupture may be complete or involve one or more of the heads and is 6 to 12 times more frequent in the posteromedial papillary muscle because of its single artery blood supply. Papillary muscle rupture usually presents as sudden hemodynamic deterioration with acute dyspnea, pulmonary edema, and/or cardiogenic shock. Immediate treatment is based on afterload reduction to reduce regurgitant volume and pulmonary congestion. Intravenous diuretic and vasodilator/inotropic support, as well as IABP, may stabilize patients in preparation for angiography and

surgery. Emergency surgery is the treatment of choice although it carries a high operative mortality (20–25%).

Recommendation 8.4.4.A.

(Acute severe mitral regurgitation)
Patients with a significant acute mitral regurgitation should undergo urgent surgery.

↑↑/EO

Author's Contribution

All authors participated sufficiently in the work to take public responsibility for the content. K.W. and M.R. designed the research/study and analyzed data. K.P., K.W., M.R., H.T., G.M., U.B., and M.T. performed study research. K.P., U.B., M.T. wrote the paper. U.B., H.T., and K.W. did the supervision and all the authors did the critical revision of the manuscript.

Conflict of Interest

None.

References

- Nguyen HL, Yarzelski J, Lessard D, Gore JM, McManus DD, Goldberg RJ. Ten-year (2001–2011) trends in the incidence rates and short-term outcomes of early versus late onset cardiogenic shock after hospitalization for acute myocardial infarction. *J Am Heart Assoc* 2017;6(06):e005566
- Werdan K, Ruß M, Engemann L, et al. Deutsch-österreichische S3-Leitlinie "Infarktbedingter kardiogener Schock—diagnose, monitoring und therapie". *Kardiologie* 2011;5(03):166–224
- Prondzinsky R, Lemm H, Swyter M, et al. Intra-aortic balloon counterpulsation in patients with acute myocardial infarction complicated by cardiogenic shock: the prospective, randomized IABP SHOCK Trial for attenuation of multiorgan dysfunction syndrome. *Crit Care Med* 2010;38(01):152–160
- Ibanez B, James S, Agewall S, et al; ESC Scientific Document Group. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: the Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2018;39(02):119–177
- Neumann FJ, Sousa-Uva M, Ahlsson A, et al; ESC Scientific Document Group. 2018 ESC/EACTS guidelines on myocardial revascularization. *Eur Heart J* 2019;40(02):87–165
- AWMF-RN 019/013. Deutsch-österreichische S3-Leitlinie "Infarktbedingter kardiogener Schock—Diagnose, Monitoring und Therapie"; 2020 Available at: <https://leitlinien.dgk.org/2020/deutsch-oesterreichische-s3-leitlinie-infarktbedingter-kardiogener-schock-diagnose-monitoring-und-therapie/>. Accessed June 03, 2020
- Standl T, Annecke T, Cascorbi I, Heller AR, Sabashnikov A, Teske W. The nomenclature, definition and distinction of types of shock. *Dtsch Arztebl Int* 2018;115(45):757–768
- Vincent J-L, De Backer D. Circulatory shock. *N Engl J Med* 2013;369(18):1726–1734
- Hochman JS, Sleeper LA, Webb JG, et al. for the SHOCK Investigators. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. SHOCK investigators. Should we emergently revascularize occluded coronaries for cardiogenic shock. *N Engl J Med* 1999;341(09):625–634
- Hochman JS, Sleeper LA, White HD, et al; SHOCK Investigators. Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock. One-year survival following early revascularization for cardiogenic shock. *JAMA* 2001;285(02):190–192
- Urban P, Stauffer J-C, Bleed D, et al. A randomized evaluation of early revascularization to treat shock complicating acute myocardial infarction. The (Swiss) Multicenter Trial of Angioplasty for Shock-(S)MASH *Eur Heart J* 1999;20(14):1030–1038
- Scholz KH, Maier SKG, Maier LS, et al. Impact of treatment delay on mortality in ST-segment elevation myocardial infarction (STEMI) patients presenting with and without haemodynamic instability: results from the German prospective, multicentre FITT-STEMI trial. *Eur Heart J* 2018;39(13):1065–1074
- Wijns W, Naber CK. Reperfusion delay in patients with high-risk ST-segment elevation myocardial infarction: every minute counts, much more than suspected. *Eur Heart J* 2018;39(13):1075–1077
- Roffi M, Patrono C, Collet JP, et al; ESC Scientific Document Group. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: task force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2016;37(03):267–315
- Thiele H, Akin I, Sandri M, et al; CULPRIT-SHOCK Investigators. One-year outcomes after PCI strategies in cardiogenic shock. *N Engl J Med* 2018;379(18):1699–1710
- Engström T, Kelbæk H, Helqvist S, et al; DANAMI-3—PRIMULTI Investigators. Complete revascularisation versus treatment of the culprit lesion only in patients with ST-segment elevation myocardial infarction and multivessel disease (DANAMI-3—PRIMULTI): an open-label, randomised controlled trial. *Lancet* 2015;386(9994):665–671
- Wald DS, Morris JK, Wald NJ, et al; PRAMI Investigators. Randomized trial of preventive angioplasty in myocardial infarction. *N Engl J Med* 2013;369(12):1115–1123
- Gershlick AH, Banning AS, Parker E, et al. Long-term follow-up of complete versus lesion-only revascularization in STEMI and multivessel disease: the CvLPRIT trial. *J Am Coll Cardiol* 2019;74(25):3083–3094
- Boersma E, Maas AC, Deckers JW, Simoons ML. Early thrombolytic treatment in acute myocardial infarction: reappraisal of the golden hour. *Lancet* 1996;348(9030):771–775
- Ben-Gal Y, Moses JW, Mehran R, et al. Surgical versus percutaneous revascularization for multivessel disease in patients with acute coronary syndromes: analysis from the ACUTY (acute catheterization and urgent intervention triage strategy) trial. *JACC Cardiovasc Interv* 2010;3(10):1059–1067
- Ben-Gal Y, Mohr R, Feit F, et al. Surgical versus percutaneous coronary revascularization for multivessel disease in diabetic patients with non-ST-segment-elevation acute coronary syndrome: analysis from the Acute Catheterization and Early Intervention Triage Strategy trial. *Circ Cardiovasc Interv* 2015;8(06):e002032
- Thiele H, Zeymer U, Neumann F-J, et al; IABP-SHOCK II Trial Investigators. Intraaortic balloon support for myocardial infarction with cardiogenic shock. *N Engl J Med* 2012;367(14):1287–1296
- Thiele H, Zeymer U, Thelemann N, et al. IABPSHOCK II Trial (Intraaortic Balloon Pump in Cardiogenic Shock II) Investigators. Intraaortic balloon pump in cardiogenic shock complicating acute myocardial infarction: long-term 6-year outcome of the randomized IABP-SHOCK II trial. *Circulation* 2018 (e-pub ahead of print). Doi: 10.1161/CIRCULATIONAHA.118.038201
- Ouweneel DM, Schotborgh JV, Limpens J, et al. Extracorporeal life support during cardiac arrest and cardiogenic shock: a systematic review and meta-analysis. *Intensive Care Med* 2016;42(12):1922–1934
- Crenshaw BS, Granger CB, Birnbaum Y, et al; GUSTO-I (Global Utilization of Streptokinase and TPA for Occluded Coronary

- Arteries) Trial Investigators. Risk factors, angiographic patterns, and outcomes in patients with ventricular septal defect complicating acute myocardial infarction. *Circulation* 2000;101(01):27–32
- 26 Holzer R, Balzer D, Amin Z, et al. Transcatheter closure of post-infarction ventricular septal defects using the new Amplatzer muscular VSD occluder: results of a U.S. Registry. *Catheter Cardiovasc Interv* 2004;61(02):196–201
- 27 Bajaj A, Sethi A, Rathor P, Suppogu N, Sethi A. Acute complications of myocardial infarction in the current era: diagnosis and management. *J Investig Med* 2015;63(07):844–855