Return to Work after Primary Total Knee Arthroplasty under the Age of 65 Years: A Systematic Review

Dries Van Leemput, MD¹ Jef Neiryndck, MD¹ Pieter Berger, MD¹ Hilde Vandenneucker, MD, PhD²

¹ Division of Orthopaedics, University Hospitals Leuven, B-3000 Leuven, Belgium

² Division of Orthopaedics, Department of Development and Regeneration—Organ Systems Cluster, KU Leuven, University Hospitals Leuven, Belgium, B-3000 Leuven, Belgium

Address for correspondence Dries Van Leemput, MD, Division of Orthopaedics, University Hospitals Leuven, B-3000 Leuven, Belgium (e-mail: driesvanleemput@hotmail.com).

J Knee Surg 2022;35:1249–1259.

Abstract

A growing number of patients undergoing total knee arthroplasty (TKA) is at working age and need to return to work (RTW) after surgery. The aim of this systematic review is to give an overview of the literature regarding RTW after TKA and beneficial and limiting factors influencing this process. A systematic search in four electronic databases was conducted in November 2019 to identify studies describing RTW after primary TKA in patients aged 65 years or younger. Study characteristics and data on work status before and after surgery were extracted. All studies were assessed for risk of bias. Fourteen studies published between 2009 and 2019 were included in this review, accounting for a total of 3,073 patients. The percentage of patients working after TKA ranged from 36 to 89%, and the fraction of patients working before and returning to work after surgery ranged from 40 to 98%. Mean time of RTW ranged from 7.7 to 16.6 weeks. Most important factors associated with a slower or no RTW were a more physical nature of employment and preoperative absence from work. The majority of patients undergoing TKA returned to work postoperatively. However, comparison between studies is seriously hampered by the wide variation regarding the definition and timeframe used to measure the work status. Therefore, standardized outcome measures for studies investigating RTW after TKA are warranted. We identify this review as level-I evidence (systematic review of level-I and level-II studies).

Keywords

- work
- employment
- sick leave
- arthroplasty
- replacement
- knee

Osteoarthritis (OA) is one of the most common diseases of the knee, causing pain and often leading to disability and sick leave. Painful, severely disabling OA of the knee affects approximately 1.5% of adults over the age of 55 years.¹

Total knee arthroplasty (TKA) is an effective treatment for patients with OA, reducing pain and improving function and quality of life.² This intervention has shown to have a good satisfaction rate, especially in older patients.³ However, in recent years, a growing portion of patients is undergoing TKA at working age and this is expected to rise even further in the

next years.^{4,5} Also, the average retirement age has increased to an average of 65 years in Europe and is expected to rise to 67 years in the next 10 years.⁶ Therefore, an increasing number of knee OA patients will need to be able to return to work (RTW) after TKA. A previous systematic review showed that the proportions of RTW patients are ranging from 71 to 83% at 3 to 6 months after TKA.⁷

It is important to have insight in the rate of RTW for both individual and socioeconomic interests. Knowledge of determinants associated with a delayed return or no RTW could help

received

June 4, 2020

accepted after revision

November 29, 2020

published online

January 20, 2021

© 2021. Thieme. All rights reserved.
Thieme Medical Publishers, Inc.,
333 Seventh Avenue, 18th Floor,
New York, NY 10001, USA

DOI <https://doi.org/10.1055/s-0040-1722626>.
ISSN 1538-8506.

to focus on the right subpopulation of patients when applying appropriate management and preventive measures.

Systematic literature on this subject is scarce and outdated. The most recent systematic review, publicized in 2014, concluded that the literature on work status after total hip arthroplasty (THA) is more extensive than for TKA, as they only had four studies providing data on RTW after TKA.⁷ In addition, RTW after TKA seems to be a field of growing interest, with several studies being published in recent years. Therefore, a systematic summarization of the recent literature would be very useful.

The aim of this study is to provide an overview of the available literature on work status and RTW in patients under the age of 65 years undergoing primary TKA, and to determine beneficial and limiting factors associated with RTW postoperatively.

Methods

The selection process was conducted by two different authors independently (D.V.L. and J.N.). Any discrepancies in study selection were discussed until consensus was reached.

Database Search

A systematic database search was conducted in PubMed, Embase, Cochrane library, and Web of Science on November 18, 2019. A search strategy was developed for each database according to their specific search methods. The search strategy for PubMed consisted of the Medical Subject Headings (MeSH) terms "Return to work," "Sick Leave," and "Arthroplasty, Replacement, Knee." These terms were combined with synonyms in free words to obtain articles not covered by MeSH terms. The search strategy for each database can be found in the Supplementary data (available online only).

Study Selection

After removing the duplicates and the articles publicized before January 1, 2020, all titles and abstracts were screened. Articles were included for further selection if they met the following inclusion criteria: (1) patients underwent primary TKA, (2) articles reported on RTW after surgery, and (3) patients aged 65 years or below.

Next, using these same criteria, full-text articles were screened. Articles including revision TKA or hemiarthroplasty were excluded, unless they reported data for primary TKA separately. We included articles regardless of the indication for primary TKA. Articles investigating RTW after both TKA and THA were included if data were available for TKA separately.

Data Extraction

Study characteristics extracted were first author, year of publication, country, study type (retrospective or prospective), data collection method, indication for TKA, population size undergoing primary TKA, mean age, male/female ratio, and follow-up time.

Data extracted on work status were work status before surgery, description of work status after surgery, fraction of patients returning to work after surgery, time after surgery

that RTW was measured, mean time taken to RTW, and influencing factors associated with RTW.

Methodological Quality

We assessed the risk for potential bias in included articles using the Quality in Prognosis Studies (QUIPS) checklist. This is a quality assessment tool developed to assess risk of bias in prognostic factor studies. It is based on a review discussing quality of prognostic studies in systematic reviews.⁸ This checklist consists of six domains being "study participation," "study attrition," "prognostic factor measurement," "outcome measurement," "study confounding," and "statistical analysis and reporting." Risk of bias was rated as high, moderate, or low depending on how many criteria in each domain were met. Two authors independently assessed the risk of bias in each study using this tool.

Results

Search Results

The initial search strategy provided 378 articles of which 292 remained after removing the duplicates. Thirteen articles published before the year 2000 were removed. After selection based on title and abstract, 53 articles remained for further analysis. Full-text screening of the selected articles resulted in 14 studies that met our eligibility criteria which were included in this review. A flow diagram of the selection process is shown in ► Fig. 1.

Study Characteristics

Study characteristics of the included articles are presented in ► Table 1. Articles were published between 2009 and 2019. Four studies had a prospective observational design and 10 studies had a retrospective design. Five studies were conducted in the United Kingdom, three in the Netherlands, two in the United States, one in Sweden, one in Canada, one in Finland, and one in Australia. The population sizes of the studies varied from six patients to 996 patients, and the articles together accounted for a total study population of 3,073 patients. The mean age ranged from 46 to 59 years. The indications for TKA were primarily OA. Two articles were qualitative studies using convenience sampling and with total population sizes of 6 and 10 patients, respectively.^{9,10} Therefore, we did not include RTW rates extracted from these two studies, as results would not be representative for the population. We did, however, include the patients' experiences and influencing factors reported in these studies.

Return to Work

► Table 2 shows the pre- and postoperative work status and the mean time taken to RTW. Work status prior to TKA was described in all studies, but the definition that was used differed among studies. Some articles reported the number of patients on sick leave in the months before surgery, others reported the number of patients in employment. Overall, the portion of patients working preoperatively ranged from 50 to 100%. Four studies included only patients who were working before surgery.^{10–13}

Fig. 1 Flowchart of the selection process. WoS, Web of Science.

All articles provided information on RTW after surgery. However, there was great variability concerning both the definition of work status postoperatively, as well as time after surgery when RTW was measured. The portion of patients working after TKA ranged from 36 to 95% (mean, 82%), and the fraction of patients working before and returning to work after surgery ranged from 40 to 98% (mean, 89%). Mean time taken to RTW ranged from 7.7 to 16.6 weeks (mean, 13 weeks).

Three articles specified whether patients returned to the same job or to a different job. In these three articles, 85, 91, and 100% of the RTW patients were returned to the same job.^{14–16} One article reported RTW rates for men and women separately.¹⁷ In this study, more men returned to work than women (men, 88%; women, 83%). No consistent differences in RTW rate were found between studies performed in different countries.

Associated Factors

An overview of the associated factors is shown in ► **Table 3**. All studies mentioned factors that could be influencing RTW. Ten studies performed statistical analysis to investigate relations between RTW and variables. Seven of them did find one or more correlations to be statistically significant, and three studies performed statistical analysis but did not find any significant correlations. Six studies found significant correlation while applying multivariate analysis.^{11–14,18,19}

Four studies did not perform any statistics but mentioned preferences or influencing factors reported by patients to be influencing their RTW.^{9,10,16,17}

Factors shown to be correlated with RTW by multivariate analysis were the physical nature of employment ($p = 0.03$,¹⁴ 0.012 ,¹¹ and <0.0001 ^{12,18}), preoperative absence from work ($p < 0.0001$ ¹⁸ and 0.001 ¹⁹), flexible work conditions

Table 1 Study characteristics

Year	First author	Country	Study type	Data collection method	Indication	Population size	Mean age (range)	Male/female	Follow-up	Mean follow-up	Ref.
2019	A.R. Boersma	The Netherlands	Prospective	Questionnaire	OA	146	56	43/57	12 months	–	11
2019	P. Lankinen	Finland	Retrospective	Surveys, employers' records, and national health registers	–	452	57	20/80	/	4.3 years	18
2019	L. McConagle	Australia	Retrospective	Questionnaire	–	31	56	64/52	6–12 months	–	12
2017	K. Stigmar	Sweden	Retrospective	Health care register: Swedish Social Insurance Agency data	OA	996	55 (40–59)	44/56	1 year before–2 year after surgery	2 years	17
2017	P. Maillette	Canada	Qualitative retrospective	Semistructured interviews	OA (83%), trauma (17%)	6	58 (52–62)	50/50	6–12 months	–	9
2017	C.E.H. Scott	The United Kingdom	Prospective observational	Questionnaire	OA (95.5%), RA (4.5%)	289	59 (42–65)	49/51	2–3.9 years	3.4 years	14
2016	M. Bardgett	The United Kingdom	Qualitative retrospective	Semistructured interviews	OA	10	54 (40–59)	50/50	6 months–3 years	–	10
2016	C.S. Leichtenberg	The Netherlands	Prospective observational	Questionnaire	OA	56	56 (18–65)	45/55	1 year	1 year	19
2015	P.J. Belmont	The United States	Retrospective	Health care register: Military Health System Management Analysis and Reporting Tool database	OA (99%), RA (1%)	159	46 (24–61)	76/24	Minimum 2 years	4.1 years	13
2014	C. Tilbury	The Netherlands	Prospective observational	Questionnaire	OA	120	57 (18–65)	34/66	1 year	1 year	7
2015	B.D. Kleim	The United Kingdom	Retrospective	Questionnaire	OA	50	54 (20–59)	34/66	6 months–3 years	21 months	21
2014	A.V. Lombardi	The United States	Retrospective multicenter	Telephone questionnaire	OA (98%), PTA (2%), avasc necr (0.003%)	661	54 (19–60)	39/61	Minimum 1 year	–	15
2010	J.A.J. Foote	The United Kingdom	Retrospective	Questionnaire	OA (88%), Trauma (12%), other (10%)	41	53 (40–60)	38/71	14–61 months	36 months	20
2009	H. Lyall	The United Kingdom	Retrospective	Postal questionnaire	OA (70%), RA (17%), trauma (11%)	56	58 (48–60)	59/41	47–112 months	64 months	16

Abbreviations: OA, osteoarthritis; RA, rheumatoid arthritis; Ref. reference.

Table 2 Return to work after TKA

Year	First author	Population size	Preoperative work status	Working before TKA (%)	Definition of postoperative work-status	Postoperative work-status	Working after TKA (%)	Working before TKA and RTW (%)	Time post-op that RTW was measured	Time to RTW	Ref.
2019	A.R. Boersma	146	Patients were eligible if they had a paid job	100	First time participants returned to work partially or fully after surgery	<ul style="list-style-type: none"> • 6 weeks: 18% partial, 6% full time • 3 months: 36% partial, 26% full time • 6 months: 26% partial, 59 full time • 12 months: 11% partial, 84% full time 	95	95	6 weeks, 3, 6, and 12 months	Mean: 93 days (13.3 weeks)	11
2019	P. Lankinen	452	Preoperative sickness absence: no: 71.2%, yes: 28.8%	71	The number of days between the date of discharge and the date of the end of the sick leave within one year	After the surgery, 87% (n = 394) of patients returned to work on average after 116 (SD = 53, range: 28–356) days of sickness absence	87	–	1 year	Mean 116 days (16.6 weeks)	18
2019	L. McGonagle	31	Patients were included if they were engaged in paid work in the 3 months prior to surgery	100	The resumption of paid employment at any capacity	91.4% returned to work, and 8.6% did not	91	91	6–12 months	Average: 7.7 weeks	12
2017	K. Stigmar	996	21% of the women and 16% of the men were on sick leave 1 year before TKR	Men: 84, women: 79	On sick leave 2 years after surgery	12% of men and 17% of women were still on sick leave	85 (men: 88, women: 83)	–	2 years	Mean: 15.4 weeks (men: mean = 96 days; women: mean = 117 days)	17
2017	P. Maillette	6	3 participants had been on sick leave before surgery (2 weeks–6 years)	50	RTW at time of interview (occupation classified as manual, nonmanual or mixed manual)	4 returned to work: 2 nonmanual and 2 mixed, 2 patients that were disabled did not RTW: 2 manual	66 ^a	–	6–12 months	Average: 42.9 weeks ^a (10 months)	9
2017	C.E.H. Scott	289	Prior to TKA, 261 patients were working	90	Return to same work, return to any work, nature of work (sedentary, light manual, moderate manual, heavy manual, other) at time of questionnaire	105 (40%) returned to work, 108 (41%) retired. None of the patients were unemployed prior to TKA returned to work. 6% returned to different work, 34% to the same work	36	40	Mean: 3.3 (2–4) year	Mean: 13.5 weeks (2–104 weeks)	14
2016	M. Bardgett	10	Patients were excluded from interview if they were unemployed in the 3 months prior to surgery	100	RTW at time of interview	All patients returned to work	100 ^a	100 ^a	8–35 months	Average: 8.7 weeks ^a	10
2016	C.S. Leichtenberg	56	32% had preoperatively been on sick leave from work due to hip/knee complaints	68	currently working (yes/no) 12 months after TKA (complete return, partial return, no RTW)	40 (71%) of patients that were working prior to TKA fully returned to work. 10 (18%) worked fewer hours and 6 (11%) did not RTW at all.	89	89 (71 full, 18 partial)	1 year	–	19
2015	P.J. Belmont	159	All were army active duty service members	100	Active duty with combat deployment, active duty without combat deployment, no longer with the	18% who medically separated and 82% who either returned to active duty or	82	82	Minimum 2 years	–	13

(Continued)

Table 2 (Continued)

Year	First author	Population size	Preoperative work status	Working before TKA (%)	Definition of postoperative work-status	Postoperative work-status	Working after TKA (%)	Working before TKA and RTW (%)	Time post-op that RTW was measured	Time to RTW	Ref.
2015	C. Tilbury	120	53% TKA patients were working, 8% were unemployed and/or looking for a job, 12% were disabled, 15% were doing household and/or volunteer work, and 8% were retired	53	service due to retirement or expiration of term of service, or having been classified with knee-related medical separation Working currently (yes/no) 1 year postoperatively	56 (89%) returned to work, 5% were on sick leave, 8% retired	47	89	1 year	Mean: 12.9 weeks	7
2015	B.D. Kleim	50	37 patients were employed preoperatively	74	Returned to work at time of questionnaire	41 patients (82% of the total study population) returned to work	82	–	6 months–3 years	Mean: 13 weeks	21
2014	A.V. Lombardi	661	494 patients reported to be working for pay in the 3 months before their TKA	75	Returned to work at some point after recovery from surgery at time of telephone survey	98% returned to work, 91% returned to their usual job either with or without restrictions, 9% did not return to the same job	73	98	1–3 year	Average: 8.9 weeks	15
2010	J.A.J. Foote	41	66% was in employment before surgery	66	In employment following surgery, working prior to surgery, and able to return to it at time of questionnaire	56% were in employment following surgery, 82% of patients working prior to surgery were able to return to it	56	82	Mean: 41.4 (14–58) months	Mean: 12 weeks	20
2009	H. Lyall	56	41 patients were employed prior to their total knee replacement	73	Employment status postoperatively at time of the questionnaire	98% who were employed before their operation returned to work. All 15 patients were unemployed before their total knee replacement and none returned to work postoperatively	71	98	Mean: 64 (47–112) months	Average: 10 weeks	16

Abbreviations: Ref. reference; RTW, return to work; SD, standard deviation; TKA, total knee arthroplasty; TKR, total knee replacement.

^aQualitative study: “RTW rates” and “time to RTW” not included in reported range.

Table 3 Factors influencing return to work after TKA

First Author	Associated factors	Limiting of beneficial factors	Statistical analysis	Ref
A.R. Boerma	In the final model, two covariates remained: having primarily physically demanding work (total PA level: $B = 0.499$, $SE = 0.193$, $p = 0.012$; leisure-time PA level: $B = 0.552$, $SE = 0.193$, $p = 0.005$), and having mixed work tasks (total PA level: $B = 0.456$, $SE = 0.171$, $p = 0.009$, leisure-time PA level: $B = 0.486$, $SE = 0.176$, $p = 0.007$). Patients who had a more physically demanding job or a combination of a physically and mentally demanding job needed more time to RTW than those with a nonphysically demanding job	Limiting: <ul style="list-style-type: none"> Physically demanding job^b ($p = 0.012$) Having mixed work tasks^b ($p = 0.009$) 	Multivariate analysis	11
P. Lankinen	Patients with higher non-manual occupational status had 2.8 (95% CI: 2.2–3.7; in mutually adjusted model HR = 2.62; CI: 1.95–3.52; $p < 0.0001$) times higher rate of RTW as compared with patients with manual labor occupational status. Low level of sickness absence (<30 days) before the surgery was associated with a 2.4 (95% CI: 1.9–3.0) times higher rate of RTW as compared with patients with longer sickness absence. Those patients whose self-rated health was good were 1.4 (95% CI: 1.1–1.7) times more likely to RTW comparing to those with poor self-rated general health. Self-rated health was not associated with RTW in the mutually adjusted model	Beneficial: <ul style="list-style-type: none"> Nonmanual work^b ($p < 0.0001$) Low number of preoperative sickness absence days^b ($p < 0.0001$) Good self-rated health ($p = 0.0012$)^a 	Multivariate analysis	18
L. McGonagle	Multiple regression analysis also identified a significantly earlier time of RTW if flexible work conditions were resumed ($p = 0.003$). Furthermore, those in less physically demanding jobs (sedentary, light, and medium) were more likely to RTW with unchanged conditions of employment compared with those with more physically demanding jobs (heavy and very heavy) (49.5 vs. 11.8%). Those with more physically demanding jobs were more likely to RTW with both reduced hours and reduced level of duties	Beneficial: <ul style="list-style-type: none"> Flexible work conditions^b ($p = 0.003$) Less physically demanding jobs and RTW with unchanged conditions of employment^b 	Multivariate analysis	12
K. Stigmar	Women generally had more sick leave before surgery and a slower RTW after both THR and TKR	Limiting: Women	–	17
P. Maillette	Participants' representations arose from expectations of the surgery, representations of their current condition, and perceived support from the various environmental systems. The patients that did RTW felt they had experienced greater improvement after surgery and received concrete support from their workplace, which facilitated their RTW in their view	Limiting: <ul style="list-style-type: none"> High expectations of surgery Absence of support from various environmental systems Beneficial: <ul style="list-style-type: none"> Support from the workplace 	–	9
C.E.H. Scott	The functional outcome of TKA, as assessed by PROMs, was significantly associated with a return to any work on univariate analysis. Multivariate analysis found that, in patients aged ≤ 65 years working prior to TKA, age ($B = 0.156$, CI: 0.07–0.24; $p < 0.001$) and heavy or moderate manual work ($B = 0.960$, CI:	Limiting: <ul style="list-style-type: none"> Older age^b ($p < 0.001$) Physical nature of employment (manual jobs;^b $p = 0.03$) Beneficial: <ul style="list-style-type: none"> Functional outcome of TKA^a 	Univariate and multivariate analysis	14

(Continued)

Table 3 (Continued)

First Author	Associated factors	Limiting of beneficial factors	Statistical analysis	Ref
	0.18–1.74; $p = 0.03$) independently predicted both return to any work and return to the same work			
M. Bardgett	Three themes were identified that influenced the process of RTW, from the patient's perspective. These were delays in surgical intervention, limited and often inconsistent advice from healthcare professionals regarding RTW, and finally the absence of rehabilitation to optimize patient's recovery and facilitate RTW	Limiting: <ul style="list-style-type: none"> • Delay in surgical intervention • Limited advice from healthcare professionals • Absence of rehabilitation 	–	10
C.H. Leichtenberg	In TKA patients, the only variable associated with a full RTW was change in KOOS Sport subscale score from baseline ($p = 0.039$). Exploratory multivariable stepwise logistic regression indicated that self-employment ($p = 0.019$), preoperative absence from work ($p = 0.001$) and baseline KOOS ADL subscale scores ($p = 0.272$) were associated with a RTW. On univariate analysis, patients who did not or only partially returned to work were significantly more likely be older ($p = 0.010$), have a lower level of education ($p = 0.043$), be self-employed ($p = 0.019$) and have preoperative absence from work ($p = 0.001$)	Limiting: <ul style="list-style-type: none"> • Preoperative absence from work^b ($p = 0.001$) • Being self-employed^b ($p = 0.019$) • Older age^a • Lower level of education^a Beneficial: <ul style="list-style-type: none"> • Change in KOOS Sport subscale score from baseline^a • KOOS ADL subscale scores ($p = 0.272$) 	Univariate and multivariate analysis	19
P.J. Belmont	Multivariate analysis showed that the <45 years age group had a significantly increased odds ratios for being medically separated (OR = 2.36; 95% CI: 1.14–4.90; $p = 0.0206$)	Limiting: <ul style="list-style-type: none"> • <45 years group^b ($p = 0.0206$) 	Univariate and multivariate analysis	13
C. Tilbury	Comparison between those who had returned to work ($n = 64$ and $n = 56$) as compared with those who had not returned to work after 1 year and were not retired ($n = 5$ and $n = 6$), did not show any statistically significant differences	No statistically significant differences shown	Mann–Whitney <i>U</i> -test, Chi-squared test	7
B.D. Kleim	Patients who have preoperative sick leave due to their hip or knee arthritis take 4.6 weeks longer to RTW than those who do not. Patients with level 2 or 3 qualifications returned to work (mean = 9.9 weeks) significantly more quickly than patients with no qualifications or level 1 (secondary education) qualifications (mean = 12.6 weeks). Those with further education beyond the required secondary education, represented by levels 2 and 3, returned more quickly. Patients in occupations with a manual level of 0 on average returned to work 2.5 weeks faster than those in occupations with a manual level of 1 ($p = 0.026$). Patients in occupations with a manual level of 1 returned to work on average 6.2 weeks faster than patients in occupations with a manual level of 2	Limiting: <ul style="list-style-type: none"> • Preoperative sick leave^a Beneficial: <ul style="list-style-type: none"> • Higher education^a • Nonmanual jobs^a 	Univariate analysis, <i>t</i> -test, Chi-squared test	21
A.V. Lombardi	Stratified by preoperative physical demand category, there was no difference between groups in either need for restrictions or length of time worked with postoperative restrictions	No statistically significant differences shown for physical demand category	Chi-squared test	15

Table 3 (Continued)

First Author	Associated factors	Limiting of beneficial factors	Statistical analysis	Ref
J.A.J. Foote	No statistically significant difference in physical intensity of pre- versus postoperative occupation was found. 12% found that their ability to work was significantly worse, 37% found that their ability to work was significantly better	No statistically significant differences shown for physical intensity of occupation	Chi-squared test, Fisher's exact test	²⁰
H. Lyall	None of the unemployed patients returned to work. Of the 15 patients unemployed before surgery 12 had previously undertaken manual work. 98% of the patients employed before surgery did RTW	Limiting: • Being unemployed before surgery • Manual work Beneficial: • Being employed before surgery	–	¹⁶

Abbreviations: ADL, activities of daily living; CI, confidence interval; HR, hazard ratio; OR, odds ratio; KOOS Score, Knee Injury and Osteoarthritis Outcome Score; PA, physical activity; PROM, patient reported outcome measures; Ref., reference; RTW, return to work; SE, standard error; THR, total hip replacement; TKA, total knee arthroplasty; TKR, total knee replacement.

^aStatistically significant.

^bStatistically significant with multivariate analysis.

($p = 0.003^{12}$), older age ($p < 0.001^{14}$), being self-employed ($p = 0.019^{19}$), and age less than 45 years in a military population ($p = 0.0206^{13}$). Factors also shown to be statistically significant were level of education, functional outcome of TKA, change in Knee Injury Osteoarthritis Outcome Score (KOOS) Sport subscale score from baseline, being self-employed, and a good self-rated health.

The factor most frequently reported was the physical nature of the employment. Six different studies showed that a physically demanding job had a negative impact on RTW of which four studies reported after applying multivariate analysis. Preoperative absence from work was reported by four different studies to be influencing RTW negatively, of which two using multivariate analysis. Factors reported by two different studies were older age, level of education,

functional outcome after TKA, and support from the environment and from health care professionals.

Methodological Quality

► **Table 4** summarizes the risk of bias in each domain for all included studies according to the QUIPS tool. Four studies scored low risk of bias in each domain. In eight studies, risk of bias was scored as moderate in one or more domains. In two studies, high risk of bias was found in the “study participation” domain. These last two were qualitative studies using purposive sampling.

Discussion

This systematic review provides important insights regarding RTW after TKA. Overall, the majority of patients reported RTW

Table 4 Risk of bias according to QUIPS tool

First author	Study participation	Study attrition	Prognostic factor measurement	Outcome measurement	Study confounding	Statistical analysis and reporting	Ref.
A.R. Boerma	Low bias	Low bias	Low bias	Low bias	Low bias	Low bias	¹¹
P. Lankinen	Moderate bias	Low bias	Moderate bias	Low bias	Moderate bias	Low bias	¹⁸
L. McGonagle	Moderate bias	Low bias	Moderate bias	Low bias	Moderate bias	Low bias	¹²
K. Stigmar	Low bias	Low bias	Low bias	Low bias	Low bias	Low bias	¹⁷
P. Maillette	High bias	Low bias	Moderate bias	Moderate bias	Low bias	Low bias	⁹
C.E.H. Scott	Low bias	Moderate bias	Low bias	Low bias	Low bias	Low bias	¹⁴
M. Bardgett	High bias	Moderate bias	Low bias	Low bias	Moderate bias	Moderate bias	¹⁰
C.S. Leichtenberg	Low bias	Moderate bias	Low bias	Low bias	Low bias	Low bias	¹⁹
P.J. Belmont	Low bias	Low bias	Low bias	Low bias	Low bias	Low bias	¹³
C. Tilbury	Low bias	Low bias	Low bias	Low bias	Low bias	Low bias	⁷
B.D. Kleim	Moderate bias	Moderate bias	Low bias	Low bias	Moderate bias	Moderate bias	²¹
A.V. Lombardi	Low bias	Moderate bias	Low bias	Low bias	Low bias	Low bias	¹⁵
J.A.J. Foote	Low bias	Moderate bias	Low bias	Low bias	Moderate bias	Low bias	²⁰
H. Lyall	Low bias	Moderate bias	Low bias	Low bias	Moderate bias	Moderate bias	¹⁶

Abbreviations: QUIPS, quality in prognosis studies; Ref., reference.

after TKA but the numbers reported by the included studies were quite divergent.

RTW rate ranged from 40 to 98% (mean, 89%) and the mean time taken to RTW ranged from 7.7 to 16.6 weeks (mean, 13 weeks). In three studies specifying whether patients returned to the same or different jobs, the majority of patients returned to the same job.^{14–16}

One study conducted in the United Kingdom by Scott et al reported an RTW rate of 40%.¹⁴ This is remarkably lower than the other included studies, reporting RTW rates ranging from 71 to 98%. This outlier cannot be explained by the circumstances of the country, as three other studies also conducted in the United Kingdom presented higher RTW rates (82–98%).^{16,20,21} The majority of patients in this study had a manual job (39% moderate or heavy manual labor). This was difficult to compare with other studies, as other criteria for physical nature of work were used. However, higher RTW rates were reported in other studies where the majority of patients also had manual jobs (52% manual jobs and 87% RTW).¹⁸ Moreover, Scott et al could not find any statistically significant correlation between heavy- or moderate-manual work and RTW ($p = 0.03$). It also has to be noted that this study has the highest mean age among included studies, and that in this study, the majority of patients not returning to work was aged >60 years. However, this age difference compared with other included studies is too small to explain the big difference in RTW rate. Lastly, quality of this study scored good according to the QUIPS tool.

Only two previous systematic reviews on RTW after TKA exist to our knowledge. In 2009, Kuijer et al reported on beneficial and limiting factors affecting RTW in patients undergoing TKA or THA.²² They concluded to almost a complete lack of literature on influencing factors on RTW after TKA as only three studies were included in the review of which only one regarding TKA. Later in 2014, Tilbury et al showed that the literature on work status after THA is more extensive than for TKA.⁷ Out of 19 studies included, only 4 provided information on work status after TKA and only 3 were determinants of work status after TKA. They found factors associated with a faster RTW to be female sex, self-employment, higher mental and physical health scores, higher functional comorbidity index scores, and a handicap accessible workspace. Factors associated with a slower RTW were having less pain preoperatively, having a more physically demanding job, and having worker's compensation. These factors were all shown to be significant after multivariate adjustment. However, none of these factors were significantly associated with RTW within 3 months after surgery after multivariate adjustment.²³ In our study, we also found a physically more demanding job to be associated with a slower or no RTW. Regarding the self-employment and female sex, we found contradictory results. However, in the article included in our review, female sex was not investigated using statistical analysis or multivariate analysis.¹⁷ Concerning for self-employment, the study included in our review also used multivariate analysis. The authors of this study suggest that this may be related to the observation that self-employed patients generally work more hours than wage earners.

In recent years, there seems to be an increase in interest concerning RTW after TKA. Twelve of the 14 studies included in this review were only recently publicized between 2014 and 2019. Since RTW might have changed over the last years due to rising retirement age, with economic changes and the continuous development of new techniques and prosthesis, the present review can provide valuable new insights in this area.

RTW after major surgery is dependent on various factors such as social and cultural factors, adaptations at the workspace, or the social safety net in the country. These are all factors we could not take into account in this review. We did, however, find no differences in RTW rate between studies conducted in different countries.

It is often unclear whether not returning to work was indeed attributable to the TKA, or if the decision to retire after the operation was already made before surgery. Some patients might postpone the surgery until a time when retirement was possible. However, by limiting the age to 65 years, we partially reduced this bias.

We observed a wide variation in both the definition and timeframe used to measure the work status pre- and postoperatively. Therefore, we did not attempt any statistical analyses as comparisons between studies would be seriously distorted by this heterogeneity. However, efforts were made to minimize the study heterogeneity by applying an upper age limit unlike previous systematic publications. We also obtained a much bigger study population compared with previous reviews, including 14 articles and a total study population of 3,073 patients. To decrease this heterogeneity in future research, standardized uniform outcome measures and protocols for studies investigating RTW after TKA are warranted.

Limitations and Strengths

We acknowledge several limitations to this study. First of all, we included only articles in English, excluding possibly valuable articles written in any other language. Also, all studies included were conducted in Western countries. Since Eastern countries might differ from Western countries both economically and culturally, conclusions drawn from this review may not be expandable to Eastern countries.

Another weakness to this review is the fact that only 4 of the 14 studies have a prospective design. RTW after a joint replacement is a process with multiple factors involved like retirement age, economic situation and social safety net in the country, surgical techniques, and type of the prosthesis used. A retrospective study is limited by several forms of bias and is therefore not optimal to evaluate the RTW process within its context. More qualitative prospective studies with a sufficient population are therefore warranted.

Furthermore, from the 14 studies included only 6 used multivariate analysis to investigate the correlation between risk factors and RTW. Because of the multitude of factors influencing RTW, adjustment for confounders is essential in these kinds of studies.

At last we encountered a wide variation in study quality, with a risk of bias ranging from low to high.

Conclusion

In conclusion, this systematic review found the rate of RTW after TKA to be ranging from 40 to 98% at a mean of 7.7 to 16.6 weeks. Most important factors associated with a slower or no RTW were a more physical nature of employment and preoperative absence from work.

RTW after TKA is a field of growing interest, with more articles being publicized in the recent years. However, there is still a lack of qualitative large studies applying multivariate analysis. Also, the implication of uniform standardized outcome measures could accommodate a better comparison of studies, benefiting our knowledge regarding this subject. A better understanding of determinants that influence RTW after TKA could after all contribute a more effective and focused management and better preventive measures.

Authors' Contributions

All authors contributed to the study conception and design. Conceptualization was done by D.V.L. and H.V. Data collection and analysis were performed by D.V.L. and J.N. The first draft of the manuscript was written by D.V.L. Review and editing were done by D.V.L., J.N., H.V., and P.B. All authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding

None.

Conflict of Interest

None declared.

References

- 1 Peat G, McCarney R, Croft P. Knee pain and osteoarthritis in older adults: a review of community burden and current use of primary health care. *Ann Rheum Dis* 2001;60(02):91–97
- 2 Singh JA, Lewallen DG. Patient-level improvements in pain and activities of daily living after total knee arthroplasty. *Rheumatology (Oxford)* 2014;53(02):313–320
- 3 Lange JK, Lee Y-YY, Spiro SK, Haas SB. Satisfaction rates and quality of life changes following total knee arthroplasty in age-differentiated cohorts. *J Arthroplasty* 2018;33(05):1373–1378
- 4 Leskinen J, Eskelinen A, Huhtala H, Paavolainen P, Remes V. The incidence of knee arthroplasty for primary osteoarthritis grows rapidly among baby boomers: a population-based study in Finland. *Arthritis Rheum* 2012;64(02):423–428
- 5 Kurtz SM, Lau E, Ong K, Zhao K, Kelly M, Bozic KJ. Future young patient demand for primary and revision joint replacement: national projections from 2010 to 2030. *Clin Orthop Relat Res* 2009;467(10):2606–2612
- 6 Finnish Centre for Pensions. Retirement ages. Accessed September 10, 2020 at: <https://www.etk.fi/en/work-and-pensions-abroad/international-comparisons/retirement-ages/>
- 7 Tilbury C, Schaasberg W, Plevier JWMJWM, Fiocco M, Nelissen RGHH, Vliet Vlieland TP. Return to work after total hip and knee arthroplasty: a systematic review. *Rheumatology (Oxford)* 2014;53(03):512–525
- 8 Hayden JA, Côté P, Bombardier C. Evaluation of the quality of prognosis studies in systematic reviews. *Ann Intern Med* 2006;144(06):427–437
- 9 Maillette P, Coutu M-FF, Gaudreault N. Workers' perspectives on return to work after total knee arthroplasty. *Ann Phys Rehabil Med* 2017;60(05):299–305
- 10 Bardgett M, Lally J, Malviya A, Deehan D. Return to work after knee replacement: a qualitative study of patient experiences. *BMJ Open* 2016;6(02):e007912
- 11 Boersma AR, Brouwer S, Koolhaas W, et al. No association between preoperative physical activity level and time to return to work in patients after total hip or knee arthroplasty: a prospective cohort study. *PLoS One* 2019;14(09):e0221932
- 12 McGonagle L, Convery-Chan L, DeCruz P, Haebich S, Fick DP, Khan RJK. Factors influencing return to work after hip and knee arthroplasty. *J Orthop Traumatol* 2019;20(01):9
- 13 Belmont PJ Jr., Heida K, Keeney JA, Hamilton W, Burks R, Waterman BR. Return to work and functional outcomes following primary total knee arthroplasty in U.S. military servicemembers. *J Arthroplasty* 2015;30(06):968–972
- 14 Scott CEH, Turnbull GS, MacDonald D, Breusch SJ. Activity levels and return to work following total knee arthroplasty in patients under 65 years of age. *Bone Joint J* 2017;99-B(08):1037–1046
- 15 Lombardi AV Jr., Nunley RM, Berend KR, et al. Do patients return to work after total knee arthroplasty? *Clin Orthop Relat Res* 2014;472(01):138–146
- 16 Lyall H, Ireland J, El-Zebdeh MY. The effect of total knee replacement on employment in patients under 60 years of age. *Ann R Coll Surg Engl* 2009;91(05):410–413
- 17 Stigmar K, Dahlberg LE, Zhou C, Jacobson Lidgren H, Petersson IF, Englund M. Sick leave in Sweden before and after total joint replacement in hip and knee osteoarthritis patients. *Acta Orthop* 2017;88(02):152–157
- 18 Lankinen P, Laasik R, Kivimäki M, et al. Are patient-related preoperative factors influencing return to work after total knee arthroplasty. *Knee* 2019;26(04):853–860
- 19 Leichtenberg CS, Tilbury C, Kuijjer P, et al. Determinants of return to work 12 months after total hip and knee arthroplasty. *Ann R Coll Surg Engl* 2016;98(06):387–395
- 20 Foote JAJ, Smith HK, Jonas SC, Greenwood R, Weale AE. Return to work following knee arthroplasty. *Knee* 2010;17(01):19–22
- 21 Kleim BD, Malviya A, Rushton S, Bardgett M, Deehan DJ. Understanding the patient-reported factors determining time taken to return to work after hip and knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2015;23(12):3646–3652
- 22 Kuijjer PPFM, de Beer MJPM, Houdijk JHP, Frings-Dresen MHW. Beneficial and limiting factors affecting return to work after total knee and hip arthroplasty: a systematic review. *J Occup Rehabil* 2009;19(04):375–381
- 23 Styron JF, Barsoum WK, Smyth KA, Singer ME. Preoperative predictors of returning to work following primary total knee arthroplasty. *J Bone Joint Surg Am* 2011;93(01):2–10