Pleurabrade: A Spiral Brush for Mechanical Pleurodesis and a Review of the Literature

John S. Young¹ Desiree A. Steimer¹ Emily Polhemus¹ Raphael Bueno¹

¹Division of Thoracic Surgery, Brigham and Women's Hospital, Boston, Massachusetts, United States

Thorac Cardiovasc Surg Rep 2021;10:e36–e38.

Address for correspondence John Young, MD, Department of Thoracic Surgery, Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115, United States (e-mail: jsyoungthoracic@gmail.com).

Abstract

Keywords

- ▶ mechanical pleurodesis
- ▶ spontaneous pneumothorax
- ▶ pleural abrasion

Background While the optimal treatment for primary spontaneous pneumothorax remains unclear, mechanical pleurodesis is a well-established treatment. The Pleurabrade is a spiral brush designed for mechanical pleurodesis during thoracoscopy. We present two patients who underwent mechanical pleurodesis with the Pleurabrade.

Case Description Two patients with spontaneous pneumothorax underwent operative intervention including mechanical pleurodesis with the Pleurabrade. Chest tubes were removed within 48 hours postoperatively and they were discharged home. Both patients remain recurrence free at 11 and 22 months, respectively.

Conclusion While further testing is needed, these case reports and operative video highlight the Pleurabrade as an efficient device for thoracoscopic mechanical pleurodesis.

Introduction

Primary spontaneous pneumothorax is frequently seen in young, tall men and is typically the result of rupture of apical blebs.¹ Recurrence of an ipsilateral spontaneous pneumothorax remains relatively low, around 25%, so many clinicians will manage an initial occurrence less invasively (observation, needle decompression, or tube thoracostomy).^{2,3} However, recurrence risk increases with subsequent events. Thus, operative intervention is typically reserved for a recurrence or those cases with high-risk features such as tension pneumothorax, or occupational or recreational risks (pilots or scuba divers).² Surgical options include mechanical pleurodesis with bleb resection, or chemical pleurodesis; however, the optimal means of preventing recurrence remains unclear.

Pleural abrasion is an accepted means of mechanical pleurodesis and is thought to have a lower rate of bleeding complications compared with pleurectomy, which is usually reserved for recurrence after a surgical treatment for pneumothorax.⁴ Maintaining the pleura also makes reoperation in the chest easier compared with pleurectomy.⁵ Many surgeons will abrade the pleura with an electrocautery scratch pad or a sponge.⁶

These have a small point of contact with the pleura and are therefore somewhat more cumbersome, time consuming, and are at risk of becoming a retained foreign body.⁶ We report herein two patients who successfully underwent mechanical pleurodesis with the Pleurabrade (Butterfield Technologies, Portland, Maine, United States), an instrument with nylon bristle brushes specifically designed for mechanical pleurodesis. The Pleurabrade (Butterfield Technologies) overcomes some of the limitations of the other techniques of pleural abrasion.

Case Presentations

Case 1

An otherwise healthy 24-year-old man presented at an outside hospital with 2 days of right-sided chest pain and dyspnea. A chest X-ray showed a moderate to large right pneumothorax and a small associated effusion. A thoracic vent was placed in the emergency room and he was discharged home with the thoracic vent in place. He presented in our clinic 6 days later and repeat imaging showed increased size of his pneumothorax. Based on our recommendation, the patient agreed to operative intervention. He was taken to the operating room

received
July 20, 2020
accepted after revision
November 16, 2020

DOI <https://doi.org/10.1055/s-0041-1723950>.
ISSN 2194-7635.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

and underwent a right video-assisted thoracoscopic surgery (VATS) where a ruptured bleb was identified in the apex and resected using a stapler. The Pleurabrade (Butterfield Technologies) was used to perform a mechanical pleurodesis. After 48 hours of -20 cm H₂O of suction, he tolerated a water seal trial and his chest tube was removed. He was discharged on postoperative day 3. He remains without recurrence at 22 months at the time of reporting.

Case 2

A 25-year-old man with a history of a right spontaneous pneumothorax managed with a tube thoracostomy 3 months prior presented to an outside hospital with acute left-sided chest pain and dyspnea. A chest X-ray showed a left apical pneumothorax and he was transferred to our institution for management. A left pleural 14 Fr pigtail was placed on arrival. He underwent a left VATS; multiple apical blebs were seen and surgically resected. A partial and apical pleurectomy was performed, and the remainder of the pleura was abraded with the Pleurabrade (Butterfield Technologies). Postoperatively, his chest tube was removed after 48 hours of suction and a successful water seal trial. He was discharged on postoperative day 2 and remains without recurrence at 11 months of follow-up.

Discussion

Primary spontaneous pneumothorax is a subset of spontaneous pneumothorax and occurs in patients without prior lung disease and typically affects tall thin males between the ages of 10 and 30 years.¹ It has an incidence of 18 to 28/100,000 cases per year for men and 1.6 to 6/100,000 cases for women.¹ The optimal management of primary spontaneous pneumothorax remains controversial. For initial occurrences, management options include observation, aspiration, tube drainage, chemical pleurodesis, or mechanical pleurodesis with wedge resection of culprit lesions.^{2,3} The option chosen depends on patient as well as clinician factors and preferences.² Large blebs, for instance, may be a reason to operate with a VATS wedge resection and pleurodesis on an initial presentation.⁷ Given the increasing rates of recurrence with each subsequent episode, many surgeons elect to perform a pleurodesis at a second occurrence.²

Pleurodesis aims to obliterate the pleural space by producing extensive adhesion of the visceral and parietal pleura.⁸ While a direct apposition between the two pleural surfaces is necessary, the inflammatory process involving mesothelial cell release of inflammatory mediators, production of fibrin adhesions, and recruitment of fibroblasts and collagen production is critical for a successful pleurodesis.⁸ Techniques for achieving pleurodesis include both mechanical (either by pleural abrasion or parietal pleurectomy) or pleurodesis with a sclerosing agent.

Whether there is a benefit to chemical or mechanical pleurodesis with bleb resection is an area of debate. A 2014 randomized controlled study suggested that the addition of a mechanical pleurodesis did not significantly decrease the risk of recurrence but did add additional bleeding and postoperative drainage.⁹ The report, however, does not clarify which patients

were brought to the operating room at an initial versus recurrent spontaneous pneumothorax.⁹ Several other studies have examined this issue and while a 2019 systemic review and meta-analysis suggested that following bullectomy, chemical pleurodesis is superior to mechanical pleurodesis in reducing hospital stay and recurrence rate, they acknowledged that more directed randomized controlled trials with a longer follow-up period are necessary.¹⁰ Thoracoscopic pleural abrasion has been established as a safe means of mechanical pleurodesis,¹¹ although some argue that it has a higher recurrence rate than pleurectomy or chemical pleurodesis.^{4,10} One hypothesis for the reduced recurrence with chemical pleurodesis is that instilled agents may reach areas that are hard to reach with conventional instruments during thoracoscopy.¹⁰

The concept of utilizing a brush for pleurodesis is not a new one and was described in 1976, predating video-assisted thoracoscopy.¹² In fact, an electric rotating brush has also been described, although it has not gained widespread popularity.⁶ A simpler variation of a rotating brush is the Pleurabrade (Butterfield Technologies) that was utilized in these two patients. It is a nylon bristle brush on a twisted stainless steel wire at the end of a stainless steel shaft. It can be bent to better contour to the curved pleural surfaces and can fit through a 5-mm port. Compared with other means of achieving pleural abrasion such as using a scratch pad or a sponge, the benefits of the Pleurabrade (Butterfield Technologies) for mechanical pleurodesis are that it is less likely to be left in the chest as a retained foreign body and that it can abrade the pleural surfaces more widely leading to a more efficient mechanical pleurodesis.

As is evident from the operative videos, the Pleurabrade (Butterfield Technologies) can be efficiently used to perform a mechanical pleurodesis in patients with primary spontaneous pneumothorax. Both patients in this series had no postoperative complications and are recurrence free. We believe this is a safe instrument in the arsenal of the thoracic surgeon managing primary spontaneous pneumothorax, although further studies need to be performed to document its safety profile, efficacy, and durability compared with other modalities (► **Video 1**).

Video 1

Video showing the case presentations and operative usage of the Pleurabrade (Butterfield Technologies, Portland, Maine, United States). Online content including video sequences viewable at: <https://www.thieme-connect.com/products/ejournals/html/10.1055/s-0041-1723950>

Conclusion

VATS mechanical pleurodesis by pleural abrasion as a treatment for primary spontaneous pneumothorax (particularly in recurrent cases or cases with high-risk features) can be thoroughly and efficiently accomplished using the

Pleurabrade (Butterfield Technologies) as depicted in the operative video. The device reduces the risk of foreign body retention and can be better contoured than other abrasion techniques (such as using a scratch pad or sponge) to reach all the pleural surfaces. We believe it is a safe and effective instrument for mechanical pleurodesis, although further studies are needed.

Conflict of Interest

Dr. Bueno reports personal fees from Butterfield Technologies/Pleurabrade, during the conduct of the study; grants from NCI, grants from NIBIB, grants from DOD, grants from Roche, grants from GeneTech, grants from Siemens, grants from Merck, grants from Verastem, grants from Gristone, grants from MedGenome, grants from Epizyme, other from Navigation Sciences, personal fees from Novocure, personal fees from Intuitive, outside the submitted work.

References

- 1 Melton LJ III, Hepper NG, Offord KP. Incidence of spontaneous pneumothorax in Olmsted County, Minnesota: 1950 to 1974. *Am Rev Respir Dis* 1979;120(06):1379–1382
- 2 MacDuff A, Arnold A, Harvey JBTS Pleural Disease Guideline Group. Management of spontaneous pneumothorax: British Thoracic Society Pleural Disease Guideline 2010. *Thorax* 2010;65 (Suppl 2):ii18–ii31
- 3 Brown SGA, Ball EL, Perrin K, et al;PSP Investigators. Conservative versus interventional treatment for spontaneous pneumothorax. *N Engl J Med* 2020;382(05):405–415
- 4 Ng C, Maier HT, Kocher F, et al. VATS partial pleurectomy versus VATS pleural abrasion: significant reduction in pneumothorax recurrence rates after pleurectomy. *World J Surg* 2018;42(10): 3256–3262
- 5 Coosemans W, Lerut TE, Van Raemdonck DE. Thoracoscopic surgery: the Belgian experience. *Ann Thorac Surg* 1993;56(03): 721–730
- 6 Maier A, Anegg U, Renner H, et al. Four-year experience with pleural abrasion using a rotating brush during video-assisted thoracoscopy. *Surg Endosc* 2000;14(01):75–78
- 7 Olesen WH, Katballe N, Sindby JE, et al. Surgical treatment versus conventional chest tube drainage in primary spontaneous pneumothorax: a randomized controlled trial. *Eur J Cardiothorac Surg* 2018;54(01):113–121
- 8 Rodriguez-Panadero F, Montes-Worboys A. Mechanisms of pleurodesis. *Respiration* 2012;83(02):91–98
- 9 Min X, Huang Y, Yang Y, et al. Mechanical pleurodesis does not reduce recurrence of spontaneous pneumothorax: a randomized trial. *Ann Thorac Surg* 2014;98(05):1790–1796, discussion 1796
- 10 Sim SKR, Nah SA, Loh AHP, Ong LY, Chen Y. Mechanical versus chemical pleurodesis after bullectomy for primary spontaneous pneumothorax: a systemic review and meta-analysis. *Eur J Pediatr Surg* 2019;s-0039–s-1697959
- 11 Gossot D, Galetta D, Stern JB, et al. Results of thoracoscopic pleural abrasion for primary spontaneous pneumothorax. *Surg Endosc* 2004;18(03):466–471
- 12 Kurzweg FT, Williamson JW, Daron PB, Danna SJ. Brush pleurodesis. *Am Surg* 1976;42(08):581