Recurrent Sphenocavernous Meningioma

Mizuho Inoue¹ Mohamed Labib² Alexander Yang¹ A. Samy Youssef³

¹Department of Neurosurgery, University of Colorado, Aurora, Colorado, United States

²Department of Neurosurgery, Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, Phoenix, Arizona, United States

³Departments of Neurosurgery and Otolaryngology, University of Colorado, Aurora, Colorado, United States

Address for correspondence A. Samy Youssef, MD, PhD, Departments of Neurosurgery and Otolaryngology, University of Colorado, Mail Stop C307, 12631 East 17th Avenue, Aurora, CO 80045, United States (e-mail: samy.youssef@cuanschutz.edu).

J Neurol Surg B Skull Base 2022;83(suppl S3):e610.

Abstract

A case of a recurrent sphenocavernous meningioma is presented. The patient is a 42-year-old male who presented with an episode of transient right-sided numbness. A magnetic resonance imaging (MRI) revealed a large left sphenocavernous meningioma. The patient underwent a frontotemporal craniotomy for tumor resection. Near total resection was achieved with minimal residual in the left cavernous sinus (CS) and orbital apex. The pathology was consistent with meningioma, World Health Organization (WHO) grade I. A follow-up MRI was done 9 months after surgery and showed a growth of the residual tumor, which was treated with intensity modulated radiotherapy. Tumor growth was detected on serial imaging over a 4-year period. Surgical resection was offered. A left frontotemporal craniotomy with pretemporal transcavernous approach was performed. The bone flap was reopened and the dura was opened in a Y-shaped fashion. The roof of the optic canal was drilled off, and the falciform ligament was opened to decompress the optic nerve. The tumor was disconnected from the anterior clinoid region (the anterior clinoid process was eroded by the tumor) and reflected off the wall of the lateral CS. Tumor was adherent to the V2 fascicles (the lateral CS wall was resected in the first surgery) and was sharply dissected off. Gross total resection was achieved. The pathology was consistent with meningioma, WHO grade I. The patient had an unremarkable postoperative course without any new neurological deficits. The link to the video can be found at: https://youtu.be/KVBVw_86JqM.

Keywords

- recurrent
- sphenocavernous
- meningioma
- cavernous sinus

Conflict of Interest

A.S.Y. is a consultant for Stryker Inc. and obtains royalties from Mizuho America Inc.

www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

received
October 19, 2020
accepted
October 27, 2020
published online
June 8, 2021

DOI <https://doi.org/10.1055/s-0041-1725941>.
ISSN 2193-6331.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany