



Extended Endoscopic Endonasal Approach for a Giant Parasellar Epidermoid Cyst

Sorin Aldea¹  Mathieu Veyrat² Pierre Bourdillon¹ Denis Ayache² Caroline Le Guérinel¹

¹Department of Neurosurgery, Rothschild Foundation Hospital, Paris, France

²Department of Otolaryngology—Head and Neck Surgery, Rothschild Foundation Hospital, Paris, France

Address for correspondence Sorin Aldea, MD, Department of Neurosurgery, Rothschild Foundation Hospital, 25 rue Manin, 75019 Paris, France (e-mail: saldea@for.paris).

J Neurol Surg B Skull Base 2022;83(suppl S3):e653–e654.

Abstract

Epidermoid cysts are rare lesions which typically grow slowly. For this reason, these lesions are usually discovered when they are already very large. The parasellar location is no exception to this rule and may involve the cavernous sinus or the Meckel cave. We present a 34-year-old female patient without past medical history who was admitted in our tertiary referral center for episodes of diplopia in the right lateral gaze and right trigeminal dyesthesias. Brain magnetic resonance imaging (MRI) showed a large right parasellar mass with mixed intensity signal on the T1 and T2 sequences, without contrast enhancement and a typical hypersignal intensity on diffusion-weighted sequences evoking an epidermoid cyst. We discuss the radiologic criteria which differentiate the lesions originating in the cavernous sinus from those of the Meckel cave (► **Figs. 1** and **2**).

Parasellar tumors may be approached through classical transcranial approaches such the epidural temporopolar or the subtemporal approach which involve a significant degree of brain retraction. The last decade witnessed the advent of extended endonasal approaches which offer an interesting alternative and avoid the manipulation of the brain. We used the endoscopic transpterygoid approach in our patient and we were able to achieve an excellent clinical and radiological result. We discuss the nuances of the technique and present the surgical steps of the procedure (► **Figs. 3** and **4**).

Keywords

- endoscopy
- parasellar area
- epidermoid cyst

The endoscopic endonasal approach represents an excellent therapeutic option for parasellar lesions. A thorough knowledge of the anatomy and experience with endoscopic techniques are obvious prerequisite.

The link to the video can be found at: <https://youtu.be/QonSvHrCwOU>.



www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

received
March 29, 2020
accepted after revision
January 10, 2021
published online
May 3, 2021

DOI <https://doi.org/10.1055/s-0041-1727127>.
ISSN 2193-6331.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany




Fig. 1 Preoperative contrast-enhanced T1 MRI showing a large parasellar epidermoid cyst. MRI, magnetic resonance imaging.




Fig. 2 Initial exposure of the tumor bulging in the sphenoid sinus.




Fig. 3 Postoperative contrast-enhanced T1 MRI showing subtotal resection of the epidermoid cyst. MRI, magnetic resonance imaging.




Fig. 4 The operative cavity at the end of the resection showing the junction of the right petrous and paraclival internal carotid artery.

Conflict of Interest
None declared.