associated with significantly longer fluoroscopy time and higher DAP and Dose reference (DR). Although nephrostomy, gastrostomy, and biliary interventions were performed in significantly shorter fluoroscopy times compared to prior studies, this did not translate into significant reduction in DAP or DR in most cases. The result showing a significant positive correlation of BMI and DAP ($R = 0.09410$, $P < 0.0001$). **Conclusion:** This analysis establishes the dosimetry of the most commonly performed vascular and interventional procedures in a tertiary care center in Saudi Arabia. This comparison indicates the need for stricter radiation precautions to further comply with international standards.

P101

Role of Positron Emission Tomography/Computed Tomography in Lesion Characterization and Biopsy Guidance in Malignant Biliary Obstruction

Ashraf Bayoumi Abdallah, Mohamed Mohamed Houseni

National Liver Institute, Menoufia University, Shebin Elkom, Egypt.
E-mail: ashrafbayoumi87@gmail.com

Objectives: This prospective study was conducted to evaluate the diagnostic impact of positron emission tomography/computed tomography (PET/CT) in patients with malignant biliary obstruction (MTBO) and if the metabolic information provided by PET/CT scan adds an incremental benefit while performing PET/CT-guided biopsies. **Methods:** This study was carried out from March 2016 to March 2018 for 52 patients diagnosed/suspected to have MTBO. They were investigated by PET/CT. Results were compared with contrast-enhanced CT in about 42 cases and with magnetic resonance cholangiopancreatography (MRCP) in about 15 cases. A group of patients have undergone PET/CT-guided biopsy (Number 15). **Results:** Our study revealed that sensitivity, specificity, positive predictive value, and negative predictive value for PET/CT were 92%, 75%, 96%, and 60%, respectively, while for CT and MRCP were 68%, 60%, 85%, 20% and 60% 50%, 85%, 20%, respectively. PET/CT is more sensitive than CT and MRCP for lymph nodal and distant metastases detection. In our study, 20/52 patients have changed their management after PET/CT. PET/CT-guided biopsy providing representative sample and definitive diagnosis with technically successful in 100% of patients (15/15). **Conclusion:** PET/CT is more sensitive and specific than CT and MRCP in primary detection and staging of tumors causing MTBO. PET/CT has significant impact on patient management. PET/CT-guided biopsy increases the chance of obtaining a representative sample, minimizing sampling errors, and achieving definitive diagnosis both by directing biopsy to most metabolically active part of lesion and by choosing most feasible and most metabolically active lesion among multiple lesions.

P102

Role of the Cardiac-Synchronized Computed Tomography Angiography in Diagnosis and Follow-Up of the Dissection of Descending Thoracic Aorta

Maria Antonella Ruffino, Chiara Ruggieri, Simona Veglia, Ottavio Davini, Paolo Fonio

A.O.U. Città della Salute e della Scienza, Torino, Italy.
E-mail: mariaantonellaruffino@gmail.com

Objectives: Computed tomography (CT) is crucial for the diagnosis of descending thoracic aortic (DTA) dissection, especially in emergency setting, due to its accuracy and ready availability. An appropriate and accurate imaging protocol permits not only to perform the diagnosis but to improve the clinical outcomes in these cases. Our aim is to identify a possible role of the cardio-synchronized CT angiography (CCTA) in the DTA. **Methods:** Twenty-three CCTAs (retrospective gating) in suspected acute aortic syndromes or in chronic thoracic dissections were retrospectively analyzed measuring total vessel areas (ToTA) and areas of the true lumen (TLA) and false lumen (FLA) (at two levels: 2 cm below the isthmus and 3 cm above the diaphragm), with measurements in two arterial phases (40% and 75% of the cardiac cycle) and in the venous one (no cardio-synchronized acquisition). **Results:** TLA average in the 40% and 75% phases was, respectively, 462 (SD=283) and 419 (SD=276) in the proximal site and 436 (SD=269) and 388 (SD=267) in the distal site, significantly larger in the 40% versus 75% phase at both sites ($P < 0.0001$). The data were inverted with FLA: It was smaller in the arterial phase at 40% than at 75% in both sites (proximal and distal). No significant differences emerged for the ToTA. In the venous phase, no statistically significant differences were seen between TLA, FLA, and ToTA. **Conclusion:** Intimal flap dynamic and lumen variations are significantly affected by the various phases of the cardiac cycle. CCTA in DTA dissection can represent a reliable imaging technique to accurately compare the patient examinations during the follow-up. Furthermore, in the presence of symptoms, CCTA can help promptly identify patients who could require treatment because of transient blood flow reduction, due to intimal flap motion along the cardiac cycle.

P201

Best Response during Repeated Chemoembolization Is the Most Significant Predictor of Survival in Hepatocellular Carcinoma

Ibrahim Alrashidi^{1,2}, Jin Hyoung Kim¹, Chan Park¹, Hee Ho Chu¹, So Yeon Kim¹, Dong Il Gwon¹, Hyun-Ki Yoon¹

¹Prince Sultan Military Medical City, Riyadh, Saudi Arabia,

²Asan Medical City, Seoul, South Korea.

E-mail: dr.ialrashidi@gmail.com

Objectives: Whether initial or best response is the better predictor of overall survival after transarterial chemoembolization (TACE) for hepatocellular carcinoma (HCC) remains controversial. We retrospectively evaluated the clinical implications of initial and best responses during repeated TACE for HCC. **Methods:** This study included adult patients who received a diagnosis of intermediate-stage HCC with preserved liver function between 2007 and 2016 and who were treated with TACE as the first-line treatment. Evaluation of the treatment response was based on the modified Response Evaluation Criteria in Solid Tumors. **Results:** Of the 726 included patients, an objective response (complete response [CR] or partial response [PR]) was observed as the