and can often identify the anatomical cause of the PVT and corrective measures can be taken which might lead to long term improved results. However more studies with larger sample size are required to establish the safety and outcome of this approach. Our experience suggests that IR treatment is feasible and safe in early PVT in LDLT patients with good outcomes in cases where concomitant HAT is not present.

OC4.10

Increasing Efficiency in the Interventional Radiology Division: Multiple Changes in Workflow from Patient Registration to Discharge

Umar Tariq, Jennifer Brent, Gordon Evans, Ashok Bhanushali, Tariq Rahman

Geisinger Medical Center, Danville, USA.

E-mail: utariq@geisinger.edu

Background: We aim to present our experience with measures resulting in increased workflow efficiency in our clinical practice, and to evaluate the effect of these changes on the time elapsed at each stage of the workflow process and how it impact overall workflow efficiency. **Method(s):** We implemented a set of changes at each stage of the workflow process from patient registration, pre-operative workup, procedure, post-operative care, and discharge. Average time for completion of each stage of the process was determined before and after implementing the changes. Weekly case volume (WCV), weekly mean overtime hours (WMOH) and monthly percentage of first case on time (FCOT%) were compared before and after the intervention. Patient profiles including age, sex, and BMI were tracked to account for confounding variables. Student's t-test was used to compare variables before and after intervention. F- test was used to compare variance before and after intervention. A p value of less than 0.05 was considered statistically significant. **Result(s):** No statistically significant difference was seen in the age, sex or BMI of patient population before and after intervention ($p > 0.05$). There was a statistically significant 20% increase in WCV from 200 to 240, 45% decrease in WMOH from 10.8 hours to 4.9 hours and 25% increase in monthly FCOT% from 50% to 75% ($p < 0.01$). **Conclusion(s):** Our workflow intervention resulted in better WCV, WMOH and FCOT%. Improved workflow efficiency is critical in the success of an interventional radiology department, and results in better patient care and overall patient satisfaction.

OC4.11

Association of Concomitant Disease in the Profunda and Femoro-Popliteal Veins to Cumulative Patency and Re-Intervention Rates Following Ilio-Femoral Venous Stenting of Limbs with Postthrombotic Occlusion

Leslie Fiengo, Adam Gwozdz¹, Prakash Saha¹, Lawrence Stephenson¹, Leonardo Jones¹, Nicholas Jackson¹, Justinas Silickas¹, Taha Kahn¹, Soundrie Padayachee¹, Alberto Smith¹, Stephen Black¹

*Department of Vascular, London, ¹Academic Department of Vascular Surgery, School of Cardiovascular Medicine and Science, St Thomas' Hospital, King's College London, London, UK.
E-mail: lesliEFIengo@gmail.com*

Background: Ilio-femoral stent patency is inferior in post-thrombotic disease compared with non-thrombotic venous obstruction. The aim of this study was to examine whether decreased inflow to the stent, caused by intraluminal obstructive disease, was associated with greater risk of re-intervention and inferior long-term patency outcomes. **Method(s):** Consecutive patients (2012-2017) receiving a nitinol venous stent for post-thrombotic disease were included for analysis. Pre-operative ultrasound was used to identify femoral vein (FV), profunda vein (PV), and/or popliteal vein (POPV) intraluminal scarring and/or residual thrombosis, and categorised into one of 3 groups: absence of disease; disease in a single inflow vessel; or disease in more than one inflow vessel. Stent patency was assessed using duplex ultrasonography post-intervention, and re-interventions performed when there was a reduction in stent diameter of $>50\%$ or occlusion. **Result(s):** Of 164 patients treated, cumulative patency was 89% (median follow-up 2.4 yrs; range 46-308 wks). However, 70/164 (43%) patients required re-intervention to maintain patency (median number of re-interventions 2; range 1-6). The respective disease state of inflow vessels are shown in Table 1. Cumulative patency and re-intervention rates were significantly worse in patients with more than one diseased inflow vessel ($P=0.47$, $P=0.004$, respectively). Disease in the FV+PV+POPV was associated with a higher risk of re-intervention (16/25 (64%); HR 2.76; $P=0.009$, 95% CI [1.29, 5.92]), and was a strong predictor of cumulative patency loss compared with patients that had no inflow vessel disease (18/25 (72%) HR 17.26; $P=0.009$, 95% CI [2.02, 147.07]). **Conclusion(s):** Maintaining stent patency in post-thrombotic limbs is influenced by the quality of inflow vessels. Patients with intraluminal scarring and/or residual thrombosis in the FV+PV+POPV should be counselled on their increased risk of patency loss.

P101

Relative Atherosclerotic Sparing of the External Iliac Artery: Possibility of a Less Vasculopathic Arterial Graft Target

Pooya Torkian, Taraneh Faghihi, Reza Talaie¹

Shahid Beheshti University of Medical Sciences, Tehran, Iran, ¹Minnesota Hospital, Minnesota, USA.

E-mail: Pooyaag@yahoo.com

Background: Calcified plaque build-up can increase the risk of subsequent vascular complications in pelvic solid organ transplantation and vascular bypass procedures. The purpose of this study was to assess the degree of atherosclerotic calcification in pelvic vessels. **Method(s):** We retrospectively reviewed the unenhanced computed tomography scans of 197 patients from August 2016 to March 2018. Using calcium-scoring software, we examined the distribution map of calcified plaques, focusing on four different arterial segments—the abdominal aorta, the common iliac artery (CIA), the EIA, and the common femoral artery (CFA)—to determine which one showed the least mural calcification. **Result(s):** A total of 197 patients (118 men, 79 women) with a mean age of 61.19 ± 10.8 years were included in our study. The right EIA segment had the lowest average