omental, cystic, adrenal, intercostal, internal mammary, renal, superior mesenteric and gastroduodenal arteries. Suspect ECS in large subcapsular tumors with exophytic growth, adjacent organ invasion, hypertrophied extrahepatic collaterals and marginal recurrence abutting the liver capsule after TACE or local ablation. During TACE, no or incomplete tumor blush on selective hepatic arterial run, or defect in lipiodol deposition in the mass suggest ECS. Search for ECS is mandatory if follow up imaging shows peripheral defect in lipiodol deposition or enhancing residual component of primary mass. An alternative treatment should be undertaken if TACE through ECS fails. **Conclusion(s):** ECS is common in HCC at initial presentation and increases with repeated TACE sessions. For achieving complete tumor response, active search for signs of ECS should be done before, during and after TACE.

P411

Crush Stent Technique: A New Approach for Occluded Iliac Stent

Ben Mrad Melek, Ben Hammamia Mohamed¹, Miri Rim¹, Gdira Faker¹, Denguir Raouef¹

*Departement of Cardiovascular, La Rabta Hospital of Tunis, ¹Faculty of Medecine, University of Tunis El Manar, Tunis, Tunisia.
E-mail: benmradmelek@yahoo.fr*

Background: Iliac stent thrombosis is problematic. Surgery is often indicated because endovascular recanalization of the occluded stent is difficult. Some interventional cardiologists described a new technique to manage occlusion of coronary stent, the “Crush stent technique” consist on subintimal recanalization and swatting of the “old stent”. The aim of this report was to demonstrate the feasibility and safety of this technique also for iliac artery. **Method(s):** A 55 years old woman was admitted in our department for a critical lower limb ischemia due to thrombosis of left common iliac stent implanted 2 years earlier for claudication. The ankle-brachial index was 0.5. Our initial strategy was to cross intraluminally the occlusion and to use a drug-eluting balloon. However, we failed to achieve an intraluminally stent recanalization, so we performed a subintimal recanalization of the occluded stent. The wire crossed the occlusion completely outside stent through the subintimal space. Firstly, we performed a balloon angioplasty, a 7 mm balloon was inflated at eight ATM pressure and contributed in crushing of the thrombosed stent. Therefore, we deployed an 8 x 80 mm Wall-Stent (Boston Scientific), covering all the common and external iliac artery. **Result(s):** Final angiography was satisfactory showing a patent iliac conduit with a total flatterring of the old stent. Post operative course was favorable with symptomatic relieve, the left ankle-brachial index was improved to 0.85 and duplex ultrasound showed patency of the wall stent at 1 and 6 months. There was no hemodynamic disturbance particularly at the site of the subintimal crush. **Conclusion(s):** Crush stent technique is not only feasible for coronary stent thrombosis but can also be performed for peripheral arterial stent thrombosis especially when intraluminally stent recanalization cannot be achieved. In our knowledge, it is the first report in the literature of a “crushed stent” technique for a thrombosed iliac stent.

P412

Uterine Artery Embolization for the Symptomatic Adenomyosis: Short-Term Follow up using Uterine Fibroid Symptom Quality of Life Questionnaire

Ibrahim Alrashidi, Faisal Alahmari¹, Hayat Matouk Alharthy¹, Abdulaziz Almat'hami¹, Abdulwaheed Alruhaimi¹, Abdulrahman Alkhalifa¹, Fares Garad¹

*Prince Sultan Military Hospital, ¹Ministry of Health, King Fahad General Hospital, Riyadh, Saudi Arabia.
E-mail: dr.ialrashidi@gmail.com*

Background: The purpose of this study was to assess clinical outcomes 3 months after uterine artery embolization (UAE) in the treatment of symptomatic adenomyosis. **Method(s):** in this prospective cohort study, at Prince Sultan Military Medical City recruited patients with symptomatic adenomyosis for UAE. The 3 months' post-intervention outcomes were health-related quality of life (HRQOL), symptom severity scores (SSS). **Result(s):** Eighteen patients with adenomyosis have more than 12 mm thickening of the junctional zone based on MRI imaging (11 with Fibroids are excluded) were treated with UAE between January 2012- April 2018. The 3 months' questionnaire was interviewed in 7 patients. The median age of 47 (28-55). The main clinical presentation was abnormal uterine bleeding and Dysmenorrhea. The median duration of symptom is 24 (12-84) months. A 71.4 % of patients received hormonal therapy prior to intervention. The median hemoglobin pre and post-intervention 11.5 and 12.6 respectively. A 28.6 % of patients received blood transfusion pre-UAE. No patient underwent a secondary hysterectomy during the follow-up period. The HRQOL and SSS scores as measured by UFS-QOL at 3 months after UAE showed The score for distress level has been significantly reduced (Z value= -2.366, p value=0.018) from baseline 4.6 (3.6-5) to after 3 months 2.25 (1-3.37). As well as, the score of symptoms level has been significantly reduced Z value= -2.366, p value=0.018) from baseline 4 (3.6-5) to after 3 months 1.6 (1-3.58). **Conclusion(s):** After 3 months of follow-up, UAE improved the distress and symptom of the selected patients with adenomyosis, however larger number of patients and longer duration of follow-up is recommended to support the UAE for the management symptomatic adenomyosis.

P413

Balloon Occlusion of Internal Iliac Arteries in Management of Morbidly Adherent Placenta: Initial Experience at Our Centre

Ummara Siddique, Aman Nawaz Khan, Shahjehan Alam, Syed Ghulam Ghaus, Seema Gul

*Rehman Medical Institute, Peshawar, Pakistan.
E-mail: ummara_81@hotmail.com*

Background: Abnormal placentation is a potential cause of maternal morbidity and mortality from massive postpartum bleeding. Interventional radiology has impressive role in

management of various obstetrical emergencies. Pelvic arterial embolization has added a new dimension in the management of obstetric hemorrhage. **Method(s):** This is a retrospective study conducted at radiology and gynecology departments of Rehman Medical Institute Peshawar. We retrospectively studied eleven cases over a period of two years from May 2015 to May 2017 and patients were diagnosed of having morbidly adherent placenta during antenatal period either by ultrasound or magnetic resonance imaging (MRI). The mean gestational age at presentation was 36 weeks. All patients were in age group of 30-36 yrs with mean age of +33.37 yrs. All were multiparous with a mean parity of +4-5. Previous Cesarean Section delivery was the major independent risk factor in most of the cases. About 60% cases were diagnosed during antenatal checkup and in most of the cases the placenta were type IV Placenta previa. Patients were divided into two groups. 5 patients were managed by conventional treatment (conventional group). In 6 cases, interventional radiologist was involved for trans-catheter arterial balloon occlusion balloon occlusion (study group). The internal iliac balloons were inserted pre-operatively and arterial occlusion was done just after the delivery and in one case additional uterine artery embolization was done post-operatively. In Conventional treatment, no prophylactic temporary balloon occlusion was done and patients were treated by either removal of placenta and oversewing of the placental bed or caesarean hysterectomy. One patient presented postoperatively with placenta in situ and septicemia. **Result(s):** The results of the two groups were compared taking different variables into account. In comparing the operating time the mean operating time in conventional group was +2.37 hr and that in interventional group was +1.25 hr with a total increase of approx.1.12 hr, which is quiet significant. The average blood loss was 962 ml more in conventional group (mean 2037 ml vs 1075 ml) and consequently increased transfusion of blood (mean+6.5 packs vs. 2.25 packs i.e 4.25 packs more). In our comparative study platelets and fresh frozen plasma (FFP) was exclusively needed in conventional group. Mean stay in intensive care unit (ICU) was more in conventional group with a mean of 2.7 days as compared to 1 day in interventional group. In 2 cases there was per-operative injury to surrounding structures attributed to difficult obstetric emergency and excessive blood loss. In one case the patient had ureteric injury and in other case there was bladder injury. In interventional group, not even a single case of damage to surrounding structure was noted, which can be partly attributed to less stress of the surgeon due to comparatively better hemodynamic stability of patient and partly due to clear field of the surgeon. **Conclusion(s):** We conclude that use of per-operative arterial occlusive balloons for managing morbidly adherent placenta at our center showed good initial results with decreased patient morbidity.

Keywords: Intra-arterial occlusive balloons, morbidly adherent placenta, placenta accreta, placenta increta, placenta percreta

P414

Aspiration Thrombectomy for Acute Deep Vein Thrombosis of the Left Lower Extremity of a Child after Cardiac Radiofrequency Ablation

Maria Antonella Ruffino,
Maria Antonella Ruffino, Marco Fronda¹,
Andrea Discalzi, Pierluigi Muratore,
Dorico Righi, Paolo Fonio¹

Department of Diagnostic Imaging and Radiotherapy, Vascular Radiology, A.O.U. Città della Salute e della Scienza di Torino,
¹Department of Surgical Sciences, Radiology Unit, University of Torino, Torino, Italy.
E-mail: ummara_81@hotmail.com

Background: Deep vein thrombosis (DVT) of the lower extremity is rare in children and most of them or history of CVC placement. The therapy includes anticoagulation, systemic thrombolysis and, in rare cases, thrombectomy. Systemic thrombolysis with TPA has been used to treat patients who do not respond to anticoagulation and those with extensive DVT. Thrombo-aspiration with penumbra/Indigo System is a safe and effective treatment for adult patient with ALI, PE, and AVF thrombosis. We report of the case of an adolescent with DVT in the left lower extremity who underwent thrombo-aspiration with Indigo System. **Method(s):** A 14-year-old boy, with previous RFA for wolff-parkinson-white syndrome, with DVT of left popliteal, superficial, common femoral, and iliac veins not responding to anticoagulation underwent thrombo-aspiration with indigo system. After US-guided puncture of the popliteal vein, the dilator (3f) of a micropuncture set kit was positioned. An 0.018 guidewire was advanced along the venous axis, and the dilator was changed for an 8f sheath. Before starting the aspiration, a removable caval filter was positioned. Venography after the aspiration revealed partial recanalization of the left venous axis. Locoregional thrombolysis was performed with 0.6 mg of TPA. There were not complications during any phase of treatment. Anticoagulant therapy was administered. **Result(s):** One week after the procedure, the swelling of the left lower extremity had resolved completely. Caval filter was removed three months later. Duplex us at 1, 3 and 6 months confirmed recanalization of popliteal, common femoral and iliac veins and only residual thrombosis of the superficial femoral vein. The patient had no symptoms. **Conclusion(s):** DVT mechanical aspiration can be safe and effective to treat acute DVT of the lower extremity in older children and adolescents. Thrombo-aspiration before thrombolysis helps to decrease the thrombus burden, improve outflow for effective thrombolysis, and reduce the dose of thrombolytic required.

P415

Parastomal Bleeding Embolization Technique

Sultan R. Alharbi

King Saud University, Riyadh, Saudi Arabia.
E-mail: drsultan000@gmail.com

Background: In this poster presentation we describe and illustrate parastomal bleeding embolization technique in two cases. one case we did direct percutaneous and other case we use trans-hepatic approach. in this poster we also explain why we use direct percutaneous or trans-hepatic approach. Both cases were technically and clinically successful without complication. **Method(s):** N/A. **Result(s):** N/A. **Conclusion(s):** N/A.

P416

Tips and Tricks in Chimney-Graft Technique Endovascular Repair: Steps for Successful Outcome

Samer Koussayer

King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia.
E-mail: sakous@msn.com