Background: ChEVAR becoming very popular option to treat juxtarenal AAA in emergency or urgent situation. The technique is very diverse and this result in conflict data about the outcome and the rate of type 1 endoleak. **Method(s):** In this presentation i will review the steps and tricks in ChEVAR deployment for best outcome and review literature comparing ChEVAR with FEVAR. **Result(s):** Chevar has very good outcome with very low rate of type 1 endoleak if carried out in standard technique. These results are very comparable with FEVAR. **Conclusion(s):** Chevar is an alternative technique to treat juxtarenal AAA especially in emergency situation and it carry very low rate of type 1 endoleak if deployed correctly.

P417

The Effect of Partial Splenic Arterial Embolization in the Control of Hypersplenism

**Mohamed Haroun Hassan Ibrahim,
Shrief Hamed Mostafa Abogamra**

*Ain-Shams University Hospitals, Cairo, Egypt.
E-mail: dr.mharoun@hotmail.com*

Background: The use of splenic artery embolization is used for the intentional infarction of splenic tissues to reduce their consumptive activity. **Method(s):** Fifteen patients with hypersplenism due to chronic liver disease were included. This was a prospective study that included patients with thrombocytopenia (platelet count <70.000) who referred from the Tropical Department to the Interventional Radiology Unit for partial splenic arterial embolization. Partial splenic arterial embolization was performed to reduce the hyperfunction of the spleen in patient with hypersplenism. **Result(s):** The mean age of the selected patients was about 48 years old. The most common cause of hypersplenism is chronic liver disease due to HCV infection in 12 patients (80%), bilharzial disease in 2 patients (13%) and mixed HCV and bilharzial infection in 1 patient (7%). The most frequent risk factors were past history of blood transfusion in 7 patients (47%), past history of surgery in 5 patients (33%), and past history of parenteral antibilharzial treatment in 3 patients (20%). Bleeding tendency was found in 15 patients (100%), while abdominal pain was found in 12 patients (80%). As regard other symptoms, lower limb swelling and abdominal enlargement were noticed in 30% and 20% of patients respectively, anemic manifestations and recurrent infection were noticed in 53% and 47% of patients respectively. Post embolization syndrome was the most common complication and occurred in all patients (100%). Newly developed ascites occurred in 2 patients (13%). Splenic abscess was developed in 2 patients (13%). No other complications had been reported. **Conclusion(s):** Partial splenic arterial embolization is an effective method for the treatment of thrombocytopenia resulting from hypersplenism in patients with liver cirrhosis; however, the procedure is associated with complications, and hence must be performed only if clearly indicated and by an expert person.

P418

My Escape Through Chimney

Yasir Khattak, Ayman Al Sibaie

*Rashid Hospital, Dubai, UAE.
E-mail: dryasirjamil@gmail.com*

Background: Open Repair (OR) remains the “gold standard” for treatment of AAAs and Juxtarenal Aneurysms (JAAA), although there is an increased morbidity and mortality rate, longer operating time, and higher need for transfusions. Endovascular repair (EVAR) offers an alternative as it obviates the need for extensive surgical dissection. Fenestrated EVAR (f-EVAR) devices are used in JAAA to overcome the insufficient neck length resulting in inadequate sealing of standard endografts. Chimney-graft technique EVAR (Ch-EVAR) has been employed to preserve the visceral aortic branches, deploying a stent parallel to the aortic endograft allowing the sealing in a healthier aortic zone. A recent review of the Ch-EVAR showed promising results in terms of morbidity, mortality, and durability at 6 and 12 months follow-up. **Method(s):** We present a unique case of a juxtarenal AAA (J-IAAA) with contained leak in the presence of an indwelling aortic stent with disintegration and kinkage of stent successfully treated with Ch-EVAR. **Result(s):** Patient was successfully treated with EVAR and bilateral renal artery chimney-grafts (Ch-EVAR). **Conclusion(s):** Ch-EVAR shows is valid and viable treatment as well as bail out option for challenging cases of JAAA, where a custom-made device is either not available or not amenable to deployment due to difficult access.

P419

High Flow Arterio-Venous Fistula Closure with Amplatzer Vascular Plug in with High Cardiac Output Failure Patient with CKD

**Amol Madanlal Lahoti, Vivek Ukirde,
Saurabh Anant Joshi, Ashank Bansal,
Arvind Borde**

*LTMCC and LTMGH, Sion Hospital, Mumbai, Maharashtra, India.
E-mail: amollahoti20031987@gmail.com*

Background: The Amplatzer Vascular Plug (AVP) is a cylindrical plug made of self-expanding nitinol wire mesh with precise delivery control, which can be used for a variety of vascular pathologies. An AVP is an ideal vascular occlusion device particularly in high-flow vessels, where there is high risk of migration and systemic embolization with traditional occlusion devices. **Method(s):** 20 year old, young male, recently diagnosed chronic kidney disease presented with palpitation, breathlessness and chest pain. We present a case where through tranfemoral approach, An amplatzer vascular plug is deployed at the site of a -v fistula with precise delivery and complete occlusion of the fistula. **Result(s):** Succesful closure of high flow a-v fistula causing high flow cardiac output failure in a chronic kidney disease patient resulting in improved signs and symptoms with maintained distal circulation. **Conclusion(s):** Vascular occlusion