From Anti-Severe Acute Respiratory Syndrome Coronavirus 2 Immune Response to Cancer Onset via Molecular Mimicry and Cross-Reactivity

Darja Kanduc¹

¹Department of Biosciences, Biotechnologies, and Biopharmaceutics, University of Bari, Bari, Italy

Address for correspondence Darja Kanduc, PhD, Department of Biosciences, Biotechnologies, and Biopharmaceutics, University of Bari, Bari 70121, Italy (e-mail: dkanduc@gmail.com).

Glob Med Genet 2021;8:176–182.

Abstract

Background and Objectives Whether exposure to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) may predispose to the risk of cancer in individuals with no prior cancers is a crucial question that remains unclear. To confirm/refute possible relationships between exposure to the virus and ex novo insurgence of tumors, this study analyzed molecular mimicry and the related cross-reactive potential between SARS-CoV-2 spike glycoprotein (gp) antigen and human tumor-suppressor proteins.

Materials and Methods Tumor-associated proteins were retrieved from UniProt database and analyzed for pentapeptide sharing with SARS-CoV-2 spike gp by using publicly available databases.

Results An impressively high level of molecular mimicry exists between SARS-CoV-2 spike gp and tumor-associated proteins. Numerically, 294 tumor-suppressor proteins share 308 pentapeptides with the viral antigen. Crucially, the shared peptides have a relevant immunologic potential by repeatedly occurring in experimentally validated epitopes. Such immunologic potential is of further relevancy in that most of the shared peptides are also present in infectious pathogens to which, in general, human population has already been exposed, thus indicating the possibility of immunologic imprint phenomena.

Conclusion This article described a vast peptide overlap between SARS-CoV-2 spike gp and tumor-suppressor proteins, and supports autoimmune cross-reactivity as a potential mechanism underlying prospective cancer insurgence following exposure to SARS-CoV-2. Clinically, the findings call for close surveillance of tumor sequelae that possibly could result from the current coronavirus pandemic.

Keywords

- SARS-CoV-2 spike gp
- tumor-suppressor proteins
- molecular mimicry
- cross-reactivity
- long COVID
- cancer epidemic

Introduction

From lung damages to skin diseases and excessive immune responses, the disorders associated with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), that is, coronavirus disease 2019 (COVID-19), are progressively being defined and diagnostically cataloged.^{1–7} Among the many

diseases encompassed by COVID-19, clinical attention has focused on the relationship between SARS-CoV-2 and cancer.^{8–10} Indeed, when compared with the pre-COVID-19 era, COVID-19 pandemic appears to be characterized by higher hospitalization and mortality rates in prostate cancer patients¹¹; increased breast cancer dimensions¹²; increased proportion of patients with advanced non-small cell lung

received

July 16, 2021

accepted after revision

August 2, 2021

published online

September 7, 2021

DOI <https://doi.org/10.1055/s-0041-1735590>.

ISSN 2699-9404.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

cancer¹³; and a higher number of diagnosed head and neck cancers (2.9–8.06% in January–April 2020).¹⁴ Such data have been interpreted as due to the pressure exerted by the viral pandemic on the health care system, so cancer treatments have been delayed and, also, have been related to the viral infection per se.¹⁵ However, how SARS-CoV-2 infection might relate to cancer diseases remains unclear.

According to the research paradigm that peptide sequences common to pathogens and the human host may lead to autoimmunity through cross-reactivity,^{16–21} a previous report²² has proposed cross-reactivity as a likely mechanism that can explain the immunopathology related to SARS-CoV-2 exposure. As a matter of fact, many SARS-CoV-2-derived epitopes were shown to share peptide sequences with human proteins that are involved—when altered, mutated, deficient, and/or improperly functioning—in the etiology of the diseases encompassed by COVID-19.²² Moreover, and of special importance, it was noted that the viral versus human peptide sharing also involved human proteins related to pleuropulmonary blastoma, non-small cell lung cancer, breast invasive ductal carcinoma, multiple human cancers, tumor predisposition syndrome, and mesothelioma, inter alia. That is, the data suggested the possibility that morbidity/mortality increases in various tumors might represent long-term sequelae following exposure to SARS-CoV-2 (Kanduc²² and pertinent references therein).

Hence, this study was undertaken to further explore the relationship between SARS-CoV-2 infection/active immunization and carcinogenesis, and specifically focused on the amino acid (aa) sequence identities between SARS-CoV-2 spike glycoprotein (gp) and tumor-suppressor human proteins. Analyses revealed a vast peptide sharing potentially able to generate pathogenic autoantibodies via cross-reactivity and immunologic imprinting phenomena, thus possibly leading to or enhancing the onset of a wide spectrum of cancer diseases.

Materials and Methods

Peptide sharing between SARS-CoV-2 spike gp (NCBI, GenBank Protein Accession ID = QHD43416.1) and cancer-related human proteins was analyzed using pentapeptides as sequence probes as already described.^{16–22} Pentapeptides were used as minimal immune determinant units since a peptide grouping formed of five aa residues defines an immune unit that can (1) induce highly specific antibodies and (2) determine antigen–antibody-specific interaction (Kanduc^{23,24} and further references therein). Seven hundred eighty-two human proteins (in)directly linked to cancer were obtained from UniProtKB database (www.uniprot.org)²⁵ using “tumor suppressor” as keywords and are listed by UniProt entry in ►Supplementary Table S1 (available in online version only).

Methodologically, the spike gp primary sequence was dissected into pentapeptides offset by one residue (i.e., MFVFL, FVFLV, VFLVL, FLVLL, and so forth) and the resulting viral pentapeptides were analyzed for occurrences within the human proteins related to cancer. The shared peptides were also controlled for occurrences in the pathogens *Bor-*

detella pertussis, *Corynebacterium diphtheriae*, *Clostridium tetani*, *Haemophilus influenzae*, and *Neisseria meningitidis*. The publicly available peptide match and peptide search programs (www.uniprot.org) were used.²⁵

The immunologic potential of the peptides shared between SARS-CoV-2 spike gp and cancer-related proteins was investigated by searching the Immune Epitope Database (IEDB, www.iedb.org)²⁶ for experimentally validated immunoreactive SARS-CoV-2 spike gp-derived epitopes hosting the shared pentapeptides.

Results and Discussion

Searching UniProt database for tumor-suppressor proteins produced 782 protein entries (in)directly related to tumor-suppressor activity and listed in ►Supplementary Table S1 (available in online version only). Of the 782 proteins, 294 have pentapeptides in common with the spike gp, in a total of 308 occurrences (in all, 462, including multiple occurrences). These numbers certify the existence of an impressive, unexpected level of molecular mimicry between the viral antigen and the cancer-related human proteins. Obvious reasons of space prevent a detailed analysis peptide-by-peptide of the peptide overlap that is given in its entirety in ►Supplementary Table S2 (available in online version only). Here in text, a snapshot of the peptide sharing is reported and discussed.

Peptide Sharing between SARS-CoV-2 Spike gp and Tumor-Suppressor Proteins

►Table 1 shows data relative to a representative sample of 19 tumor-suppressor proteins and documents that the peptide commonality with the viral antigen amounts to 29 pentapeptides. From a pathological perspective, ►Table 1 clearly illustrates that even hitting only 19 out of the 294 tumor-suppressor proteins described in ►Supplementary Table S2 (available in online version only) might equate to induce or enhance carcinogenesis in almost all of the human organs, from brain and liver to lung and bones. Examples of the cancers that might be evoked/potentiased by exposure to SARS-CoV-2 in the next future are T cell acute lymphoblastic leukemia, oligodendrogliomas, breast/ovarian cancers, sarcoma, malignant mesothelioma, B cell chronic lymphocytic leukemia, and non-small cell lung carcinoma, among the others.

Immunologic Potential of the Peptide Sharing between SARS-CoV-2 Spike gp and Tumor-Suppressor Proteins

The gloomy outlook hinted at by the findings described in ►Table 1 becomes all the more likely in light of the high immunologic potential of the shared peptides. De facto, investigation of IEDB shows that the 29 pentapeptides shared by the spike gp antigen and the 19 tumor-suppressor proteins (►Table 1) occur and recur in 150 epitopes derived from SARS-CoV-2 that have been experimentally validated and are cataloged as immunoreactive (►Table 2).

In essence, ►Table 2 factually supports the possibility that cross-reactions can be triggered by SARS-CoV-2 infection/active immunization and hit human proteins

Table 1 Peptide sharing between SARS-CoV-2 spike gp and a representative sample of 19 tumor-suppressor proteins

Shared peptides	Tumor-suppressor proteins and related cancer diseases ^a	Refs ^b
DPFLG	<i>BC11B. B cell lymphoma/leukemia 11B. T cell acute lymphoblastic leukemia</i>	27,28
LPPLL, GAGAA, QDVVN, SPDVD	<i>BICRA. Glioma tumor suppressor candidate. Oligodendrogliomas</i>	29
EPQII	<i>BRCA1. Breast cancer type 1 susceptibility protein. Breast/ovarian cancer</i>	30
SLGAE, LAATK, EPVLK	<i>BRCA2. Breast cancer type 2 susceptibility protein. Breast cancer</i>	31
RVVVL	<i>DCC. Netrin receptor DCC. Deleted in colorectal carcinoma. Gallbladder cancer</i>	32
YRVVV, SALGK	<i>DIRA1. GTP-binding protein Di-Ras1. Small GTP-binding tumor suppressor 1. Lost/downregulated in neural tumors</i>	33
ITDAV	<i>EXT1. Exostosin-1. Putative tumor suppressor protein EXT1. Bone tumors</i>	34
ALLAG	<i>EXT2. Exostosin-2. Putative tumor suppressor protein EXT2. Bone tumors</i>	34
TLKSF, RLQSL	<i>IL24. Interleukin-24. Suppression of tumorigenicity 16 protein. Melanoma</i>	35,36
SKPSK	<i>LATS1. Large tumor suppressor homolog 1. Soft tissue sarcoma.</i>	37
ARDLI	<i>LATS2. Large tumor suppressor homolog 2. Malignant mesothelioma</i>	38
YSNNS	<i>MTUS1. Microtubule-associated tumor suppressor 1. Hepatocellular carcinoma</i>	39
GAGAA	<i>PLAT2. Phospholipase A and acyltransferase 2. Gastric cancer</i>	40
GAGAA	<i>PLAT3. Phospholipase A and acyltransferase 3. Ovarian carcinoma cells</i>	41
ADAGE, TYVPA	<i>RBM5. Putative tumor suppressor LUCA15. Lung cancer</i>	42
RDL PQ, NSVAY	<i>SCAI. Suppressor of cancer cell invasion. Downregulated in human tumors</i>	43
LLTDE	<i>SDS3. Suppressor of defective silencing 3 protein homolog. Antitumor activity</i>	44
TQSLL, NFKNL, AGAAA	<i>TASOR. Transcription activation suppressor. Clear cell renal cell carcinoma</i>	45
LSRLD, GDSSS	<i>TRI13. B cell chronic lymphocytic leukemia tumor suppressor Leu5. B cell chronic lymphocytic leukemia. Non-small cell lung carcinoma</i>	46,47

Abbreviations: gp, glycoprotein; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

^aTumor-suppressor proteins given by UniProt entry are in italic.

^bFurther references on cancer diseases are available at UniProt, OMIM, and PubMed.

related to carcinogenesis. Very much this conclusion applies when considering that the extent of the potential immunologic cross-reactivity as well as the spectrum of potentially inducible tumors may be exponentially higher in light of the fact that ▶ **Tables 1** and **2** refer to the peptide commonality involving only a tiny part (19 out of 294) of the human proteins that—if altered—may lead to cancer (see ▶ **Supplementary Table S2** [available in online version only] for the peptide sharing in its totality).

Potential Immunologic Imprint

The 29 pentapeptides common to SARS-CoV-2 spike gp and tumor-suppressor proteins (▶ **Table 1**) are not only present in immunoreactive epitopes (▶ **Table 2**) but, in addition, almost all of them (24 out of 29) are also present in microbial organisms such as *Bordetella pertussis*, *C. diphtheriae*, *C. tetani*, *H. influenzae*, and *N. meningitidis* (▶ **Table 3**). That is, most of the shared peptides are also present in pathogens that an individual possibly encountered during his life because of infections and/or vaccinal routes.

Such interpathogen peptide commonality introduces the immunologic memory as a factor capable of enhancing the extent of the immune cross-reactive response against the tumor-suppressor proteins. That is, as already described

since 1947,^{48,49} the immune system does not induce ex novo primary responses toward a recent infection. Rather, the immune system recalls, amplifies, and intensifies preexisting memory responses toward past infections. In this way, what should have been a primary response to a recent infection is transformed into an anamnestic, secondary, and magnified response to past infections. Simply put, as already discussed in previous reports,^{50–55} the early history of the individual's infections/vaccinations dictates the immune outcomes of any successive infections/vaccinations.

The immunologic imprint phenomenon has its molecular foundations in the massive peptide sharing that characterizes microbial and human proteins^{17,56,57} and of which ▶ **Table 3** is an example. The implications are noteworthy. In the case object, following exposure to SARS-CoV-2 by infection or vaccination, the expected primary response to the virus can turn into a secondary response to previously encountered pathogens against which the immune system already reacted and of which has stored an immunologic memory, that is, the microbial organisms reported in ▶ **Table 3**. However, the previously encountered pathogens are no more present in the human organism, so that the anamnestic immune response triggered by the exposure to SARS-CoV-2 by infection or vaccination ends

Table 2 Immunoreactive SARS-CoV-2 spike gp-derived epitopes containing peptides shared between SARS-CoV-2 spike gp and tumor-suppressor human proteins

IEDB ID ^a	Epitope sequence ^b	IEDB ID ^a	Epitope sequence ^b
36724	litgRLQSL	1329082	ADAGFikqygdcldgia
38831	IQDVVNqnaqalntl	1329083	ADAGFikqygdcldgiaa
51999	qpYRVVVLsf	1329254	demiaqytsALLAG
54725	RLQSLQtyv	1329256	demiaqytsALLAGt
533447	raaeirasanaLAAATK	1329258	demiaqytsALLAGti
1069290	cTLKSFTvekgyiqt	1329260	demiaqytsALLAGtit
1069445	EPQIIIttdntfvsgn	1329323	efqfcnDPFLGvyy
1073938	vqidrlitgRLQSLq	1329325	efqfcnDPFLGvyyh
1074928	ilpdpSKPSK	1329327	efqfcnDPFLGvyyhk
1125063	gltpLPPLL	1329329	efqfcnDPFLGvyyhkn
1309132	nfsqilpdpSKPSKr	1329342	emiaqytsALLAG
1309418	aeirasanaLAATKmsecvlg	1329344	emiaqytsALLAGt
1309447	dfggfnfsqilpdpSKPSKr	1329345	emiaqytsALLAGtit
1309450	dplsetkcTLKSFTvekgyi	1329353	EPQIIIttdntfvsg
1309451	dsfkeeldkyfknhtSPDVD	1329390	fcnDPFLGvyyh
1309467	fdeddsEPVLKgvklhyt	1329414	fqfcnDPFLGvyy
1309478	gNFKNLrefvfkidgyfki	1329416	fqfcnDPFLGvyyh
1309482	gyqpYRVVVLsfellhapat	1329422	fsqilpdpSKPSKr
1309515	lhryltpGDSSSgwttagaa	1329571	idrlitgRLQSLq
1309516	litgRLQSLQtyvtqqlira	1329572	idrlitgRLQSLqt
1309519	lpdpSKPSKrsfiedllfnk	1329595	iqdlsstaSALGKlq
1309523	lssnfgaissvlnidLSRLD	1329597	iraaeirasanaLAAATK
1309532	ngltvLPPLLTDEmiaqyts	1329606	ITDAVdcaldpse
1309534	nitrfqTLALhryltpgd	1329627	khtpinlvRDLQPg
1309546	pflmdlegkqgNFKNLrefv	1329659	IADAGFikqygdcldgiaa
1309556	qfcnDPFLGvyyhknnkswm	1329710	lpdpSKPSKrsfiedllfnkvt
1309561	qrnfyEPQIIIttdntfvsgn	1329762	miaqytsALLAG
1309566	qygdcldgiaARDLlcaqkf	1329764	miaqytsALLAGt
1309567	RDLQPgfsaleplvdlpigi	1329793	ndilSRLDKveaevq
1309585	sssgwtAGAAAYyvgylqpr	1329940	qidrlitgRLQSLqt
1309589	sygfqptngvgypYRVVVL	1329966	qpYRVVVLsfellhapa
1309593	tiTDAVdcaldpsetkctl	1329969	qsiiaytmSLGAE
1309599	TYVPAqeknftapaichdg	1329978	raaeirasanaLAATKm
1309605	vsngthwftqrnfEPQII	1330138	staSALGKIQDVVN
1310254	aeNSVAYSNNsIaip	1330167	tdemiaqytsALLAGt
1310284	ARDLlcaqkfngltv	1330169	tdemiaqytsALLAGti
1310303	caqkfngltvLPPLL	1330171	tdemiaqytsALLAGtit
1310362	eldkyfknhtSPDVD	1330209	TLKSFTvekgyiqt
1310392	fgttldskTQSLLiv	1330210	TLKSFTvekgyiqt
1310415	fngltvLPPLLTDEm	1330211	TLKSFTvekgyiqt
1310448	gklQDVVNqnaqaln	1330219	tpGDSSSgwtAGAAA
1310586	litgRLQSLQtyvtq	1330220	tpinlvRDLQPg
1310609	lpdpSKPSKrsfied	1330305	vqidrlitgRLQSLqt

(Continued)

Table 2 (Continued)

IEDB ID ^a	Epitope sequence ^b	IEDB ID ^a	Epitope sequence ^b
1310611	LPPLLTDEmiaqys	1330306	vqidrlitgRLQSLqtyv
1310747	qpYRVVVLsfellha	1330368	yfkiskhtpinlvRDLQPQ
1310750	qrnfYEPQIIIttdnt	1330391	ytsALLAGtit
1310847	titsgwtfGAGAAIq	1330433	dILSRLD
1310947	wtfGAGAAIqipfam	1330434	dILSRLDppeaevq
1311657	ccscgscckfdeddsEPVLKgvkl	1330437	dsIsstaSALGKI
1311782	pdpSKPSKrsfiedllfnkvltad	1330438	dsIsstaSALGKIq
1312257	cckfdeddsEPVLKg	1330439	dsIsstaSALGKIqdv
1312283	deddsEPVLKgvklh	1330447	EPQIIIttdntfvsngc
1312733	ilpdpSKPSKrsfie	1330456	fsqilpdpSKPSK
1312780	ITDAVdcaldplset	1330457	fsqilpdpSKPSKrs
1313154	miaqysALLAGtit	1330463	gfnfsqilpdpSKPSKr
1313286	pinlvRDLQPQgfwal	1330487	ilpdpSKPSKr
1313756	TLKSFTvek	1330489	iqdsIsstaSALGKI
1313930	vTYVPAqeknftap	1330490	iqdsIsstaSALGKIq
1314170	ADAGFIkqy	1330515	IADAGFIkqy
1315180	aYSNNSiai	1330551	pSKPSKrsf
1316068	etkcTLKSF	1330552	pSKPSKrsfi
1316945	fsqilpdpSKPSKrsfie	1330557	qilpdpSKPSKr
1318209	hvTYVPAqek	1330589	sIsstaSALGKIq
1320443	lgaeNSVAY	1330597	sqilpdpSKPSK
1321084	LPPLLTDEm	1330598	sqilpdpSKPSKr
1323467	qpYRVVVL	1330623	tpinlvRDLQPQgfs
1323750	rasanLAATK	1330624	tpinlvRDLQPQgfsa
1323919	RLQSLqty	1330625	tpinlvRDLQPQgfsalepl
1324353	setkcTLKSF	1331139	cnDPFLGvy
1325536	tlADAGFIk	1332424	itgRLQSLqty
1327824	wtAGAAAYy	1332664	LLTDEmiaqy
1327836	wtfGAGAAI	1334122	TYVPAqeknft
1328800	ytmslgaeNSVAY	1334394	yqpYRVVVL
1328800	ytmslgAEEnsvey	1334452	alhrsyltpGDSSSg
1329076	aaeirasanaLAATK	1334473	NSVAYSNNNSiaiptnft

Abbreviations: gp, glycoprotein; IEDB, Immune Epitope Database; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

^aEpitopes listed according to the IEDB ID number.

^bShared sequences given are capitalized.

to divert onto available immune determinants that, in the present case, are the common determinants present in the tumor-suppressor human proteins. Pathologically, one has to consider that usually an anamnestic secondary immune response is characterized by high avidity and high affinity, besides being quantitatively relevant. Therefore, as a final result, exposure to SARS-CoV-2 by infection and/or vaccination can trigger immediate and violent cross-reactive attacks against the proteins that protect the human being from carcinogenesis.

Conclusion

The findings described in ► **Tables 1 to 3** and ► **Supplementary Table S2** (available in online version only) indicate that molecular mimicry and cross-reactivity between peptides common to SARS-CoV-2 and tumor-related proteins might cause/contribute to cancer epidemics worldwide in the next future. The potential cancer risk might be enhanced by immunologic imprinting phenomena, given the fact that the comparative analyses shown in ► **Table 3** indicate the possibility

Table 3 Occurrences in microbial organisms of pentapeptides common to SARS-CoV-2 spike gp, human proteins related to cancer, and SARS-CoV-2 spike gp-derived epitopes

Organism	Shared peptides
<i>Bordetella pertussis</i>	ADAGF, AGAAA, ALLAG, GAGAA, ITDAV, RLQSL, SLGAE, SPDVD, TYVPA
<i>Clostridium tetani</i>	AGAAA, LAATK, LLTDE, YSNNs
<i>Corynebacterium diphtheriae</i>	AGAAA, ALLAG, EPQII, GAGAA, ITDAV, SALGK, YRVVV
<i>Haemophilus influenzae</i>	AGAAA, GAGAA, LLTDE, LPPLL, LSRLD, NFKNL, NSVAY, RDLPO, RLQSL, RVVVL, SALGK, SLGAE, TLKSF, TQSLL, YSNNs
<i>Neisseria meningitidis</i>	AGAAA, ALLAG, EPVLK, GAGAA, LLTDE, LPPLL

Abbreviations: gp, glycoprotein; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

that a preexisting immune response to previously encountered pathogens could be magnified and intensified following SARS-CoV-2 infection/active immunization. These data are disturbing and invite to immediately intensify clinical surveillance in oncology and to undertake rigid cancer prevention actions, including healthy lifestyle and continuous controls. It will be vital to formulate/implement actions that contemplate fast and safe procedures for clinical trials, development of specific and reliable tumor markers for diagnosis, accurate follow-up of treatments, and, administratively, medical health records, detailed registries, biobanks, health surveys, and coordinated observational studies. Never before do all the recommendations of the European plan for the fight against cancer appear current and necessary.⁵⁸ De facto, tumors appear to be the predominant pathologies that will populate the post pandemic long COVID-19.

Funding

None.

Conflict of Interest

None declared.

References

- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395(10223):497–506
- Sadeghzadeh-Bazargan A, Rezai M, Najar Nobari N, Mozafarpour S, Goodarzi A. Skin manifestations as potential symptoms of diffuse vascular injury in critical COVID-19 patients. *J Cutan Pathol* 2021. Doi: 10.1111/cup.14059
- Boechat JL, Chora I, Morais A, Delgado L. The immune response to SARS-CoV-2 and COVID-19 immunopathology - current perspectives. *Pulmonology* 2021;27(05):423–437
- Cabrera Martimbiano AL, Pacheco RL, Bagattini ÂM, Riera R. Frequency, signs and symptoms, and criteria adopted for long COVID-19: a systematic review. *Int J Clin Pract* 2021:e14357
- Rosen HR, O'Connell C, Nadim MK, et al. Extrapulmonary manifestations of severe acute respiratory syndrome coronavirus-2 infection. *J Med Virol* 2021;93(05):2645–2653
- Ramos-Casals M, Brito-Zerón P, Mariette X. Systemic and organ-specific immune-related manifestations of COVID-19. *Nat Rev Rheumatol* 2021;17(06):315–332
- Suárez-Reyes A, Villegas-Valverde CA. Implications of low-grade inflammation in SARS-CoV-2 immunopathology. *MEDICC Rev* 2021;23(02):42
- Han HJ, Nwagwu C, Anyim O, Ekwereadu C, Kim S. COVID-19 and cancer: from basic mechanisms to vaccine development using nanotechnology. *Int Immunopharmacol* 2021;90:107247
- Corti C, Curigliano G. Commentary: SARS-CoV-2 vaccines and cancer patients. *Ann Oncol* 2021;32(04):569–571
- Xia P, Dubrovskaya A. Tumor markers as an entry for SARS-CoV-2 infection? *FEBS J* 2020;287(17):3677–3680
- Chakravarty D, Ratnani P, Sobotka S, et al. Increased hospitalization and mortality from COVID-19 in prostate cancer patients. *Cancers (Basel)* 2021;13(07):1630
- Vanni G, Pellicciaro M, Materazzo M, et al. Advanced stages and increased need for adjuvant treatments in breast cancer patients: the effect of the one-year COVID-19 pandemic. *Anticancer Res* 2021;41(05):2689–2696
- Park JY, Lee YJ, Kim T, et al. Collateral effects of the coronavirus disease 2019 pandemic on lung cancer diagnosis in Korea. *BMC Cancer* 2020;20(01):1040
- Taylor R, Omakobia E, Sood S, Glore RJ. The impact of coronavirus disease 2019 on head and neck cancer services: a UK tertiary centre study. *J Laryngol Otol* 2020;134(08):684–687
- Derosa L, Melenotte C, Griscelli F, et al. The immuno-oncological challenge of COVID-19. *Nat Can* 2020;1:946–964
- Natale C, Giannini T, Lucchese A, Kanduc D. Computer-assisted analysis of molecular mimicry between human papillomavirus 16 E7 oncoprotein and human protein sequences. *Immunol Cell Biol* 2000;78(06):580–585
- Kanduc D, Stufano A, Lucchese G, Kusalik A. Massive peptide sharing between viral and human proteomes. *Peptides* 2008;29(10):1755–1766
- Lucchese G, Capone G, Kanduc D. Peptide sharing between influenza A H1N1 hemagglutinin and human axon guidance proteins. *Schizophr Bull* 2014;40(02):362–375
- Kanduc D. Peptide cross-reactivity: the original sin of vaccines. *Front Biosci (Schol Ed)* 2012;4:1393–1401
- Lucchese G, Kanduc D. Potential crossreactivity of human immune responses against HCMV glycoprotein B. *Curr Drug Discov Technol* 2016;13(01):16–24
- Lucchese G, Kanduc D. Zika virus and autoimmunity: from microcephaly to Guillain-Barré syndrome, and beyond. *Autoimmun Rev* 2016;15(08):801–808
- Kanduc D. From anti-SARS-CoV-2 immune responses to COVID-19 via molecular mimicry. *Antibodies (Basel)* 2020;9(03):33
- Kanduc D. Pentapeptides as minimal functional units in cell biology and immunology. *Curr Protein Pept Sci* 2013;14(02):111–120
- Kanduc D. Homology, similarity, and identity in peptide epitope immunodefinition. *J Pept Sci* 2012;18(08):487–494
- UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 2019;47(D1):D506–D515
- Vita R, Mahajan S, Overton JA, et al. The Immune Epitope Database (IEDB): 2018 update. *Nucleic Acids Res* 2019;47(D1):D339–D343

- 27 Gutierrez A, Kentsis A, Sanda T, et al. The BCL11B tumor suppressor is mutated across the major molecular subtypes of T-cell acute lymphoblastic leukemia. *Blood* 2011;118(15):4169–4173
- 28 Bartram I, Gökbuget N, Schlee C, et al. Low expression of T-cell transcription factor BCL11b predicts inferior survival in adult standard risk T-cell acute lymphoblastic leukemia patients. *J Hematol Oncol* 2014;7:51
- 29 Yang P, Kollmeyer TM, Buckner K, Bamlet W, Ballman KV, Jenkins RB. Polymorphisms in GLTSCR1 and ERCC2 are associated with the development of oligodendrogliomas. *Cancer* 2005;103(11):2363–2372
- 30 Clark SL, Rodriguez AM, Snyder RR, Hankins GD, Boehning D. Structure-function of the tumor suppressor BRCA1. *Comput Struct Biotechnol J* 2012;1(01):e201204005
- 31 Angeli D, Salvi S, Tedaldi G. Genetic predisposition to breast and ovarian cancers: how many and which genes to test? *Int J Mol Sci* 2020;21(03):1128
- 32 Rai R, Sharma KL, Tiwari S, Misra S, Kumar A, Mittal B. DCC (deleted in colorectal carcinoma) gene variants confer increased susceptibility to gallbladder cancer (Ref. No.: Gene-D-12-01446). *Gene* 2013;518(02):303–309
- 33 Ellis CA, Vos MD, Howell H, Vallecorsa T, Fuets DW, Clark GJ. Rig is a novel Ras-related protein and potential neural tumor suppressor. *Proc Natl Acad Sci U S A* 2002;99(15):9876–9881
- 34 Duncan G, McCormick C, Tufaro F. The link between heparan sulfate and hereditary bone disease: finding a function for the EXT family of putative tumor suppressor proteins. *J Clin Invest* 2001;108(04):511–516
- 35 Huang EY, Madireddi MT, Gopalkrishnan RV, et al. Genomic structure, chromosomal localization and expression profile of a novel melanoma differentiation associated (mda-7) gene with cancer specific growth suppressing and apoptosis inducing properties. *Oncogene* 2001;20(48):7051–7063
- 36 Allen M, Pratscher B, Roka F, et al. Loss of novel mda-7 splice variant (mda-7s) expression is associated with metastatic melanoma. *J Invest Dermatol* 2004;123(03):583–588
- 37 Hisaoka M, Tanaka A, Hashimoto H. Molecular alterations of h-warts/LATS1 tumor suppressor in human soft tissue sarcoma. *Lab Invest* 2002;82(10):1427–1435
- 38 Murakami H, Mizuno T, Taniguchi T, et al. LATS2 is a tumor suppressor gene of malignant mesothelioma. *Cancer Res* 2011;71(03):873–883
- 39 Di Benedetto M, Pineau P, Nouet S, et al. Mutation analysis of the 8p22 candidate tumor suppressor gene ATIP/MTUS1 in hepatocellular carcinoma. *Mol Cell Endocrinol* 2006;252(1-2):207–215
- 40 Liang Y, Zhang C, Dai DQ. Identification of DNA methylation-regulated differentially-expressed genes and related pathways using Illumina 450K BeadChip and bioinformatic analysis in gastric cancer. *Pathol Res Pract* 2019;215(10):152570
- 41 Nazarenko I, Schäfer R, Sers C. Mechanisms of the HRSL3 tumor suppressor function in ovarian carcinoma cells. *J Cell Sci* 2007;120(Pt 8):1393–1404
- 42 Oh JJ, Razfar A, Delgado I, et al. 3p21.3 tumor suppressor gene H37/Luca15/RBM5 inhibits growth of human lung cancer cells through cell cycle arrest and apoptosis. *Cancer Res* 2006;66(07):3419–3427
- 43 Brandt DT, Baarlink C, Kitzing TM, et al. SCAI acts as a suppressor of cancer cell invasion through the transcriptional control of beta1-integrin. *Nat Cell Biol* 2009;11(05):557–568
- 44 Ramakrishna S, Suresh B, Bae SM, Ahn WS, Lim KH, Baek KH. Hyaluronan binding motifs of USP17 and SDS3 exhibit anti-tumor activity. *PLoS One* 2012;7(05):e37772
- 45 Iacobas DA, Mgbemena VE, Iacobas S, Menezes KM, Wang H, Saganti PB. Genomic fabric remodeling in metastatic clear cell renal cell carcinoma (ccRCC): a new paradigm and proposal for a personalized gene therapy approach. *Cancers (Basel)* 2020;12(12):3678
- 46 van Everdink WJ, Baranova A, Lummen C, et al. RFP2, c13ORF1, and FAM10A4 are the most likely tumor suppressor gene candidates for B-cell chronic lymphocytic leukemia. *Cancer Genet Cytogenet* 2003;146(01):48–57
- 47 Xu L, Wu Q, Zhou X, Wu Q, Fang M. TRIM13 inhibited cell proliferation and induced cell apoptosis by regulating NF-κB pathway in non-small-cell lung carcinoma cells. *Gene* 2019;715:144015
- 48 Francis T, Salk JE, Quilligan JJ. Experience with vaccination against influenza in the spring of 1947: a preliminary report. *Am J Public Health Nations Health* 1947;37(08):1013–1016
- 49 Davenport FM, Hennessy AV, Francis T Jr. Epidemiologic and immunologic significance of age distribution of antibody to antigenic variants of influenza virus. *J Exp Med* 1953;98(06):641–656
- 50 Lucchese G, Kanduc D. The Guillain-Barré peptide signatures: from Zika virus to *Campylobacter*, and beyond. *Virus Adaptation and Treatment* 2017;9:1–11
- 51 Lucchese G, Kanduc D. Minimal immune determinants connect Zika virus, human cytomegalovirus, and *Toxoplasma gondii* to microcephaly-related human proteins. *Am J Reprod Immunol* 2017;77(02):e12608
- 52 Kanduc D. Immunobiology: on the inexistence of a negative selection process. *Adv Stud Biol* 2020;12(01):19–28
- 53 Kanduc D. Anti-SARS-CoV-2 Immune response and sudden death: titin as a link. *Adv Stud Biol* 2021;13(01):37–44
- 54 Kanduc D, Shoenfeld Y. Inter-pathogen peptide sharing and the original antigenic sin: solving a paradox. *Open Immunol J* 2018;8:16–27
- 55 Kanduc D. Thromboses and hemostasis disorders associated with coronavirus disease 2019: the possible causal role of cross-reactivity and immunological imprinting. *Glob Med Genet* 2021. Doi: 10.1055/s-0041-1731068
- 56 Trost B, Lucchese G, Stufano A, Bickis M, Kusalik A, Kanduc D. No human protein is exempt from bacterial motifs, not even one. *Self Nonself* 2010;1(04):328–334
- 57 Kanduc D. The comparative biochemistry of viruses and humans: an evolutionary path towards autoimmunity. *Biol Chem* 2019;400(05):629–638
- 58 Europe's Beating Cancer Plan: A new EU approach to prevention, treatment and care. [Internet] Accessed on May 2021 at: https://ec.europa.eu/commission/presscorner/detail/en/ip_21_342